

# Ekeland's variational principle

In mathematical analysis, **Ekeland's variational principle**, discovered by Ivar Ekeland,<sup>[1][2][3]</sup> is a theorem that asserts that there exist nearly optimal solutions to some optimization problems.

Ekeland's variational principle can be used when the lower level set of a minimization problems is not compact, so that the Bolzano–Weierstrass theorem cannot be applied. Ekeland's principle relies on the completeness of the metric space.<sup>[4]</sup>

Ekeland's principle leads to a quick proof of the Caristi fixed point theorem.<sup>[4][5]</sup>

Ekeland's principle has been shown to be equivalent to completeness of metric spaces.<sup>[6]</sup>

Ekeland was associated with the Paris Dauphine University when he proposed this theorem.<sup>[1]</sup>

## Contents

**Ekeland's variational principle**

Preliminary definitions

Statement of the theorem

Corollaries

**References**

**Bibliography**

## Ekeland's variational principle

### Preliminary definitions

Let  $f : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  be a function valued in the extended real numbers  $\mathbb{R} \cup \{-\infty, +\infty\} = [-\infty, +\infty]$ . Then

- $\text{dom } f := \{x \in X : f(x) \neq +\infty\}$  denotes the effective domain of  $f$ .
- $f$  is **proper** if  $\text{dom } f \neq \emptyset$  (that is, if  $f$  is not identically  $+\infty$ ).
- $f$  is **bounded below** if  $\inf_{x \in X} f(x) > -\infty$ .
- given  $x_0 \in X$ , say that  $f$  is *lower semicontinuous* at  $x_0$  if for every real  $y < f(x_0)$  there exists a neighborhood  $U$  of  $x_0$  such that  $f(x) > y$  for all  $x$  in  $U$ .
- $f$  is lower semicontinuous if it is lower semicontinuous at every point of  $X$ .
  - A function is lower semi-continuous if and only if  $\{x \in X : f(x) > y\}$  is an open set for every  $y \in \mathbb{R}$ ; alternatively, a function is lower semicontinuous if and only if all of its lower level sets  $\{x \in X : f(x) \leq y\}$  are closed.

## Statement of the theorem

**Ekeland's variational principle**<sup>[7]</sup> — Let  $(X, d)$  be a complete metric space and  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  a proper (that is, not identically  $+\infty$ ) lower semicontinuous function that is bounded below. Pick  $\epsilon > 0$  and  $x_0 \in X$  such that  $f(x_0) \neq +\infty$  (or equivalently,  $f(x_0) \in \mathbb{R}$ ). There exists some  $v \in X$  such that

$$f(v) \leq f(x_0) - \epsilon d(x_0, v)$$

and for all  $x \neq v$ ,

$$f(v) < f(x) + \epsilon d(v, x).$$

### Proof

Define a function  $G : X \times X \rightarrow \mathbb{R} \cup \{+\infty\}$  by

$$G(x, y) := f(x) + \epsilon d(x, y)$$

which is lower semicontinuous because it is the sum of the lower semicontinuous function  $f$  and the continuous function  $(x, y) \mapsto \epsilon d(x, y)$ . Given  $z \in X$ , define the functions

$$G_z := G(z, \cdot) : X \rightarrow \mathbb{R} \cup \{+\infty\} \quad \text{and} \quad G^z := G(\cdot, z) : X \rightarrow \mathbb{R} \cup \{+\infty\}$$

and define the set

$$F(z) := \{y \in Y : G_z(y) \leq f(z)\} = \{y \in Y : f(z) + \epsilon d(z, y) \leq f(z)\}.$$

It may be verified that for all  $x \in X$ ,

1.  $F(x)$  is closed (because  $G_x := G(x, \cdot) : X \rightarrow \mathbb{R} \cup \{+\infty\}$  is lower semicontinuous);
2. if  $x \notin \text{dom } f$  then  $F(x) = X$ ;
3. if  $x \in \text{dom } f$  then  $x \in F(x) \subseteq \text{dom } f$ ; in particular,  $x_0 \in F(x_0) \subseteq \text{dom } f$ ;
4. if  $y \in F(x)$  then  $F(y) \subseteq F(x)$ .

Let  $s_0 = \inf_{x \in F(x_0)} f(x)$ , which is a real number because  $f$  was assumed to be bounded below.

Pick  $x_1 \in F(x_0)$  such that  $f(x_1) < s_0 + 2^{-1}$ . Having defined  $s_{n-1}$  and  $x_n$ , let

$$s_n := \inf_{x \in F(x_n)} f(x)$$

and pick  $x_{n+1} \in F(x_n)$  such that  $f(x_{n+1}) < s_n + 2^{-(n+1)}$ .

These sequences have the following properties:

- for all  $n \geq 0$ ,  $F(x_{n+1}) \subseteq F(x_n)$  because  $x_{n+1} \in F(x_n)$ , where this now implies that  $s_{n+1} \geq s_n$ ;
- for all  $n \geq 1$ , because  $x_{n+1} \in F(x_n)$

$$\epsilon d(x_{n+1}, x_n) \leq f(x_n) - f(x_{n+1}) \leq f(x_n) - s_n \leq f(x_n) - s_{n-1} < 2^{-n}.$$

It follows that for all  $n, p \geq 1$ ,

$$d(x_{n+1}, x_n) \leq \epsilon 2^{1-n},$$

which proves that  $x_\bullet := (x_n)_{n=0}^\infty$  is a Cauchy sequence. Because  $X$  is a complete metric space, there exists some  $v \in X$  such that  $x_\bullet$  converges to  $v$ . The fact that  $x_m \in F(x_n)$  for all  $m \geq n$  implies that  $v \in \text{cl}_Y F(x_n) = F(x_n)$  for all  $n \geq 0$ , where in particular,  $v \in F(x_0)$ .

The conclusion of the theorem will follow once it is shown that  $F(v) = \{v\}$ . So let  $x \in F(v)$ . Because  $x \in F(x_n)$  for all  $n \geq 0$ , it follows as above that  $\epsilon d(x, x_n) \leq 2^{-n}$ , which implies that  $x_\bullet$  converges to  $x$ . Since the limit of  $x_\bullet$  is unique, it follows that  $x = v$ . Thus  $F(v) = \{v\}$ , as desired. ■

## Corollaries

**Corollary<sup>[8]</sup> — Corollary:** Let  $(X, d)$  be a complete metric space, and let  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be a lower semicontinuous functional on  $X$  that is bounded below and not identically equal to  $+\infty$ . Fix  $\epsilon > 0$  and a point  $x_0 \in X$  such that

$$f(x_0) \leq \epsilon + \inf_{x \in X} f(x).$$

Then, for every  $\lambda > 0$ , there exists a point  $v \in X$  such that

$$f(v) \leq f(x_0),$$

$$d(x_0, v) \leq \lambda,$$

and, for all  $x \neq v$ ,

$$f(x) > f(v) - \frac{\epsilon}{\lambda} d(v, x).$$

A good compromise is to take  $\lambda := \sqrt{\epsilon}$  in the preceding result.<sup>[8]</sup>

## References

1. Ekeland, Ivar (1974). "On the variational principle" (<https://doi.org/10.1016%2F0022-247X%2874%2990025-0>). *J. Math. Anal. Appl.* **47** (2): 324–353. doi:10.1016/0022-247X(74)90025-0 (<https://doi.org/10.1016%2F0022-247X%2874%2990025-0>). ISSN 0022-247X (<https://www.worldcat.org/issn/0022-247X>).
2. Ekeland, Ivar (1979). "Nonconvex minimization problems" (<https://doi.org/10.1090%2FS0273-0979-1979-14595-6>). *Bulletin of the American Mathematical Society. New Series.* **1** (3): 443–474. doi:10.1090/S0273-0979-1979-14595-6 (<https://doi.org/10.1090%2FS0273-0979-1979-14595-6>). MR 0526967 (<https://www.ams.org/mathscinet-getitem?mr=0526967>).
3. Ekeland, Ivar; Temam, Roger (1999). *Convex analysis and variational problems*. Classics in applied mathematics. Vol. 28 (Corrected reprinting of the (1976) North-Holland ed.). Philadelphia, PA: Society for Industrial and Applied Mathematics (SIAM). pp. 357–373. ISBN 0-89871-450-8. MR 1727362 (<https://www.ams.org/mathscinet-getitem?mr=1727362>).
4. Kirk, William A.; Goebel, Kazimierz (1990). *Topics in Metric Fixed Point Theory*. Cambridge University Press. ISBN 0-521-38289-0.
5. Ok, Efe (2007). "D: Continuity I". *Real Analysis with Economic Applications* (<http://homepages.nyu.edu/~eo1/Book-PDF/Ekeland.pdf>) (PDF). Princeton University Press. p. 664. ISBN 978-0-691-11768-3. Retrieved January 31, 2009.
6. Sullivan, Francis (October 1981). "A characterization of complete metric spaces" (<https://doi.org/10.1090%2FS0002-9939-1981-0624927-9>). *Proceedings of the American Mathematical Society.* **83** (2): 345–346. doi:10.1090/S0002-9939-1981-0624927-9 (<https://doi.org/10.1090%2FS0002-9939-1981-0624927-9>). MR 0624927 (<https://www.ams.org/mathscinet-getitem?mr=0624927>).
7. Zalinescu 2002, p. 29.
8. Zalinescu 2002, p. 30.

## Bibliography

- Ekeland, Ivar (1979). "Nonconvex minimization problems" (<https://doi.org/10.1090%2FS0273-0979-1979-14595-6>). *Bulletin of the American Mathematical Society. New Series.* **1** (3): 443–474. doi:10.1090/S0273-0979-1979-14595-6 (<https://doi.org/10.1090%2FS0273-0979-1979-14595-6>). MR 0526967 (<https://www.ams.org/mathscinet-getitem?mr=0526967>).
- Kirk, William A.; Goebel, Kazimierz (1990). *Topics in Metric Fixed Point Theory*. Cambridge University Press. ISBN 0-521-38289-0.
- Zalinescu, C (2002). *Convex analysis in general vector spaces*. River Edge, N.J. London: World Scientific. ISBN 981-238-067-1. OCLC 285163112 (<https://www.worldcat.org/oclc/285163112>).
- Zălinescu, Constantin (30 July 2002). *Convex Analysis in General Vector Spaces* ([https://archive.org/details/convexanalysisge00zali\\_934](https://archive.org/details/convexanalysisge00zali_934)). River Edge, N.J. London: World Scientific Publishing. ISBN 978-981-4488-15-0. MR 1921556 (<https://www.ams.org/mathscinet-getitem?mr=1921556>). OCLC 285163112 (<https://www.worldcat.org/oclc/285163112>) – via Internet Archive.

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Ekeland%27s\\_variational\\_principle&oldid=1072684096](https://en.wikipedia.org/w/index.php?title=Ekeland%27s_variational_principle&oldid=1072684096)"

This page was last edited on 18 February 2022, at 23:54 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.