


Extreme point

In mathematics, an **extreme point** of a convex set S in a real or complex vector space is a point in S which does not lie in any open line segment joining two points of S . In linear programming problems, an extreme point is also called vertex or corner point of S .^[1]



A convex set in light blue, and its extreme points in red.

Contents

Definition

Characterizations

Examples

Properties

Theorems

Krein–Milman theorem

For Banach spaces

Related notions

k -extreme points

See also

Citations

Bibliography

Definition

Throughout, it is assumed that X is a real or complex vector space.

For any $p, x, y \in X$, say that **p lies between**^[2] x and y if $x \neq y$ and there exists a $0 < t < 1$ such that $p = tx + (1 - t)y$.

If K is a subset of X and $p \in K$, then p is called an **extreme point**^[2] of K if it does not lie between any two *distinct* points of K . That is, if there does *not* exist $x, y \in K$ and $0 < t < 1$ such that $x \neq y$ and $p = tx + (1 - t)y$. The set of all extreme points of K is denoted by **extreme(K)**.

Characterizations

The **midpoint**^[2] of two elements x and y in a vector space is the vector $\frac{1}{2}(x + y)$.

For any elements x and y in a vector space, the set $[x, y] = \{tx + (1 - t)y : 0 \leq t \leq 1\}$ is called the **closed line segment** or **closed interval** between x and y . The **open line segment** or **open interval** between x and y is $(x, y) = \emptyset$ when $x = y$ while it is $(x, y) = \{tx + (1 - t)y : 0 < t < 1\}$ when $x \neq y$.^[2] The points x and y are called the **endpoints** of these interval. An interval is said to be a **non-degenerate interval** or a **proper interval** if its endpoints are distinct. The **midpoint of an interval** is the midpoint of its endpoints.

The closed interval $[x, y]$ is equal to the convex hull of (x, y) if (and only if) $x \neq y$. So if K is convex and $x, y \in K$, then $[x, y] \subseteq K$.

If K is a nonempty subset of X and F is a nonempty subset of K , then F is called a **face**^[2] of K if whenever a point $p \in F$ lies between two points of K , then those two points necessarily belong to F .

Theorem^[2] — Let K be a non-empty convex subset of a vector space X and let $p \in K$. Then the following statements are equivalent:

1. p is an extreme point of K .
2. $K \setminus \{p\}$ is convex.
3. p is not the midpoint of a non-degenerate line segment contained in K .
4. for any $x, y \in K$, if $p \in [x, y]$ then $x = p$ or $y = p$.
5. if $x \in X$ is such that both $p + x$ and $p - x$ belong to K , then $x = 0$.
6. $\{p\}$ is a face of K .

Examples

If $a < b$ are two real numbers then a and b are extreme points of the interval $[a, b]$. However, the open interval (a, b) has no extreme points.^[2] Any open interval in \mathbb{R} has no extreme points while any non-degenerate closed interval not equal to \mathbb{R} does have extreme points (that is, the closed interval's endpoint(s)). More generally, any open subset of finite-dimensional Euclidean space \mathbb{R}^n has no extreme points.

The extreme points of the closed unit disk in \mathbb{R}^2 is the unit circle.

The perimeter of any convex polygon in the plane is a face of that polygon.^[2] The vertices of any convex polygon in the plane \mathbb{R}^2 are the extreme points of that polygon.

An injective linear map $F : X \rightarrow Y$ sends the extreme points of a convex set $C \subseteq X$ to the extreme points of the convex set $F(X)$.^[2] This is also true for injective affine maps.

Properties

The extreme points of a compact convex form a Baire space (with the subspace topology) but this set may *fail* to be closed in X .^[2]

Theorems

Krein–Milman theorem

The Krein–Milman theorem is arguably one of the most well-known theorems about extreme points.

Krein–Milman theorem — If S is convex and compact in a locally convex topological vector space, then S is the closed convex hull of its extreme points: In particular, such a set has extreme points.

For Banach spaces

These theorems are for Banach spaces with the Radon–Nikodym property.

A theorem of Joram Lindenstrauss states that, in a Banach space with the Radon–Nikodym property, a nonempty closed and bounded set has an extreme point. (In infinite-dimensional spaces, the property of compactness is stronger than the joint properties of being closed and being bounded).^[3]

Theorem (Gerald Edgar) — Let E be a Banach space with the Radon-Nikodym property, let C be a separable, closed, bounded, convex subset of E , and let a be a point in C . Then there is a probability measure p on the universally measurable sets in C such that a is the barycenter of p , and the set of extreme points of C has p -measure 1.^[4]

Edgar's theorem implies Lindenstrauss's theorem.

Related notions

A closed convex subset of a topological vector space is called strictly convex if every one of its (topological) boundary points is an extreme point.^[5] The unit ball of any Hilbert space is a strictly convex set.^[5]

k -extreme points

More generally, a point in a convex set S is **k -extreme** if it lies in the interior of a k -dimensional convex set within S , but not a $k + 1$ -dimensional convex set within S . Thus, an extreme point is also a **0-extreme** point. If S is a polytope, then the k -extreme points are exactly the interior points of the k -dimensional faces of S . More generally, for any convex set S , the k -extreme points are partitioned into k -dimensional open faces.

The finite-dimensional Krein-Milman theorem, which is due to Minkowski, can be quickly proved using the concept of k -extreme points. If S is closed, bounded, and n -dimensional, and if p is a point in S , then p is k -extreme for some $k \leq n$. The theorem asserts that p is a convex combination of

extreme points. If $k = 0$ then it is immediate. Otherwise p lies on a line segment in S which can be maximally extended (because S is closed and bounded). If the endpoints of the segment are q and r , then their extreme rank must be less than that of p , and the theorem follows by induction.

See also

- [Choquet theory](#)

Citations

- Saltzman, Matthew. "What is the difference between corner points and extreme points in linear programming problems?" (<https://www.quora.com/What-is-the-difference-between-corner-points-and-extreme-points-in-linear-programming-problems>).
- Narici & Beckenstein 2011, pp. 275–339.
- Artstein, Zvi (1980). "Discrete and continuous bang-bang and facial spaces, or: Look for the extreme points". *SIAM Review*. **22** (2): 172–185. doi:10.1137/1022026 (<https://doi.org/10.1137/1022026>). JSTOR 2029960 (<https://www.jstor.org/stable/2029960>). MR 0564562 (<https://www.ams.org/mathscinet-getitem?mr=0564562>).
- Edgar GA. A noncompact Choquet theorem. (<https://www.ams.org/journals/proc/1975-049-02/S0002-9939-1975-0372586-2/S0002-9939-1975-0372586-2.pdf>) Proceedings of the American Mathematical Society. 1975;49(2):354-8.
- Halmos 1982, p. 5.

Bibliography

- Adasch, Norbert; Ernst, Bruno; Keim, Dieter (1978). *Topological Vector Spaces: The Theory Without Convexity Conditions*. Lecture Notes in Mathematics. Vol. 639. Berlin New York: Springer-Verlag. ISBN 978-3-540-08662-8. OCLC 297140003 (<https://www.worldcat.org/oclc/297140003>).
- Bourbaki, Nicolas (1987) [1981]. *Sur certains espaces vectoriels topologiques* (http://www.numdam.org/item?id=AIF_1950__2__5_0) [*Topological Vector Spaces: Chapters 1–5*]. *Annales de l'Institut Fourier*. Éléments de mathématique. Vol. 2. Translated by Eggleston, H.G.; Madan, S. Berlin New York: Springer-Verlag. ISBN 978-3-540-42338-6. OCLC 17499190 (<https://www.worldcat.org/oclc/17499190>).
- Paul E. Black, ed. (2004-12-17). "extreme point" (<https://xlinux.nist.gov/dads/HTML/extremepoint.html>). *Dictionary of algorithms and data structures*. US National institute of standards and technology. Retrieved 2011-03-24.
- Borowski, Ephraim J.; Borwein, Jonathan M. (1989). "extreme point". *Dictionary of mathematics*. Collins dictionary. Harper Collins. ISBN 0-00-434347-6.
- Grothendieck, Alexander (1973). *Topological Vector Spaces* (<https://archive.org/details/topologicalvecto0000grot>). Translated by Chaljub, Orlando. New York: Gordon and Breach Science Publishers. ISBN 978-0-677-30020-7. OCLC 886098 (<https://www.worldcat.org/oclc/886098>).
- Halmos, Paul R. (8 November 1982). *A Hilbert Space Problem Book*. Graduate Texts in Mathematics. Vol. 19 (2nd ed.). New York: Springer-Verlag. ISBN 978-0-387-90685-0. OCLC 8169781 (<https://www.worldcat.org/oclc/8169781>).
- Jarchow, Hans (1981). *Locally convex spaces*. Stuttgart: B.G. Teubner. ISBN 978-3-519-02224-4. OCLC 8210342 (<https://www.worldcat.org/oclc/8210342>).
- Köthe, Gottfried (1983) [1969]. *Topological Vector Spaces I*. Grundlehren der mathematischen Wissenschaften. Vol. 159. Translated by Garling, D.J.H. New York: Springer Science & Business

- Media. ISBN 978-3-642-64988-2. MR 0248498 (<https://www.ams.org/mathscinet-getitem?mr=0248498>). OCLC 840293704 (<https://www.worldcat.org/oclc/840293704>).
- Köthe, Gottfried (1979). *Topological Vector Spaces II*. Grundlehren der mathematischen Wissenschaften. Vol. 237. New York: Springer Science & Business Media. ISBN 978-0-387-90400-9. OCLC 180577972 (<https://www.worldcat.org/oclc/180577972>).
 - Narici, Lawrence; Beckenstein, Edward (2011). *Topological Vector Spaces*. Pure and applied mathematics (Second ed.). Boca Raton, FL: CRC Press. ISBN 978-1584888666. OCLC 144216834 (<https://www.worldcat.org/oclc/144216834>).
 - Robertson, Alex P.; Robertson, Wendy J. (1980). *Topological Vector Spaces*. Cambridge Tracts in Mathematics. Vol. 53. Cambridge England: Cambridge University Press. ISBN 978-0-521-29882-7. OCLC 589250 (<https://www.worldcat.org/oclc/589250>).
 - Rudin, Walter (1991). *Functional Analysis* (<https://archive.org/details/functionalanalys00rudi>). International Series in Pure and Applied Mathematics. Vol. 8 (Second ed.). New York, NY: McGraw-Hill Science/Engineering/Math. ISBN 978-0-07-054236-5. OCLC 21163277 (<https://www.worldcat.org/oclc/21163277>).
 - Schaefer, Helmut H.; Wolff, Manfred P. (1999). *Topological Vector Spaces*. GTM. Vol. 8 (Second ed.). New York, NY: Springer New York Imprint Springer. ISBN 978-1-4612-7155-0. OCLC 840278135 (<https://www.worldcat.org/oclc/840278135>).
 - Schechter, Eric (1996). *Handbook of Analysis and Its Foundations*. San Diego, CA: Academic Press. ISBN 978-0-12-622760-4. OCLC 175294365 (<https://www.worldcat.org/oclc/175294365>).
 - Trèves, François (2006) [1967]. *Topological Vector Spaces, Distributions and Kernels*. Mineola, N.Y.: Dover Publications. ISBN 978-0-486-45352-1. OCLC 853623322 (<https://www.worldcat.org/oclc/853623322>).
 - Wilansky, Albert (2013). *Modern Methods in Topological Vector Spaces*. Mineola, New York: Dover Publications, Inc. ISBN 978-0-486-49353-4. OCLC 849801114 (<https://www.worldcat.org/oclc/849801114>).

Retrieved from "https://en.wikipedia.org/w/index.php?title=Extreme_point&oldid=1068413394"

This page was last edited on 28 January 2022, at 08:32 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.