


# Carathéodory's theorem (convex hull)

**Carathéodory's theorem** is a theorem in convex geometry. It states that if a point  $x$  of  $\mathbf{R}^d$  lies in the convex hull of a set  $P$ , then  $x$  can be written as the convex combination of at most  $d + 1$  points in  $P$ . Namely, there is a subset  $P'$  of  $P$  consisting of  $d + 1$  or fewer points such that  $x$  lies in the convex hull of  $P'$ . Equivalently,  $x$  lies in an  $r$ -simplex with vertices in  $P$ , where  $r \leq d$ . The smallest  $r$  that makes the last statement valid for each  $x$  in the convex hull of  $P$  is defined as the *Carathéodory's number* of  $P$ . Depending on the properties of  $P$ , upper bounds lower than the one provided by Carathéodory's theorem can be obtained.<sup>[1]</sup> Note that  $P$  need not be itself convex. A consequence of this is that  $P'$  can always be extremal in  $P$ , as non-extremal points can be removed from  $P$  without changing the membership of  $x$  in the convex hull.

The similar theorems of Helly and Radon are closely related to Carathéodory's theorem: the latter theorem can be used to prove the former theorems and vice versa.<sup>[2]</sup>

The result is named for Constantin Carathéodory, who proved the theorem in 1911 for the case when  $P$  is compact.<sup>[3]</sup> In 1914 Ernst Steinitz expanded Carathéodory's theorem for any sets  $P$  in  $\mathbf{R}^d$ .<sup>[4]</sup>



An illustration of Carathéodory's theorem for a square in  $\mathbf{R}^2$

## Contents

Example

Proof

Variants

Carathéodory's theorem for the conical hull

Dimensionless variant

Colorful Carathéodory theorem

See also

Notes

Further reading

External links

## Example

Consider a set  $P = \{(0,0), (0,1), (1,0), (1,1)\}$  which is a subset of  $\mathbf{R}^2$ . The convex hull of this set is a square. Consider now a point  $x = (1/4, 1/4)$ , which is in the convex hull of  $P$ . We can then construct a set  $\{(0,0), (0,1), (1,0)\} = P'$ , the convex hull of which is a triangle and encloses  $x$ , and thus the theorem works for this instance, since  $|P'| = 3$ . It may help to visualise Carathéodory's theorem in 2 dimensions, as saying that we can construct a triangle consisting of points from  $P$  that encloses any point in  $P$ .

## Proof

Let  $\mathbf{x}$  be a point in the convex hull of  $P$ . Then,  $\mathbf{x}$  is a convex combination of a finite number of points in  $P$ :

$$\mathbf{x} = \sum_{j=1}^k \lambda_j \mathbf{x}_j$$

where every  $\mathbf{x}_j$  is in  $P$ , every  $\lambda_j$  is (w.l.o.g.) positive, and  $\sum_{j=1}^k \lambda_j = 1$ .

Suppose  $k > d + 1$  (otherwise, there is nothing to prove). Then, the vectors  $\mathbf{x}_2 - \mathbf{x}_1, \dots, \mathbf{x}_k - \mathbf{x}_1$  are linearly dependent,

so there are real scalars  $\mu_2, \dots, \mu_k$ , not all zero, such that

$$\sum_{j=2}^k \mu_j (\mathbf{x}_j - \mathbf{x}_1) = \mathbf{0}.$$

If  $\mu_1$  is defined as

$$\mu_1 := - \sum_{j=2}^k \mu_j$$

then

$$\begin{aligned} \sum_{j=1}^k \mu_j \mathbf{x}_j &= \mathbf{0} \\ \sum_{j=1}^k \mu_j &= 0 \end{aligned}$$

and not all of the  $\mu_j$  are equal to zero. Therefore, at least one  $\mu_j > 0$ . Then,

$$\mathbf{x} = \sum_{j=1}^k \lambda_j \mathbf{x}_j - \alpha \sum_{j=1}^k \mu_j \mathbf{x}_j = \sum_{j=1}^k (\lambda_j - \alpha \mu_j) \mathbf{x}_j$$

for any real  $\alpha$ . In particular, the equality will hold if  $\alpha$  is defined as

$$\alpha := \min_{1 \leq j \leq k} \left\{ \frac{\lambda_j}{\mu_j} : \mu_j > 0 \right\} = \frac{\lambda_i}{\mu_i}.$$

Note that  $\alpha > 0$ , and for every  $j$  between 1 and  $k$ ,

$$\lambda_j - \alpha \mu_j \geq 0.$$

In particular,  $\lambda_i - \alpha \mu_i = 0$  by definition of  $\alpha$ . Therefore,

$$\mathbf{x} = \sum_{j=1}^k (\lambda_j - \alpha \mu_j) \mathbf{x}_j$$

where every  $\lambda_j - \alpha \mu_j$  is nonnegative, their sum is one, and furthermore,  $\lambda_i - \alpha \mu_i = 0$ . In other words,  $\mathbf{x}$  is represented as a convex combination of at most  $k-1$  points of  $P$ . This process can be repeated until  $\mathbf{x}$  is represented as a convex combination of at most  $d + 1$  points in  $P$ .

Alternative proofs use Helly's theorem or the Perron–Frobenius theorem.<sup>[5][6]</sup>

## Variants

---

### Carathéodory's theorem for the conical hull

If a point  $x$  of  $\mathbf{R}^d$  lies in the **conical hull** of a set  $P$ , then  $x$  can be written as the **conical combination** of at most  $d$  points in  $P$ . Namely, there is a subset  $P'$  of  $P$  consisting of  $d$  or fewer points, such that  $x$  lies in the conical hull of  $P'$ .<sup>[7]:257</sup> The proof is similar to the original theorem; the difference is that, in a  $d$ -dimensional space, the maximum size of a linearly-independent set is  $d$ , while the maximum size of an affinely-independent set is  $d+1$ .<sup>[8]</sup>

### Dimensionless variant

Recently, Adiprasito, Barany, Mustafa and Terpai proved a variant of Carathéodory's theorem that does not depend on the dimension of the space.<sup>[9]</sup>

### Colorful Carathéodory theorem

Let  $X_1, \dots, X_{d+1}$  be sets in  $\mathbf{R}^d$  and let  $x$  be a point contained in the intersection of the convex hulls of all these  $d+1$  sets.

Then there is a set  $T = \{x_1, \dots, x_{d+1}\}$ , where  $x_1 \in X_1, \dots, x_{d+1} \in X_{d+1}$ , such that the convex hull of  $T$  contains the point  $x$ .<sup>[10]</sup>

By viewing the sets  $X_1, \dots, X_{d+1}$  as different colors, the set  $T$  is made by points of all colors, hence the "colorful" in the theorem's name.<sup>[11]</sup> The set  $T$  is also called a *rainbow simplex*, since it is a  $d$ -dimensional simplex in which each corner has a different color.<sup>[12]</sup>

This theorem has a variant in which the convex hull is replaced by the conical hull.<sup>[10]</sup>:Thm.2.2 Let  $X_1, \dots, X_d$  be sets in  $\mathbf{R}^d$  and let  $x$  be a point contained in the intersection of the *conical hulls* of all these  $d$  sets. Then there is a set  $T = \{x_1, \dots, x_d\}$ , where  $x_1 \in X_1, \dots, x_d \in X_d$ , such that the *conical hull* of  $T$  contains the point  $x$ .<sup>[10]</sup>

Mustafa and Ray extended this colorful theorem from points to convex bodies.<sup>[12]</sup>

## See also

- Shapley–Folkman lemma
- Helly's theorem
- Kirchberger's theorem
- Radon's theorem, and its generalization Tverberg's theorem
- Krein–Milman theorem
- Choquet theory

## Notes

1. Bárány, Imre; Karasev, Roman (2012-07-20). "Notes About the Carathéodory Number". *Discrete & Computational Geometry*. **48** (3): 783–792. arXiv:1112.5942 (<https://arxiv.org/abs/1112.5942>). doi:10.1007/s00454-012-9439-z (<https://doi.org/10.1007%2Fs00454-012-9439-z>). ISSN 0179-5376 (<https://www.worldcat.org/issn/0179-5376>). S2CID 9090617 (<https://api.semanticscholar.org/CorpusID:9090617>).
2. Danzer, L.; Grünbaum, B.; Klee, V. (1963). "Helly's theorem and its relatives". *Convexity*. Proc. Symp. Pure Math. Vol. 7. American Mathematical Society. pp. 101–179. See in particular p.109
3. Carathéodory, C. (1911). "Über den Variabilitätsbereich der Fourier'schen Konstanten von positiven harmonischen Funktionen" (<https://link.springer.com/article/10.1007%2FBF03014795>). *Rendiconti del Circolo Matematico di Palermo (1884–1940)* (in German). **32** (1): 193–217[see p.200 bottom]. doi:10.1007/BF03014795 (<https://doi.org/10.1007%2FBF03014795>). S2CID 120032616 (<https://api.semanticscholar.org/CorpusID:120032616>).
4. Steinitz, Ernst (1913). "Bedingt konvergente Reihen und konvexe Systeme". *J. Reine Angew. Math.* **1913** (143): 128–175. doi:10.1515/crll.1913.143.128 (<https://doi.org/10.1515%2Fcrl.1913.143.128>). S2CID 120411668 (<https://api.semanticscholar.org/CorpusID:120411668>).
5. Eggleston, H. G. (1958). *Convexity* (<http://ebooks.cambridge.org/ref/id/CBO9780511566172>). Cambridge University Press. doi:10.1017/cbo9780511566172 (<https://doi.org/10.1017%2Fcbo9780511566172>). ISBN 9780511566172. See pages 40–41.
6. Naszódi, Márton; Polyanskii, Alexandr (2021). "Perron and Frobenius Meet Carathéodory". *The Electronic Journal of Combinatorics*. **28** (3). arXiv:1901.00540 (<https://arxiv.org/abs/1901.00540>). doi:10.37236/9996 (<https://doi.org/10.37236%2F9996>). S2CID 119656227 (<https://api.semanticscholar.org/CorpusID:119656227>).
7. Lovász, László; Plummer, M. D. (1986). *Matching Theory*. Annals of Discrete Mathematics. Vol. 29. North-Holland. ISBN 0-444-87916-1. MR 0859549 (<https://www.ams.org/mathscinet-getitem?mr=0859549>).
8. "convex geometry - Caratheodory's theorem for vectors in a cone" (<https://math.stackexchange.com/questions/3759861/caratheodorys-theorem-for-vectors-in-a-cone>). *Mathematics Stack Exchange*. Retrieved 2020-07-22.

9. Adiprasito, Karim; Bárány, Imre; Mustafa, Nabil H.; Terpai, Tamás (2019-08-28). "Theorems of Carathéodory, Helly, and Tverberg without dimension". *arXiv:1806.08725* (<https://arxiv.org/abs/1806.08725>) [[math.MG](https://arxiv.org/archive/math)] (<https://arxiv.org/archive/math>).
10. Bárány, Imre (1982-01-01). "A generalization of carathéodory's theorem" (<https://doi.org/10.1016%2F0012-365X%2882%2990115-7>). *Discrete Mathematics*. **40** (2–3): 141–152. doi:10.1016/0012-365X(82)90115-7 (<https://doi.org/10.1016%2F0012-365X%2882%2990115-7>). ISSN 0012-365X (<https://www.worldcat.org/issn/0012-365X>).
11. Montejano, Luis; Fabila, Ruy; Bracho, Javier; Bárány, Imre; Arocha, Jorge L. (2009-09-01). "Very Colorful Theorems" (<https://doi.org/10.1007%2Fs00454-009-9180-4>). *Discrete & Computational Geometry*. **42** (2): 142–154. doi:10.1007/s00454-009-9180-4 (<https://doi.org/10.1007%2Fs00454-009-9180-4>). ISSN 1432-0444 (<https://www.worldcat.org/issn/1432-0444>).
12. Mustafa, Nabil H.; Ray, Saurabh (2016-04-06). "An optimal generalization of the Colorful Carathéodory theorem" (<https://doi.org/10.1016%2Fj.disc.2015.11.019>). *Discrete Mathematics*. **339** (4): 1300–1305. doi:10.1016/j.disc.2015.11.019 (<https://doi.org/10.1016%2Fj.disc.2015.11.019>). ISSN 0012-365X (<https://www.worldcat.org/issn/0012-365X>).

## Further reading

- Eckhoff, J. (1993). "Helly, Radon, and Carathéodory type theorems". *Handbook of Convex Geometry*. Vol. A, B. Amsterdam: North-Holland. pp. 389–448.
- Mustafa, Nabil; Meunier, Frédéric; Goaoc, Xavier; De Loera, Jesús (2019). "The discrete yet ubiquitous theorems of Carathéodory, Helly, Sperner, Tucker, and Tverberg". *Bulletin of the American Mathematical Society*. **56** (3): 415–511. *arXiv:1706.05975* (<https://arxiv.org/abs/1706.05975>). doi:10.1090/bull/1653 (<https://doi.org/10.1090%2Fbull%2F1653>). ISSN 0273-0979 (<https://www.worldcat.org/issn/0273-0979>). S2CID 32071768 (<https://api.semanticscholar.org/CorpusID:32071768>).

## External links

- Concise statement of theorem (<https://planetmath.org/caratheodorystheorem>) in terms of convex hulls (at [PlanetMath](#))

Retrieved from "https://en.wikipedia.org/w/index.php?title=Carathéodory%27s\_theorem\_(convex\_hull)&oldid=1068611023"

This page was last edited on 29 January 2022, at 11:31 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.