

WIKIPEDIA

Busemann's theorem

In mathematics, **Busemann's theorem** is a theorem in Euclidean geometry and geometric tomography. It was first proved by Herbert Busemann in 1949 and was motivated by his theory of area in Finsler spaces.

Statement of the theorem

Let K be a convex body in n -dimensional Euclidean space \mathbf{R}^n containing the origin in its interior. Let S be an $(n - 2)$ -dimensional linear subspace of \mathbf{R}^n . For each unit vector θ in S^\perp , the orthogonal complement of S , let S_θ denote the $(n - 1)$ -dimensional hyperplane containing θ and S . Define $r(\theta)$ to be the $(n - 1)$ -dimensional volume of $K \cap S_\theta$. Let C be the curve $\{\theta r(\theta)\}$ in S^\perp . Then C forms the boundary of a convex body in S^\perp .

See also

- Brunn–Minkowski inequality
- Prékopa–Leindler inequality

References

- Busemann, Herbert (1949). "A theorem on convex bodies of the Brunn-Minkowski type" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1062952>). *Proc. Natl. Acad. Sci. U.S.A.* **35** (1): 27–31. doi:10.1073/pnas.35.1.27 (<https://doi.org/10.1073/pnas.35.1.27>). PMC 1062952 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1062952>). PMID 16588849 (<https://pubmed.ncbi.nlm.nih.gov/16588849>).
- Gardner, Richard J. (2002). "The Brunn-Minkowski inequality". *Bull. Amer. Math. Soc. (N.S.)*. **39** (3): 355–405 (electronic). CiteSeerX 10.1.1.106.7344 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.106.7344>). doi:10.1090/S0273-0979-02-00941-2 (<https://doi.org/10.1090/S0273-0979-02-00941-2>).

Retrieved from "https://en.wikipedia.org/w/index.php?title=Busemann%27s_theorem&oldid=1040065763"

This page was last edited on 22 August 2021, at 12:56 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.