

Convex set

In geometry, a subset of a Euclidean space, or more generally an affine space over the reals, is **convex** if, given any two points in the subset, the subset contains the whole line segment that joins them. Equivalently, a **convex set** or a **convex region** is a subset that intersects every line into a single line segment (possibly empty).^{[1][2]} For example, a solid cube is a convex set, but anything that is hollow or has an indent, for example, a crescent shape, is not convex.

The boundary of a convex set is always a convex curve. The intersection of all the convex sets that contain a given subset A of Euclidean space is called the convex hull of A . It is the smallest convex set containing A .

A convex function is a real-valued function defined on an interval with the property that its epigraph (the set of points on or above the graph of the function) is a convex set. Convex minimization is a subfield of optimization that studies the problem of minimizing convex functions over convex sets. The branch of mathematics devoted to the study of properties of convex sets and convex functions is called convex analysis.

The notion of a convex set can be generalized as described below.




Illustration of a convex set which looks somewhat like a deformed circle. The line segment, illustrated in black above, joining points x and y , lies completely within the set, illustrated in green. Since this is true for any potential locations of any two points within the above set, the set is convex.




Illustration of a non-convex set. Illustrated by the above line segment whereby it changes from the black color to the red color. Exemplifying why this above set, illustrated in green, is non-convex.

Contents

Definitions

Non-convex set

Properties

Intersections and unions

Closed convex sets

Convex sets and rectangles

Blaschke-Santaló diagrams

Other properties

Convex hulls and Minkowski sums

Convex hulls

Minkowski addition

Convex hulls of Minkowski sums

Minkowski sums of convex sets

Generalizations and extensions for convexity

Star-convex (star-shaped) sets

[Orthogonal convexity](#)

[Non-Euclidean geometry](#)

[Order topology](#)

[Convexity spaces](#)

See also

References

External links

Definitions

Let S be a [vector space](#) or an [affine space](#) over the real numbers, or, more generally, over some [ordered field](#). This includes Euclidean spaces, which are [affine spaces](#). A subset C of S is **convex** if, for all x and y in C , the [line segment](#) connecting x and y is included in C . This means that the [affine combination](#) $(1 - t)x + ty$ belongs to C , for all x and y in C , and t in the [interval](#) $[0, 1]$. This implies that convexity (the property of being convex) is invariant under [affine transformations](#). This implies also that a convex set in a [real](#) or [complex topological vector space](#) is [path-connected](#), thus [connected](#).

A set C is *strictly convex* if every point on the line segment connecting x and y other than the endpoints is inside the [topological interior](#) of C . A closed convex subset is strictly convex if and only if every one of its [boundary points](#) is an [extreme point](#).^[3]

A set C is *absolutely convex* if it is convex and [balanced](#).


The convex [subsets](#) of \mathbf{R} (the set of real numbers) are the intervals and the points of \mathbf{R} . Some examples of convex subsets of the [Euclidean plane](#) are solid regular polygons, solid triangles, and intersections of solid triangles. Some examples of convex subsets of a [Euclidean 3-dimensional space](#) are the [Archimedean solids](#) and the [Platonic solids](#). The [Kepler-Poinsot polyhedra](#) are examples of non-convex sets.

Non-convex set

A set that is not convex is called a *non-convex set*. A [polygon](#) that is not a [convex polygon](#) is sometimes called a [concave polygon](#),^[4] and some sources more generally use the term *concave set* to mean a non-convex set,^[5] but most authorities prohibit this usage.^{[6][7]}

The [complement](#) of a convex set, such as the [epigraph](#) of a [concave function](#), is sometimes called a *reverse convex set*, especially in the context of [mathematical optimization](#).^[8]

Properties



A [function](#) is convex if and only if its [epigraph](#), the region (in green) above its [graph](#) (in blue), is a convex set.

Given r points u_1, \dots, u_r in a convex set S , and r nonnegative numbers $\lambda_1, \dots, \lambda_r$ such that $\lambda_1 + \dots + \lambda_r = 1$, the affine combination

$$\sum_{k=1}^r \lambda_k u_k$$

belongs to S . As the definition of a convex set is the case $r = 2$, this property characterizes convex sets.

Such an affine combination is called a convex combination of u_1, \dots, u_r .

Intersections and unions

The collection of convex subsets of a vector space, an affine space, or a Euclidean space has the following properties:^{[9][10]}

1. The empty set and the whole space are convex.
2. The intersection of any collection of convex sets is convex.
3. The union of a sequence of convex sets is convex, if they form a non-decreasing chain for inclusion. For this property, the restriction to chains is important, as the union of two convex sets need not be convex.

Closed convex sets

Closed convex sets are convex sets that contain all their limit points. They can be characterised as the intersections of closed half-spaces (sets of point in space that lie on and to one side of a hyperplane).

From what has just been said, it is clear that such intersections are convex, and they will also be closed sets. To prove the converse, i.e., every closed convex set may be represented as such intersection, one needs the supporting hyperplane theorem in the form that for a given closed convex set C and point P outside it, there is a closed half-space H that contains C and not P . The supporting hyperplane theorem is a special case of the Hahn–Banach theorem of functional analysis.

Convex sets and rectangles

Let C be a convex body in the plane (a convex set whose interior is non-empty). We can inscribe a rectangle r in C such that a homothetic copy R of r is circumscribed about C . The positive homothety ratio is at most 2 and:^[11]

$$\frac{1}{2} \cdot \text{Area}(R) \leq \text{Area}(C) \leq 2 \cdot \text{Area}(r)$$

Blaschke-Santaló diagrams

The set \mathcal{K}^2 of all planar convex bodies can be parameterized in terms of the convex body diameter D , its inradius r (the biggest circle contained in the convex body) and its circumradius R (the smallest circle containing the convex body). In fact, this set can be described by the set of inequalities given by^{[12][13]}


$$2r \leq D \leq 2R$$

$$R \leq \frac{\sqrt{3}}{3} D$$

$$r + R \leq D$$

$$D^2 \sqrt{4R^2 - D^2} \leq 2R(2R + \sqrt{4R^2 - D^2})$$

and can be visualized as the image of the function g that maps a convex body to the \mathbf{R}^2 point given by $(r/R, D/2R)$. The image of this function is known a (r, D, R) Blaschke-Santaló diagram.^[13]



Blaschke-Santaló (r, D, R) diagram for planar convex bodies. \mathbf{L} denotes the line segment, $\mathbf{I}_{\frac{\pi}{3}}$ the equilateral triangle, \mathbf{RT} the Reuleaux triangle and \mathbf{B}_2 the unit circle.

Alternatively, the set \mathcal{K}^2 can also be parametrized by its width (the smallest distance between any two different parallel support hyperplanes), perimeter and area.^{[12][13]}

Other properties

Let X be a topological vector space and $C \subseteq X$ be convex.

- $\text{Cl } C$ and $\text{Int } C$ are both convex (i.e. the closure and interior of convex sets are convex).
- If $a \in \text{Int } C$ and $b \in \text{Cl } C$ then $[a, b] \subseteq \text{Int } C$ (where $[a, b] := \{(1 - r)a + rb : 0 \leq r < 1\}$).
- If $\text{Int } C \neq \emptyset$ then:
 - $\text{cl}(\text{Int } C) = \text{Cl } C$, and

- $\text{Int } C = \text{Int}(\text{Cl } C) = C^i$, where C^i is the algebraic interior of C .

Convex hulls and Minkowski sums

Convex hulls

Every subset A of the vector space is contained within a smallest convex set (called the convex hull of A), namely the intersection of all convex sets containing A . The convex-hull operator $\text{Conv}()$ has the characteristic properties of a hull operator:

- *extensive*: $S \subseteq \text{Conv}(S)$,
- *non-decreasing*: $S \subseteq T$ implies that $\text{Conv}(S) \subseteq \text{Conv}(T)$, and
- *idempotent*: $\text{Conv}(\text{Conv}(S)) = \text{Conv}(S)$.

The convex-hull operation is needed for the set of convex sets to form a lattice, in which the "join" operation is the convex hull of the union of two convex sets

$$\text{Conv}(S) \vee \text{Conv}(T) = \text{Conv}(S \cup T) = \text{Conv}(\text{Conv}(S) \cup \text{Conv}(T)).$$

The intersection of any collection of convex sets is itself convex, so the convex subsets of a (real or complex) vector space form a complete lattice.


Minkowski addition

In a real vector-space, the Minkowski sum of two (non-empty) sets, S_1 and S_2 , is defined to be the set $S_1 + S_2$ formed by the addition of vectors element-wise from the summand-sets

$$S_1 + S_2 = \{x_1 + x_2 : x_1 \in S_1, x_2 \in S_2\}.$$

More generally, the Minkowski sum of a finite family of (non-empty) sets S_n is the set formed by element-wise addition of vectors

$$\sum_n S_n = \left\{ \sum_n x_n : x_n \in S_n \right\}.$$



Minkowski addition of sets. The sum of the squares $Q_1=[0,1]^2$ and $Q_2=[1,2]^2$ is the square $Q_1+Q_2=[1,3]^2$.

For Minkowski addition, the *zero set* $\{0\}$ containing only the zero vector 0 has special importance: For every non-empty subset S of a vector space

$$S + \{0\} = S;$$

in algebraic terminology, $\{0\}$ is the identity element of Minkowski addition (on the collection of non-empty sets).^[14]

Convex hulls of Minkowski sums

Minkowski addition behaves well with respect to the operation of taking convex hulls, as shown by the following proposition:

Let S_1, S_2 be subsets of a real vector-space, the convex hull of their Minkowski sum is the Minkowski sum of their convex hulls

$$\text{Conv}(S_1 + S_2) = \text{Conv}(S_1) + \text{Conv}(S_2).$$

This result holds more generally for each finite collection of non-empty sets:

$$\text{Conv} \left(\sum_n S_n \right) = \sum_n \text{Conv} (S_n).$$

In mathematical terminology, the operations of Minkowski summation and of forming convex hulls are commuting operations.^{[15][16]}

Minkowski sums of convex sets

The Minkowski sum of two compact convex sets is compact. The sum of a compact convex set and a closed convex set is closed.^[17]

The following famous theorem, proved by Dieudonné in 1966, gives a sufficient condition for the difference of two closed convex subsets to be closed.^[18] It uses the concept of a **recession cone** of a non-empty convex subset S , defined as:

$$\text{rec } S = \{x \in X : x + S \subseteq S\},$$

where this set is a convex cone containing $0 \in X$ and satisfying $S + \text{rec } S = S$. Note that if S is closed and convex then $\text{rec } S$ is closed and for all $s_0 \in S$,

$$\text{rec } S = \bigcap_{t>0} t(S - s_0).$$

Theorem (Dieudonné). Let A and B be non-empty, closed, and convex subsets of a locally convex topological vector space such that $\text{rec } A \cap \text{rec } B$ is a linear subspace. If A or B is locally compact then $A - B$ is closed.

Generalizations and extensions for convexity

The notion of convexity in the Euclidean space may be generalized by modifying the definition in some or other aspects. The common name "generalized convexity" is used, because the resulting objects retain certain properties of convex sets.

Star-convex (star-shaped) sets

Let C be a set in a real or complex vector space. C is **star convex (star-shaped)** if there exists an x_0 in C such that the line segment from x_0 to any point y in C is contained in C . Hence a non-empty convex set is always star-convex but a star-convex set is not always convex.

Orthogonal convexity

An example of generalized convexity is **orthogonal convexity**.^[19]

A set S in the Euclidean space is called **orthogonally convex** or **ortho-convex**, if any segment parallel to any of the coordinate axes connecting two points of S lies totally within S . It is easy to prove that an intersection of any collection of orthoconvex sets is orthoconvex. Some other properties of convex sets are valid as well.

Non-Euclidean geometry

The definition of a convex set and a convex hull extends naturally to geometries which are not Euclidean by defining a geodesically convex set to be one that contains the geodesics joining any two points in the set.

Order topology

Convexity can be extended for a totally ordered set X endowed with the order topology.^[20]

Let $Y \subseteq X$. The subspace Y is a convex set if for each pair of points a, b in Y such that $a \leq b$, the interval $[a, b] = \{x \in X \mid a \leq x \leq b\}$ is contained in Y . That is, Y is convex if and only if for all a, b in Y , $a \leq b$ implies $[a, b] \subseteq Y$.

A convex set is **not** connected in general: a counter-example is given by the subspace $\{1,2,3\}$ in \mathbf{Z} , which is both convex and not connected.

Convexity spaces

The notion of convexity may be generalised to other objects, if certain properties of convexity are selected as axioms.

Given a set X , a **convexity** over X is a collection \mathcal{C} of subsets of X satisfying the following axioms:^{[9][10][21]}

1. The empty set and X are in \mathcal{C}
2. The intersection of any collection from \mathcal{C} is in \mathcal{C} .
3. The union of a chain (with respect to the inclusion relation) of elements of \mathcal{C} is in \mathcal{C} .

The elements of \mathcal{C} are called convex sets and the pair (X, \mathcal{C}) is called a **convexity space**. For the ordinary convexity, the first two axioms hold, and the third one is trivial.

For an alternative definition of abstract convexity, more suited to discrete geometry, see the *convex geometries* associated with antimatroids.

See also

- Absorbing set
- Bounded set (topological vector space)
- Brouwer fixed-point theorem
- Complex convexity
- Convex hull
- Convex series
- Convex metric space
- Carathéodory's theorem (convex hull)
- Choquet theory
- Helly's theorem
- Holomorphically convex hull
- Integrally-convex set
- John ellipsoid
- Pseudoconvexity
- Radon's theorem
- Shapley–Folkman lemma
- Symmetric set

References

1. Morris, Carla C.; Stark, Robert M. (24 August 2015). *Finite Mathematics: Models and Applications* (<https://books.google.com/books?id=ZgJyCgAAQBAJ&q=convex+region&pg=PA121>). John Wiley & Sons. p. 121. ISBN 9781119015383. Retrieved 5 April 2017.
2. Kjeldsen, Tinne Hoff. "History of Convexity and Mathematical Programming" (<https://web.archive.org/web/20170811100026/http://www.mathunion.org/ICM/ICM2010.4/Main/icm2010.4.3233.3257.pdf>) (PDF). *Proceedings of the International Congress of Mathematicians* (ICM 2010): 3233–3257. doi:10.1142/9789814324359_0187 (https://doi.org/10.1142%2F9789814324359_0187). Archived from the original (<http://www.mathunion.org/ICM/ICM2010.4/Main/icm2010.4.3233.3257.pdf>) (PDF) on 2017-08-11. Retrieved 5 April 2017.
3. Halmos, Paul R. (8 November 1982). *A Hilbert Space Problem Book*. Graduate Texts in Mathematics. Vol. 19 (2nd ed.). New York: Springer-Verlag. p. 5. ISBN 978-0-387-90685-0. OCLC 8169781 (<https://www.worldcat.org/oclc/8169781>).
4. McConnell, Jeffrey J. (2006). *Computer Graphics: Theory Into Practice* (<https://archive.org/details/computergraphics0000mcco/page/130>). p. 130 (<https://archive.org/details/computergraphics0000mcco/page/130>). ISBN 0-7637-2250-2.
5. Weisstein, Eric W. "Concave" (<https://mathworld.wolfram.com/Concave.html>). *MathWorld*.
6. Takayama, Akira (1994). *Analytical Methods in Economics* (https://books.google.com/books?id=_WmZA0MPImEC&pg=PA54). University of Michigan Press. p. 54. ISBN 9780472081356. "An often seen confusion is a "concave set". Concave and convex functions designate certain classes of functions, not of sets, whereas a convex set designates a certain class of sets, and not a class of functions. A "concave set" confuses sets with functions."

7. Corbae, Dean; Stinchcombe, Maxwell B.; Zeman, Juraj (2009). *An Introduction to Mathematical Analysis for Economic Theory and Econometrics* (<https://books.google.com/books?id=j5P83LtzVO8C&pg=PT347>). Princeton University Press. p. 347. ISBN 9781400833085. "There is no such thing as a concave set."
8. Meyer, Robert (1970). "The validity of a family of optimization methods" (<https://minds.wisconsin.edu/bitstream/handle/1793/57508/TR28.pdf?sequence=1>) (PDF). *SIAM Journal on Control and Optimization*. **8**: 41–54. doi:10.1137/0308003 (<https://doi.org/10.1137%2F0308003>). MR 0312915 (<https://www.ams.org/mathscinet-getitem?mr=0312915>)..
9. Soltan, Valeriu, *Introduction to the Axiomatic Theory of Convexity*, Știința, Chișinău, 1984 (in Russian).
10. Singer, Ivan (1997). *Abstract convex analysis*. Canadian Mathematical Society series of monographs and advanced texts. New York: John Wiley & Sons, Inc. pp. xxii+491. ISBN 0-471-16015-6. MR 1461544 (<https://www.ams.org/mathscinet-getitem?mr=1461544>).
11. Lassak, M. (1993). "Approximation of convex bodies by rectangles". *Geometriae Dedicata*. **47**: 111–117. doi:10.1007/BF01263495 (<https://doi.org/10.1007%2FBF01263495>). S2CID 119508642 (<https://api.semanticscholar.org/CorpusID:119508642>).
12. Santaló, L. (1961). "Sobre los sistemas completos de desigualdades entre tres elementos de una figura convexa planas". *Mathematicae Notae*. **17**: 82–104.
13. Brandenberg, René; González Merino, Bernardo (2017). "A complete 3-dimensional Blaschke-Santaló diagram" (<http://mia.ele-math.com/20-22>). *Mathematical Inequalities & Applications* (2): 301–348. doi:10.7153/mia-20-22 (<https://doi.org/10.7153%2Fmia-20-22>). ISSN 1331-4343 (<https://www.worldcat.org/issn/1331-4343>).
14. The empty set is important in Minkowski addition, because the empty set annihilates every other subset: For every subset S of a vector space, its sum with the empty set is empty: $S + \emptyset = \emptyset$.
15. Theorem 3 (pages 562–563): Krein, M.; Šmulian, V. (1940). "On regularly convex sets in the space conjugate to a Banach space". *Annals of Mathematics*. Second Series. Vol. 41. pp. 556–583. doi:10.2307/1968735 (<https://doi.org/10.2307%2F1968735>). JSTOR 1968735 (<https://www.jstor.org/stable/1968735>).
16. For the commutativity of Minkowski addition and convexification, see Theorem 1.1.2 (pages 2–3) in Schneider; this reference discusses much of the literature on the convex hulls of Minkowski sumsets in its "Chapter 3 Minkowski addition" (pages 126–196): Schneider, Rolf (1993). *Convex bodies: The Brunn–Minkowski theory* (<https://archive.org/details/convexbodiesbrun0000schn>). Encyclopedia of mathematics and its applications. Vol. 44. Cambridge: Cambridge University Press. pp. xiv+490. ISBN 0-521-35220-7. MR 1216521 (<https://www.ams.org/mathscinet-getitem?mr=1216521>).
17. Lemma 5.3: Aliprantis, C.D.; Border, K.C. (2006). *Infinite Dimensional Analysis, A Hitchhiker's Guide*. Berlin: Springer. ISBN 978-3-540-29587-7.
18. Zălinescu, C. (2002). *Convex analysis in general vector spaces* (https://archive.org/details/convexanalysisge00zali_934). River Edge, NJ: World Scientific Publishing Co., Inc. p. 7 (https://archive.org/details/convexanalysisge00zali_934/page/n27). ISBN 981-238-067-1. MR 1921556 (<https://www.ams.org/mathscinet-getitem?mr=1921556>).
19. Rawlins G.J.E. and Wood D, "Ortho-convexity and its generalizations", in: *Computational Morphology*, 137-152. Elsevier, 1988.
20. Munkres, James; *Topology*, Prentice Hall; 2nd edition (December 28, 1999). ISBN 0-13-181629-2.
21. van De Vel, Marcel L. J. (1993). *Theory of convex structures*. North-Holland Mathematical Library. Amsterdam: North-Holland Publishing Co. pp. xvi+540. ISBN 0-444-81505-8. MR 1234493 (<https://www.ams.org/mathscinet-getitem?mr=1234493>).

External links

- "Convex subset" (https://www.encyclopediaofmath.org/index.php?title=Convex_subset). *Encyclopedia of Mathematics*. EMS Press. 2001 [1994].
 - [Lectures on Convex Sets](http://www.fmf.uni-lj.si/~lavric/lauritzen.pdf) (<http://www.fmf.uni-lj.si/~lavric/lauritzen.pdf>), notes by Niels Lauritzen, at Aarhus University, March 2010.
-

Retrieved from "https://en.wikipedia.org/w/index.php?title=Convex_set&oldid=1064943728"

This page was last edited on 11 January 2022, at 00:41 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.