

# Effective domain

In convex analysis, a branch of mathematics, the **effective domain** is an extension of the domain of a function defined for functions that take values in the extended real number line  $[-\infty, \infty] = \mathbb{R} \cup \{\pm\infty\}$ .

In convex analysis and variational analysis, a point at which some given extended real-valued function is minimized is typically sought, where such a point is called a global minimum point. The effective domain of this function is defined to be the set of all points in this function's domain at which its value is not equal to  $+\infty$ ,<sup>[1]</sup> where the effective domain is defined this way because it is only these points that have even a remote chance of being a global minimum point. Indeed, it is common practice in these fields to set a function equal to  $+\infty$  at a point specifically to *excluded* that point from even being considered as a potential solution (to the minimization problem).<sup>[1]</sup> Points at which the function takes the value  $-\infty$  (if any) belong to the effective domain because such points are considered acceptable solutions to the minimization problem,<sup>[1]</sup> with the reasoning being that if such a point was not acceptable as a solution then the function would have already been set to  $+\infty$  at that point instead.

When a minimum point (in  $X$ ) of a function  $f : X \rightarrow [-\infty, \infty]$  is to be found but  $f$ 's domain  $X$  is a proper subset of some vector space  $V$ , then it often technically useful to extend  $f$  to all of  $V$  by setting  $f(x) := +\infty$  at every  $x \in V \setminus X$ .<sup>[1]</sup> By definition, no point of  $V \setminus X$  belongs to the effective domain of  $f$ , which is consistent with the desire to find a minimum point of the original function  $f : X \rightarrow [-\infty, \infty]$  rather than of the newly defined extension to all of  $V$ .

If the problem is instead a maximization problem (which would be clearly indicated) then the effective domain instead consists of all points in the function's domain at which it is not equal to  $-\infty$ .

## Contents

- [Definition](#)
- [Characterizations](#)
- [Properties](#)
- [See also](#)
- [References](#)

## Definition

Suppose  $f : X \rightarrow [-\infty, \infty]$  is a map valued in the extended real number line  $[-\infty, \infty] = \mathbb{R} \cup \{\pm\infty\}$  whose domain, which is denoted by **domain**  $f$ , is  $X$  (where  $X$  will be assumed to be a subset of some vector space whenever this assumption is necessary). Then the **effective domain** of  $f$  is denoted by **dom**  $f$  and typically defined to be the set<sup>[1][2][3]</sup>

$$\text{dom } f = \{x \in X : f(x) < +\infty\}$$

unless  $f$  is a concave function or the maximum (rather than the minimum) of  $f$  is being sought, in which case the **effective domain** of  $f$  is instead the set<sup>[2]</sup>

$$\operatorname{dom} f = \{x \in X : f(x) > -\infty\}.$$

In convex analysis and variational analysis,  $\operatorname{dom} f$  is usually assumed to be  $\operatorname{dom} f = \{x \in X : f(x) < +\infty\}$  unless clearly indicated otherwise.

## Characterizations

Let  $\pi_X : X \times \mathbb{R} \rightarrow X$  denote the canonical projection onto  $X$ , which is defined by  $(x, r) \mapsto x$ . The effective domain of  $f : X \rightarrow [-\infty, \infty]$  is equal to the image of  $f$ 's epigraph  $\operatorname{epi} f$  under the canonical projection  $\pi_X$ . That is

$$\operatorname{dom} f = \pi_X (\operatorname{epi} f) = \{x \in X : \text{there exists } y \in \mathbb{R} \text{ such that } (x, y) \in \operatorname{epi} f\}.$$
<sup>[4]</sup>

For a maximization problem (such as if the  $f$  is concave rather than convex), the effective domain is instead equal to the image under  $\pi_X$  of  $f$ 's hypograph.

## Properties

If a function *never* takes the value  $+\infty$ , such as if the function is real-valued, then its domain and effective domain are equal.

A function  $f : X \rightarrow [-\infty, \infty]$  is a proper convex function if and only if  $f$  is convex, the effective domain of  $f$  is nonempty, and  $f(x) > -\infty$  for every  $x \in X$ .<sup>[4]</sup>

## See also

- Proper convex function
- Epigraph (mathematics)
- Hypograph (mathematics)

## References

- Rockafellar & Wets 2009, pp. 1–28.
  - Aliprantis, C.D.; Border, K.C. (2007). *Infinite Dimensional Analysis: A Hitchhiker's Guide* (3 ed.). Springer. p. 254. doi:10.1007/3-540-29587-9 (https://doi.org/10.1007%2F3-540-29587-9). ISBN 978-3-540-32696-0.
  - Föllmer, Hans; Schied, Alexander (2004). *Stochastic finance: an introduction in discrete time* (2 ed.). Walter de Gruyter. p. 400. ISBN 978-3-11-018346-7.
  - Rockafellar, R. Tyrrell (1997) [1970]. *Convex Analysis*. Princeton, NJ: Princeton University Press. p. 23. ISBN 978-0-691-01586-6.
- Rockafellar, R. Tyrrell; Wets, Roger J.-B. (26 June 2009). *Variational Analysis*. Grundlehren der mathematischen Wissenschaften. Vol. 317. Berlin New York: Springer Science & Business Media.

ISBN 9783642024313. OCLC 883392544 (<https://www.worldcat.org/oclc/883392544>).

---

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Effective\\_domain&oldid=1041521836](https://en.wikipedia.org/w/index.php?title=Effective_domain&oldid=1041521836)"

---

**This page was last edited on 31 August 2021, at 00:48 (UTC).**

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.