

Ursescu theorem

In mathematics, particularly in [functional analysis](#) and [convex analysis](#), the **Ursescu theorem** is a theorem that generalizes the [closed graph theorem](#), the [open mapping theorem](#), and the [uniform boundedness principle](#).

Contents

Ursescu Theorem

Statement

Corollaries

Closed graph theorem

Uniform boundedness principle

Open mapping theorem

Additional corollaries

Related theorems

Simons' theorem

Robinson–Ursescu theorem

See also

Notes

References

Ursescu Theorem

The following notation and notions are used, where $\mathcal{R} : X \rightrightarrows Y$ is a [multivalued function](#) and S is a non-empty subset of a [topological vector space](#) X :

- the [affine span](#) of S is denoted by $\mathbf{aff} S$ and the [linear span](#) is denoted by $\mathbf{span} S$.
- $S^i := \mathbf{aint}_X S$ denotes the [algebraic interior](#) of S in X .
- $^i S := \mathbf{aint}_{\mathbf{aff}(S-S)} S$ denotes the [relative algebraic interior](#) of S (i.e. the algebraic interior of S in $\mathbf{aff}(S - S)$).
- $^{ib} S := ^i S$ if $\mathbf{span}(S - s_0)$ is [barreled](#) for some/every $s_0 \in S$ while $^{ib} S := \emptyset$ otherwise.
 - If S is convex then it can be shown that for any $x \in X$, $x \in ^{ib} S$ if and only if the cone generated by $S - x$ is a barreled linear subspace of X or equivalently, if and only if $\bigcup_{n \in \mathbb{N}} n(S - x)$ is a barreled linear subspace of X
- The **domain** of \mathcal{R} is $\mathbf{Dom} \mathcal{R} := \{x \in X : \mathcal{R}(x) \neq \emptyset\}$.
- The **image** of \mathcal{R} is $\mathbf{Im} \mathcal{R} := \bigcup_{x \in X} \mathcal{R}(x)$. For any subset $A \subseteq X$, $\mathcal{R}(A) := \bigcup_{x \in A} \mathcal{R}(x)$.
- The **graph** of \mathcal{R} is $\mathbf{gr} \mathcal{R} := \{(x, y) \in X \times Y : y \in \mathcal{R}(x)\}$.

- \mathcal{R} is **closed** (respectively, **convex**) if the graph of \mathcal{R} is closed (resp. convex) in $X \times Y$.
 - Note that \mathcal{R} is convex if and only if for all $x_0, x_1 \in X$ and all $r \in [0, 1]$, $r\mathcal{R}(x_0) + (1-r)\mathcal{R}(x_1) \subseteq \mathcal{R}(rx_0 + (1-r)x_1)$.
- The **inverse** of \mathcal{R} is the multifunction $\mathcal{R}^{-1} : Y \rightrightarrows X$ defined by $\mathcal{R}^{-1}(y) := \{x \in X : y \in \mathcal{R}(x)\}$. For any subset $B \subseteq Y$, $\mathcal{R}^{-1}(B) := \bigcup_{y \in B} \mathcal{R}^{-1}(y)$.
 - If $f : X \rightarrow Y$ is a function, then its inverse is the multifunction $f^{-1} : Y \rightrightarrows X$ obtained from canonically identifying f with the multifunction $f : X \rightrightarrows Y$ defined by $x \mapsto \{f(x)\}$.
- $\text{int}_T S$ is the topological interior of S with respect to T , where $S \subseteq T$.
- $\text{rint } S := \text{int}_{\text{aff } S} S$ is the interior of S with respect to $\text{aff } S$.

Statement

Theorem^[1] (Ursescu) — Let X be a complete semi-metrizable locally convex topological vector space and $\mathcal{R} : X \rightrightarrows Y$ be a closed convex multifunction with non-empty domain. Assume that $\text{span}(\text{Im } \mathcal{R} - y)$ is a barrelled space for some/every $y \in \text{Im } \mathcal{R}$. Assume that $y_0 \in {}^i(\text{Im } \mathcal{R})$ and let $x_0 \in \mathcal{R}^{-1}(y_0)$ (so that $y_0 \in \mathcal{R}(x_0)$). Then for every neighborhood U of x_0 in X , y_0 belongs to the relative interior of $\mathcal{R}(U)$ in $\text{aff}(\text{Im } \mathcal{R})$ (that is, $y_0 \in \text{int}_{\text{aff}(\text{Im } \mathcal{R})} \mathcal{R}(U)$). In particular, if ${}^{ib}(\text{Im } \mathcal{R}) \neq \emptyset$ then ${}^{ib}(\text{Im } \mathcal{R}) = {}^i(\text{Im } \mathcal{R}) = \text{rint}(\text{Im } \mathcal{R})$.

Corollaries

Closed graph theorem

Closed graph theorem — Let X and Y be Fréchet spaces and $T : X \rightarrow Y$ be a linear map. Then T is continuous if and only if the graph of T is closed in $X \times Y$.

Proof

For the non-trivial direction, assume that the graph of T is closed and let $\mathcal{R} := T^{-1} : Y \rightrightarrows X$. It is easy to see that $\text{gr } \mathcal{R}$ is closed and convex and that its image is X . Given $x \in X$, (Tx, x) belongs to $Y \times X$ so that for every open neighborhood V of Tx in Y , $\mathcal{R}(V) = T^{-1}(V)$ is a neighborhood of x in X . Thus T is continuous at x . ■

Uniform boundedness principle

Uniform boundedness principle — Let X and Y be Fréchet spaces and $T : X \rightarrow Y$ be a bijective linear map. Then T is continuous if and only if $T^{-1} : Y \rightarrow X$ is continuous. Furthermore, if T is continuous then T is an isomorphism of Fréchet spaces.

Proof

Apply the closed graph theorem to T and T^{-1} . ■

Open mapping theorem

Open mapping theorem — Let X and Y be Fréchet spaces and $T : X \rightarrow Y$ be a continuous surjective linear map. Then T is an open map.

Proof

Clearly, T is a closed and convex relation whose image is Y . Let U be a non-empty open subset of X , let y be in $T(U)$, and let x in U be such that $y = Tx$. From the Ursescu theorem it follows that $T(U)$ is a neighborhood of y . ■

Additional corollaries

The following notation and notions are used for these corollaries, where $\mathcal{R} : X \rightrightarrows Y$ is a multifunction, S is a non-empty subset of a topological vector space X :

- a **convex series with elements of S** is a series of the form $\sum_{i=1}^{\infty} r_i s_i$ where all $s_i \in S$ and $\sum_{i=1}^{\infty} r_i = 1$ is a series of non-negative numbers. If $\sum_{i=1}^{\infty} r_i s_i$ converges then the series is called **convergent** while if $(s_i)_{i=1}^{\infty}$ is bounded then the series is called **bounded** and **b-convex**.
- S is **ideally convex** if any convergent b-convex series of elements of S has its sum in S .
- S is **lower ideally convex** if there exists a Fréchet space Y such that S is equal to the projection onto X of some ideally convex subset B of $X \times Y$. Every ideally convex set is lower ideally convex.

Corollary — Let X be a barreled first countable space and let C be a subset of X . Then:

1. If C is lower ideally convex then $C^i = \text{int } C$.
2. If C is ideally convex then $C^i = \text{int } C = \text{int}(\text{cl } C) = (\text{cl } C)^i$.

Related theorems

Simons' theorem

Simons' theorem^[2] — Let X and Y be first countable with X locally convex. Suppose that $\mathcal{R} : X \rightrightarrows Y$ is a multimap with non-empty domain that satisfies condition (Hwx) or else assume that X is a Fréchet space and that \mathcal{R} is lower ideally convex. Assume that $\text{span}(\text{Im } \mathcal{R} - y)$ is barreled for some/every $y \in \text{Im } \mathcal{R}$. Assume that $y_0 \in {}^i(\text{Im } \mathcal{R})$ and let $x_0 \in \mathcal{R}^{-1}(y_0)$. Then for every neighborhood U of x_0 in X , y_0 belongs to the relative interior of $\mathcal{R}(U)$ in $\text{aff}(\text{Im } \mathcal{R})$ (i.e. $y_0 \in \text{int}_{\text{aff}(\text{Im } \mathcal{R})} \mathcal{R}(U)$). In particular, if ${}^{ib}(\text{Im } \mathcal{R}) \neq \emptyset$ then ${}^{ib}(\text{Im } \mathcal{R}) = {}^i(\text{Im } \mathcal{R}) = \text{rint}(\text{Im } \mathcal{R})$.

Robinson–Ursescu theorem

The implication (1) \implies (2) in the following theorem is known as the Robinson–Ursescu theorem.^[3]

Robinson–Ursescu theorem^[3] — Let $(X, \|\cdot\|)$ and $(Y, \|\cdot\|)$ be normed spaces and $\mathcal{R} : X \rightrightarrows Y$ be a multimap with non-empty domain. Suppose that Y is a barreled space, the graph of \mathcal{R} verifies condition (Hwx), and that $(x_0, y_0) \in \text{gr } \mathcal{R}$. Let C_X (resp. C_Y) denote the closed unit ball in X (resp. Y) (so $C_X = \{x \in X : \|x\| \leq 1\}$). Then the following are equivalent:

1. y_0 belongs to the algebraic interior of $\text{Im } \mathcal{R}$.
2. $y_0 \in \text{int } \mathcal{R}(x_0 + C_X)$.
3. There exists $B > 0$ such that for all $0 \leq r \leq 1$, $y_0 + BrC_Y \subseteq \mathcal{R}(x_0 + rC_X)$.
4. There exist $A > 0$ and $B > 0$ such that for all $x \in x_0 + AC_X$ and all $y \in y_0 + AC_Y$, $d(x, \mathcal{R}^{-1}(y)) \leq B \cdot d(y, \mathcal{R}(x))$.
5. There exists $B > 0$ such that for all $x \in X$ and all $y \in y_0 + BC_Y$,

$$d(x, \mathcal{R}^{-1}(y)) \leq \frac{1 + \|x - x_0\|}{B - \|y - y_0\|} \cdot d(y, \mathcal{R}(x)).$$

See also

- [Closed graph theorem](#) – Theorem relating continuity to graphs
- [Closed graph theorem \(functional analysis\)](#) – Theorems for deducing continuity
- [Open mapping theorem \(functional analysis\)](#) – Condition for a linear operator to be open
- [Surjection of Fréchet spaces](#) – Characterization of surjectivity
- [Uniform boundedness principle](#) – A theorem stating that pointwise boundedness implies uniform boundedness
- [Webbed space](#) – Spaces where open mapping and closed graphs theorems hold

Notes

1. [Zălinescu 2002](#), p. 23.
2. [Zălinescu 2002](#), p. 22-23.
3. [Zălinescu 2002](#), p. 24.

References

- [Zălinescu, Constantin \(30 July 2002\). *Convex Analysis in General Vector Spaces* \(https://archive.org/details/convexanalysisge00zali_934\). River Edge, N.J. London: World Scientific Publishing. ISBN 978-981-4488-15-0. MR 1921556 \(https://www.ams.org/mathscinet-getitem?mr=1921556\). OCLC 285163112 \(https://www.worldcat.org/oclc/285163112\) – via Internet Archive.](#)
- [Baggs, Ivan \(1974\). "Functions with a closed graph" \(https://www.ams.org/\). *Proceedings of the American Mathematical Society*. **43** \(2\): 439–442. doi:10.1090/S0002-9939-1974-0334132-8 \(https://doi.org/10.1090/S0002-9939-1974-0334132-8\). ISSN 0002-9939 \(https://www.worldcat.org/issn/0002-9939\).](#)

Retrieved from "https://en.wikipedia.org/w/index.php?title=Ursescu_theorem&oldid=1053133912"

This page was last edited on 2 November 2021, at 03:30 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.