


John ellipsoid

In mathematics, the **John ellipsoid** or **Löwner-John ellipsoid** $E(K)$ associated to a convex body K in n -dimensional Euclidean space \mathbf{R}^n can refer to the n -dimensional ellipsoid of maximal volume contained within K or the ellipsoid of minimal volume that contains K .

Often, the minimal volume ellipsoid is called as Löwner ellipsoid, and the maximal volume ellipsoid as the John ellipsoid (although John worked with the minimal volume ellipsoid in its original paper).^[1] One also refer to the minimal volume circumscribed ellipsoid as the **outer Löwner-John ellipsoid** and the maximum volume inscribed ellipsoid as the **inner Löwner-John ellipsoid**.^[2]



Outer Löwner-John ellipsoid containing a set of a points in \mathbf{R}^2

Contents

- [Properties](#)
- [Applications](#)
- [See also](#)
- [References](#)

Properties

The John ellipsoid is named after the German-American mathematician Fritz John, who proved in 1948 that each convex body in \mathbf{R}^n contains a unique circumscribed ellipsoid of minimal volume and that the dilation of this ellipsoid by factor $1/n$ is contained inside the convex body.^[3]

The inner Löwner-John ellipsoid $E(K)$ of a convex body $K \subset \mathbf{R}^n$ is a closed unit ball B in \mathbf{R}^n if and only if $B \subseteq K$ and there exists an integer $m \geq n$ and, for $i = 1, \dots, m$, real numbers $c_i > 0$ and unit vectors $u_i \in \mathbf{S}^{n-1} \cap \partial K$ such that^[4]

$$\sum_{i=1}^m c_i u_i = 0$$

and, for all $x \in \mathbf{R}^n$

$$x = \sum_{i=1}^m c_i (x \cdot u_i) u_i.$$

Applications

Computing Löwner-John ellipsoids has applications in obstacle collision detection for robotic systems, where the distance between a robot and its surrounding environment is estimated using a best ellipsoid fit.^[5]

It also has applications in portfolio optimization with transaction costs.^[6]

See also

- Banach–Mazur compactum – Set of n-dimensional subspaces of a normed space made into a compact metric space.
- Steiner inellipse, the special case of the inner Löwner-John ellipsoid for a triangle.
- Fat object, related to radius of largest contained ball.

References

1. Güler, Osman; Gürtuna, Filiz (2012). "Symmetry of convex sets and its applications to the extremal ellipsoids of convex bodies" (<http://www.tandfonline.com/doi/abs/10.1080/10556788.2011.626037>). *Optimization Methods and Software*. **27** (4–5): 735–759. doi:10.1080/10556788.2011.626037 (<https://doi.org/10.1080%2F10556788.2011.626037>). ISSN 1055-6788 (<https://www.worldcat.org/issn/1055-6788>).
2. Ben-Tal, A. (2001). *Lectures on modern convex optimization : analysis, algorithms, and engineering applications* (<https://www.worldcat.org/oclc/46538510>). Nemirovskii, Arkadii Semenovich. Philadelphia, PA: Society for Industrial and Applied Mathematics. ISBN 0-89871-491-5. OCLC 46538510 (<https://www.worldcat.org/oclc/46538510>).
3. John, Fritz. "Extremum problems with inequalities as subsidiary conditions". *Studies and Essays Presented to R. Courant on his 60th Birthday*, January 8, 1948, 187–204. Interscience Publishers, Inc., New York, N. Y., 1948. OCLC 1871554 (<https://www.worldcat.org/oclc/1871554>) MR30135 (<https://mathscinet.ams.org/mathscinet-getitem?mr=30135>)
4. Ball, Keith M. (1992). "Ellipsoids of maximal volume in convex bodies". *Geom. Dedicata*. **41** (2): 241–250. arXiv:math/9201217 (<https://arxiv.org/abs/math/9201217>). doi:10.1007/BF00182424 (<https://doi.org/10.1007%2FBF00182424>). ISSN 0046-5755 (<https://www.worldcat.org/issn/0046-5755>).
5. Rimon, Elon; Boyd, Stephen (1997). "Obstacle Collision Detection Using Best Ellipsoid Fit". *Journal of Intelligent and Robotic Systems*. **18** (2): 105–126. doi:10.1023/A:1007960531949 (<https://doi.org/10.1023%2FA%3A1007960531949>).
6. Shen, Weiwei; Wang, Jun (2015). "Transaction costs-aware portfolio optimization via fast Löwner-John ellipsoid approximation" (<https://web.archive.org/web/20170116192444/https://pdfs.semanticscholar.org/7b31/2141616f092137c12397a47d11d94ddcea78.pdf>) (PDF). *Proceedings of the Twenty-Ninth AAAI Conference on Artificial Intelligence (AAAI2015)*: 1854–1860. Archived from the original (<https://pdfs.semanticscholar.org/7b31/2141616f092137c12397a47d11d94ddcea78.pdf>) (PDF) on 2017-01-16.
- Gardner, Richard J. (2002). "The Brunn-Minkowski inequality" (<https://doi.org/10.1090%2FS0273-0979-02-00941-2>). *Bull. Amer. Math. Soc. (N.S.)*. **39** (3): 355–405 (electronic). doi:10.1090/S0273-0979-02-00941-2 (<https://doi.org/10.1090%2FS0273-0979-02-00941-2>). ISSN 0273-0979 (<https://www.worldcat.org/issn/0273-0979>).

Retrieved from "https://en.wikipedia.org/w/index.php?title=John_ellipsoid&oldid=1020696648"

This page was last edited on 30 April 2021, at 15:11 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.