


Shared KEGG pathways

Within-sample top 15%; both datasets FDR < 0.05

Oxidative phosphorylation	2.38 ***	2.33 ***
Parkinson disease	2.23 ***	2.07 ***
Allograft rejection	1.95 **	1.96 ***
Asthma	1.90 **	2.13 ***
Leishmaniasis	1.91 **	1.88 ***
Proteasome	1.88 **	1.89 **
Graft-versus-host disease	1.85 *	1.98 ***
Huntington disease	1.92 ***	1.84 ***
Th1 and Th2 cell differentiation	1.96 **	1.84 ***
Tuberculosis	1.83 **	1.96 ***
Epstein-Barr virus infection	1.84 ***	1.83 ***
Autoimmune thyroid disease	1.80 *	1.99 ***
Th17 cell differentiation	1.80 **	2.04 ***
Thermogenesis	1.92 ***	1.78 ***
Cardiac muscle contraction	1.77 *	1.94 ***
C-type lectin receptor signaling pathway	1.77 **	1.80 ***
Ribosome	2.79 ***	1.76 ***
Retrograde endocannabinoid signaling	1.76 *	1.93 ***
Alzheimer disease	1.75 ***	1.93 ***
Systemic lupus erythematosus	1.75 *	1.73 **
Pertussis	1.70 *	1.76 **
Rheumatoid arthritis	1.70 *	2.16 ***
Amoebiasis	1.70 *	1.67 **
Osteoclast differentiation	1.66 *	2.02 ***
Necroptosis	1.65 *	1.67 **
Human cytomegalovirus infection	1.65 **	1.70 ***
Intestinal immune network for IgA production	1.76 *	1.64 *
B cell receptor signaling pathway	1.74 *	1.61 *
Rap1 signaling pathway	1.59 *	1.60 **
Mitophagy	1.67 *	1.56 *
Salmonella infection	1.55 *	1.96 ***
Human T-cell leukemia virus 1 infection	1.52 *	1.53 **
Chemokine signaling pathway	1.52 *	1.85 ***
Kaposi sarcoma-associated herpesvirus infection	1.51 *	1.72 ***
Pathways in cancer	1.51 **	1.47 ***
Proteoglycans in cancer	1.56 *	1.47 *
Human immunodeficiency virus 1 infection	1.62 **	1.47 *
MAPK signaling pathway	1.50 *	1.46 *
Natural killer cell mediated cytotoxicity	2.11 ***	1.45 *
Influenza A	1.55 *	1.41 *

Relapse
scRNA-seq

STAT5B N642H
bulk RNA-seq



* FDR < 0.05; ** < 0.01; *** < 0.001
Pathway names do not imply the corresponding disease is present.