

1 Negative Niche Construction Favors the  
2 Evolution of Cooperation

3  
4 Brian D. Connelly<sup>1</sup>, Katherine J. Dickinson<sup>1</sup>, Sarah P. Hammarlund<sup>1,2</sup>, and  
5 Benjamin Kerr<sup>1,3</sup>

6 <sup>1</sup>Department of Biology and BEACON Center for the Study of Evolution in  
7 Action, University of Washington, Seattle, Washington, 98195, USA

8 <sup>2</sup>Current Address: Department of Zoology, University of Oxford, Oxford OX1  
9 3PS, United Kingdom

10 <sup>3</sup>E-mail: kerrb@uw.edu

## 11 Abstract

12 By benefitting others at a cost to themselves, cooperators face an unrelenting  
13 threat from defectors, or individuals that avail themselves of the cooperative  
14 benefit without contributing. A longstanding challenge to evolutionary biol-  
15 ogy is to understand the mechanisms that support the vast displays of cooper-  
16 ation that nevertheless exist. Hammarlund et al. recently demonstrated that  
17 cooperation can persist by hitchhiking along with beneficial non-social adapta-  
18 tions. Importantly, cooperators play an active role in this process. In spatially-  
19 structured environments, clustered cooperator populations reach greater densi-  
20 ties, which creates more mutational opportunities to gain beneficial non-social  
21 adaptations. Cooperation rises in abundance by association with these adap-  
22 tations. However once adaptive opportunities have been exhausted, the free  
23 ride ends as cooperators are displaced by equally-adapted defectors. Using an  
24 agent-based model, we demonstrate that the selective feedback that is created  
25 as populations alter their environments can maintain cooperation indefinitely.  
26 We show that cooperator success depends specifically on negative niche con-  
27 struction. Here, negative niche construction acts as a perpetual source of  
28 adaptive opportunities. As populations adapt, they further alter their envi-  
29 ronment in ways that reveal additional opportunities for adaptation. Despite  
30 being independent of niche construction in our model, cooperation feeds this  
31 cycle. We show that by reaching larger densities, populations of cooperators  
32 are better able to adapt both to changing environments and to the constant  
33 threat posed by defectors. We relate these findings to previous studies from

34 the niche construction literature and discuss how this model could be extended  
35 to provide a greater understanding of how cooperation evolves in the complex  
36 environments in which it is found.

## 37 Introduction

38 Cooperative behaviors are common across all branches of the tree of life. In-  
39 sects divide labor within their colonies, plants and soil bacteria exchange es-  
40 sential nutrients, birds care for others' young, and the trillions of cells in the  
41 human body coordinate to provide vital functions. Each instance of cooper-  
42 ation presents an evolutionary challenge: How can individuals that sacrifice  
43 their own well-being to help others avoid subversion by those that do not? Over  
44 time, we would expect these *defectors* to rise in abundance at the expense of  
45 others, eventually driving cooperators—and perhaps the entire population—to  
46 extinction.

47 Several factors can prevent this *tragedy of the commons* (Hamilton, 1964;  
48 Nowak, 2006; West *et al.*, 2007). One such factor involves non-random so-  
49 cial interaction, in which cooperators benefit more from the cooperative act  
50 than defectors. This can occur when cooperators are clustered together in  
51 spatially-structured populations (Fletcher and Doebeli, 2009; Nadell *et al.*,  
52 2010; Kuzdzal-Fick *et al.*, 2011) or when cooperators use communication  
53 (Brown and Johnstone, 2001; Darch *et al.*, 2012) or other cues (Sinervo *et al.*,  
54 2006; Gardner and West, 2010; Veelders *et al.*, 2010) to cooperate condition-  
55 ally with kin. Cooperation can also be bolstered by pleiotropic connections to  
56 personal benefits (Foster *et al.*, 2004; Dandekar *et al.*, 2012) or through associ-  
57 ation with alleles encoding self-benefitting traits (Asfahl *et al.*, 2015). In these  
58 cases, the alleles may provide private benefits that are completely independent  
59 from the public benefits of cooperation. In asexual populations of cooperators

60 and defectors, this sets the stage for an “adaptive race” in which both types  
61 vie for the first highly beneficial adaptation (Waite and Shou, 2012; Morgan  
62 *et al.*, 2012). The tragedy of the commons can be deferred if a cooperator, by  
63 chance, wins the adaptive race.

64 Hammarlund et al. (2015) recently showed that in spatially-structured pop-  
65 ulations, the “Hankshaw effect” can give cooperators a substantial leg up on  
66 defectors in an adaptive race. This advantage is reminiscent of Sissy Han-  
67 kshaw, a fictional character in Tom Robbins’ *Even Cowgirls Get the Blues*,  
68 whose oversized thumbs—which were otherwise an impairment—made her a  
69 prolific hitchhiker. Similarly, cooperation is costly, but it increases local pop-  
70 ulation density. As a result, cooperators are more likely to acquire beneficial  
71 mutations. By hitchhiking along with these adaptations, cooperation can rise  
72 in abundance. Nevertheless, this advantage is fleeting. As soon as the oppor-  
73 tunities for adaptation are exhausted, cooperators are once again at a selective  
74 disadvantage against equally-adapted defectors that arise via mutation. How-  
75 ever, Hammarlund et al. (2015) also demonstrated that cooperation can be  
76 maintained when frequent environmental changes produce a steady stream of  
77 new adaptive opportunities. Although organisms typically find themselves in  
78 dynamic environments, the nature and frequency of these changes might not  
79 ensure long-term cooperator survival.

80 However, organisms do more than simply experience changing environments  
81 passively. Through their activities, their interactions with others, and even  
82 their deaths, organisms constantly modify their environment. This niche con-  
83 struction process can produce evolutionary feedback loops in which environ-

84 mental change alters selection, which, in turn, alters the distribution of types  
85 and their corresponding influence on the environment (Odling-Smee *et al.*,  
86 2003). The nature of this feedback can have dramatic evolutionary conse-  
87 quences. One critical distinction is whether the constructing type is favored in  
88 the resulting environment. Under positive niche construction, selection favors  
89 the constructor, and evolution stagnates as this type fixes. Under negative  
90 niche construction, selection favors a type other than the constructor, which  
91 creates an opportunity for adaptation. If an adapted type also engages in  
92 negative niche construction, cycles of construction and adaptation can ensue,  
93 such that populations find themselves continually chasing beneficial mutations  
94 as their adaptive landscape perpetually shifts.

95 Here, we show that the selective feedbacks that result from niche construction  
96 can maintain cooperation indefinitely. We find that it is specifically negative  
97 niche construction that is responsible for this result because of the endless  
98 opportunities for adaptation that it produces. These results indicate that  
99 cooperators can ensure their survival when they play an active role in their  
100 own evolution.

## 101 **Methods**

102 Building upon Hammarlund et al. (2015), we describe an individual-based  
103 model in which cooperators and defectors evolve and compete in a population  
104 of subpopulations (i.e., a metapopulation). Through mutations, individuals  
105 gain adaptations to their environment, which increase reproductive fitness,

106 and allow those lineages to rise in abundance. More successful lineages spread  
107 to neighboring subpopulations by migration.

108 In this expanded model, subpopulations additionally modify their local envi-  
109 ronment. As this process occurs, environmental changes feed back to affect  
110 selection. We explore how niche construction affects the evolution of coop-  
111 eration; specifically, how cooperative behavior can hitchhike with adaptive  
112 mutations to modified environments. Box 1 provides additional information  
113 about the model.

## 114 **Model Description**

### 115 **Individual Genotypes and Adaptation**

116 Each individual has a haploid genome with  $L + 1$  loci (see [Table 1](#) for model  
117 parameters and their values). Different alleles at each locus are represented by  
118 different integers. A binary allele at the first locus (here, locus zero) determines  
119 whether that individual is a cooperator (1), which carries fitness cost  $c$ , or a  
120 defector (0). The remaining  $L$  loci are *adaptive loci*, and are each occupied by  
121 0 or a value from the set  $\{1, 2, \dots, A\}$ . Allele 0 represents a lack of adaptation,  
122 while a non-zero allele represents one of the  $A$  possible adaptations at that  
123 locus.

124 These non-zero alleles signify two types of adaptations, both of which increase  
125 fitness. First, adaptations to the external environment confer a fitness benefit  $\delta$ .  
126 This selective value is the same regardless of which non-zero allele is present  
127 and is not affected by other individuals. We assume  $\delta > c$ , which allows a

128 minimally adapted cooperator to recoup the cost of cooperation and gain a  
129 fitness advantage.

## 130 Niche Construction and Selective Feedbacks

131 Individual fitness is also affected by aspects of the local environment that are  
132 affected by organisms. We implicitly represent this constructed “niche” based  
133 on the specific allelic states present in the subpopulation. As allelic states  
134 change, the subpopulation alters its environment, creating a unique niche. As  
135 described below, the specific alleles that are present at each locus matter.

136 In our model, the feedback from niche construction takes the form of den-  
137 sity dependent selection, and individuals evolve to better match their niche.  
138 Specifically, the selective value of non-zero allele  $a$  at adaptive locus  $l$ —and  
139 consequently the fitness of an individual carrying that allele—increases with  
140 the number of individuals in the subpopulation that have allele  $a - 1$  at locus  
141  $l - 1$ . For example, when  $L = 5$  and  $A = 6$ , and allele 4 has fixed at locus 2,  
142 a genotype with allele 5 at locus 3 is favored. And once allele 5 has fixed at  
143 locus 3, the niche that this population constructs will favor allele 6 at locus 4  
144 (see Box 1). As a consequence, genotypes with sequentially increasing allelic  
145 states will tend to evolve. We treat both adaptive loci and their non-zero al-  
146 lelic states as “circular”: the selective value of an allele at locus 1 is affected by  
147 the allelic composition of the subpopulation at locus  $L$ . Similarly, the selective  
148 value of allele 1 at any locus increases with the number of individuals carrying  
149 allele  $A$  at the previous locus. This circularity is represented by the function



150  $\beta(x, X)$ , which gives the integer that is below an arbitrary value  $x$  in the set  
 151  $\{1, 2, \dots, X\}$ :

$$\beta(x, X) = \text{mod}_X(x - 2 + X) + 1 \quad (1)$$

152 Here,  $\text{mod}_X(x)$  is the integer remainder when dividing  $x$  by  $X$ . Using this  
 153 function, the selective value of allele  $a$  at adaptive locus  $l$  is increased by  $\epsilon$  for  
 154 each individual in the subpopulation that has allele  $\beta(a, A)$  at locus  $\beta(l, L)$ .  
 155 Thus,  $\epsilon$  specifies the intensity of selection due to niche construction.

156 Consider a genotype  $g$  with allelic state  $a_{g,l}$  at locus  $l$ ; the fitness of an indi-  
 157 vidual with this genotype is defined as:

$$W_g = z - \underbrace{ca_{g,0}}_{\text{cost of cooperation}} + \underbrace{\delta \sum_{l=1}^L I(a_{g,l})}_{\text{adaptation to external environment}} + \underbrace{\epsilon \sum_{l=1}^L n(\beta(a_{g,l}, A), \beta(l, L))}_{\text{adaptation to constructed environment}} \quad (2)$$

158 where  $z$  is a baseline fitness,  $n(a, l)$  is the number of individuals in the sub-  
 159 population with allele  $a$  at locus  $l$ , and  $I(a)$  indicates whether a given allele is  
 160 non-zero:

$$I(a) = \begin{cases} 1 & \text{if } a \in \{1, 2, \dots, A\} \\ 0 & \text{otherwise} \end{cases} \quad (3)$$

161 Thus, an individual's fitness is determined both by adaptations to the exter-  
 162 nal environment and by adaptations to its constructed environment. **Box 1**

163 illustrates the process of adaptation to the constructed environment. While  
 164 cooperation is costly, we assume its effects are independent of the external and  
 165 constructed components of the environment.

## 166 **Population Growth and the Benefit of Cooperation**

167 Cooperation enables a subpopulation to reach a greater density. This benefit  
 168 affects all individuals equally and accumulates linearly with the proportion  
 169 of cooperators in the subpopulation. If  $p$  is the proportion of cooperators  
 170 present at the beginning of a growth cycle, then that subpopulation reaches  
 171 the following size:

$$S(p) = S_{min} + p(S_{max} - S_{min}) \quad (4)$$

172 During subpopulation growth, individuals compete through differential repro-  
 173 duction. Each individual's probability of success is determined by its fitness.  
 174 The composition of a subpopulation with size  $P$  and cooperator proportion  $p$   
 175 after growth is multinomial with parameters  $S(p)$  and  $\{\pi_1, \pi_2, \dots, \pi_P\}$ , where  
 176  $\pi_i$  represents individual  $i$ 's reproductive fitness relative to others in the sub-  
 177 population (using Equation 2).

## 178 **Mutation**

179 For simplicity, we apply mutations after subpopulation growth. Mutations  
 180 occur independently at each locus and cause an allelic state change. At the

181 binary cooperation locus, mutations occur at rate  $\mu_c$ . These mutations flip  
 182 the allelic state, causing cooperators to become defectors and vice versa. Mu-  
 183 tations occur at rate  $\mu_a$  at each adaptive locus. These mutations replace the  
 184 existing allele with a random selection from the set  $\{0\} \cup \{1, 2, \dots, A\}$ . Be-  
 185 cause mutations are stochastic, the allelic sequences that evolve depend on  
 186 which allele arises first and at which locus.

## 187 **Migration**

188 Populations are composed of  $N^2$  patches arranged as an  $N \times N$  lattice, where  
 189 each patch can support a subpopulation. After mutation, individuals emigrate  
 190 to an adjacent patch with probability  $m$ . During each migration event, a single  
 191 destination patch is randomly chosen from each source patch's Moore neigh-  
 192 borhood, which is composed of the nearest 8 patches on the lattice. Because  
 193 the population lattice has boundaries, patches located on the periphery have  
 194 smaller neighborhoods.

## 195 **Population Initialization and Simulation**

196 Following Hammarlund et al. (2015), we begin simulations with sparse pop-  
 197 ulations. Subpopulations are first seeded at all patches with size  $S(p_0)$  and  
 198 cooperator proportion  $p_0$ . The population is then thinned. Each individual  
 199 survives this bottleneck with probability  $\sigma$ . Starting from this initial state,  
 200 simulations then proceed for  $T$  cycles, where each discrete cycle consists of  
 201 subpopulation growth, mutation, migration, and dilution. Dilution reduces

202 the population to support growth in the next cycle. Each individual remains  
203 with probability  $d$ , regardless of its genotype.

## 204 **Simulation Source Code and Software Dependencies**

205 The simulation software and configurations for the experiments reported are  
206 available online.<sup>1</sup> Simulations used Python 3.4, NumPy 1.9.1, Pandas 0.15.2  
207 (McKinney, 2010), and NetworkX 1.9.1 (Hagberg *et al.*, 2008). Data analy-  
208 ses were performed with R 3.1.3 (R Core Team, 2015). Reported confidence  
209 intervals were estimated by bootstrapping with 1000 resamples.

## 210 **Results**

211 Using the model described in the previous section, we perform simulations  
212 that follow the evolution of cooperation in a population consisting of subpopu-  
213 lations that are connected by spatially-limited migration. Individuals increase  
214 their competitiveness by gaining adaptations. While cooperation does not di-  
215 rectly affect the fitness benefits that these adaptations confer, cooperation has  
216 indirect effects on the adaptive process. Specifically, cooperation increases  
217 subpopulation density. As a result, larger subpopulations of cooperators ex-  
218 perience more mutational opportunities. Cooperation can rise in abundance  
219 by hitchhiking along with beneficial mutations, which compensate for the cost  
220 of cooperation. Importantly, subpopulations alter their local environments,

---

<sup>1</sup>These materials will be made public prior to publication.

221 which feeds back to influence selection. Here, we explore how such niche con-  
222 struction affects the evolution of cooperation.

## 223 Cooperation Persists with Niche Construction

224 Without any opportunity for adaptation ( $L = 0$ ), cooperators are swiftly elim-  
225 inated in competition with defectors (Figure 1A). Despite an initial lift in  
226 cooperator abundance due to increased productivity, the cost of cooperation  
227 becomes disadvantageous as migration mixes the initially isolated subpopula-  
228 tions. When populations can adapt to the external environment ( $L = 5$  and  
229  $\delta > 0$ ), but niche construction is absent ( $\epsilon = 0$ ), cooperators are maintained  
230 only transiently (Figure 1B). Here, larger cooperator subpopulations can more  
231 quickly adapt to their external environment. However, as previously described  
232 by Hammarlund et al. (2015), cooperation is subsequently lost once popula-  
233 tions become fully adapted to their environment. This occurs when isogenic  
234 defectors (i.e., defectors with identical adaptive loci) arise via mutation and  
235 displace cooperators due to their selective advantage. However, when niche  
236 construction feeds back to influence selection ( $\epsilon > 0$ ), cooperation persists in  
237 the majority of the replicate populations (Figure 1C). We see in Figure 2A that  
238 despite oscillations in the proportion of cooperators, cooperation is maintained  
239 at high levels in these populations.

## **Fitness Increases Alone do not Support Persisting Cooperation**

In the model, adaptations to both the external environment and the constructed environment contribute to an individual's fitness. To determine whether cooperation is maintained solely due to the larger selective values that result from the contributions of niche construction ( $\epsilon$ ), we performed simulations in which these contributions were removed ( $\epsilon = 0$ ), and we instead increased the fitness benefits conferred by adaptation to the external, non-constructed environment ( $\delta = 0.6$ ). In doing so, we conservatively estimate the selective effects of niche construction by supplementing the selective benefits of adaptations to the external environment by the maximum possible selective benefit that results from niche construction. Nevertheless, we find that simply increasing selective values does not enable cooperators to persist (Figure 2B). Niche construction, therefore, plays a decisive role here.

## **Negative Niche Construction is Critical to Cooperator Persistence**

In our model, an adaptation to a constructed aspect of the environment initiates a new instance of niche construction, leading to sequentially increasing allelic states across the adaptive loci. Under certain conditions, this construction always makes the constructor suboptimal for the niche it creates (see Box 1). This negative niche construction occurs when the number of adaptive al-

261 leles ( $A$ ) does not divide evenly into the number of adaptive loci ( $L$ ). In such  
 262 a case, any sequence of integers on the circular genome will always contain a  
 263 break in the sequence; that is, one locus with an allele that is not one less than  
 264 the allele at the next locus. Given this unavoidable mismatch, any type that  
 265 has fixed will always construct a niche that favors selection for a new type.  
 266 When negative niche construction is removed (by setting  $L = 5$ ,  $A = 5$ ), coop-  
 267 erators are again driven extinct after an initial lift in abundance (Figure 2C).  
 268 These results indicate that the type of niche construction matters. Specifically,  
 269 negative niche construction is crucial for maintaining cooperation.

## 270 **Selective Feedbacks Limit Defector Invasion**

271 The adaptation resulting from selective feedbacks can limit invasion by de-  
 272 fectors, which arise either through immigration from neighboring patches or  
 273 through mutation from a cooperator ancestor. The latter challenge is par-  
 274 ticularly threatening, as these isogenic defectors are equally adapted, yet do  
 275 not incur the cost of cooperation. As demonstrated in Figure 3A, isogenic  
 276 defectors rapidly spread when introduced at a single patch in the center of a  
 277 population of cooperators if mutations do not occur. However, when coopera-  
 278 tors can gain adaptations via mutation, cooperators resist defector invasion in  
 279 over half of the replicate populations (Figure 3B). Figure 4 depicts one such  
 280 instance. In that population, defectors quickly began to spread. However, an  
 281 adaptation arose in a neighboring cooperator population. This type spreads  
 282 more quickly, stopping the spread of defectors and eventually driving them to

283 extinction. Because this adaption occurred in a cooperator population, coop-  
284 eration was able to hitchhike to safety. Importantly, this new cooperator was  
285 favored because of the niche that its ancestor created. Here, cooperators can  
286 find safety in numbers—because their larger populations have more mutational  
287 opportunities, they are more likely to gain adaptations that rescue them from  
288 invasion. Further, the larger number of cooperators more strongly construct  
289 their niche, and thus more strongly favors an adapted type. This allows that  
290 type to appear and to spread more quickly in the population. **Figure 3C** shows  
291 how quickly an adapted cooperator type can invade a population of defectors.

## 292 Discussion

293 Despite their negative effects, deleterious traits can rise in abundance due to ge-  
294 netic linkage with other traits that are strongly favored by selection (Maynard  
295 Smith and Haigh, 1974). In a process termed the “Hankshaw effect”, Hammar-  
296 lund et al. (2015) recently demonstrated that cooperation can actively prolong  
297 its existence by increasing its likelihood of hitchhiking with a beneficial trait.  
298 In that work and in ours, populations of cooperators grow to a higher density  
299 than those of defectors. Because of this, these cooperator populations experi-  
300 ence more mutations and are therefore more likely to gain adaptations. While  
301 this process does favor cooperation in the short term, it eventually reaches a  
302 dead end; when the opportunities for adaptation are exhausted, and coopera-  
303 tors can no longer hitchhike, they face extinction. Here, we have considered  
304 whether niche construction might serve to perpetually generate new adaptive



305 opportunities and thus favor cooperation indefinitely.  
 306 When niche construction occurs, cooperation can indeed persist (Figures 1C  
 307 and 2A). In our model, niche construction introduces additional selective ef-  
 308 fects that influence the evolutionary process, leading to a more pronounced  
 309 Hankshaw effect. However, simply raising the fitness benefits conferred by  
 310 adaptations does not maintain cooperators at high proportion (Figure 2B).  
 311 This indicates that niche construction does play a crucial role.  
 312 Further, we find that it is specifically *negative* niche construction that main-  
 313 tains cooperation (Figure 2C). As cooperator and defector types gain adapta-  
 314 tions, they change the environment ways that favor other types. Because of  
 315 this, negative niche construction serves as a perpetual source of adaptation.  
 316 Here we observe another facet of the Hankshaw effect: because populations  
 317 of cooperators are larger, they are better able to respond to the adaptive op-  
 318 portunities that result from negative niche construction. These opportunities  
 319 can allow cooperators to resist invasion by defectors (Figure 3B). Although  
 320 defectors initially have an advantage by saving on the cost of cooperation, sub-  
 321 populations of cooperators can quickly gain an advantage because they are  
 322 larger. Even after defector invasion, subpopulations of cooperators are more  
 323 likely to produce the next adapted mutant, which can then displace the slower  
 324 evolving defectors. It is these recurring cycles of defector invasion and cooper-  
 325 ator adaptation that underlie the oscillations in cooperator proportion that we  
 326 see in Figure 2A. When an adaptation to the constructed environment does  
 327 not occur in the cooperator background before the defector dominates, the  
 328 defector can drive the cooperator to extinction. This is something that we see

329 occur stochastically in Figures 2A and 3B.

## 330 Cooperation as Niche Construction

331 In our model, cooperation is orthogonal to niche construction, which allows us  
332 to focus on hitchhiking. However, by increasing the size of the subpopulation,  
333 this form of cooperation can itself be seen as form of niche construction. Co-  
334 operative benefits often take similar forms in natural systems. For example,  
335 bacteria produce a host of extracellular products that scavenge soluble iron  
336 (Griffin *et al.*, 2004), digest large proteins (Diggle *et al.*, 2007; Darch *et al.*,  
337 2012), and reduce the risk of predation (Cosson *et al.*, 2002). As in our model,  
338 such cooperative acts are likely to increase local population density. While  
339 many studies have focused on how the environment affects the evolution of  
340 these cooperative traits, relatively few have addressed how the environmental  
341 changes created by public goods feed back to influence evolution.

342 Perhaps most similar to our work, Van Dyken and Wade (2012) demonstrated  
343 that when two negative niche constructing, cooperative behaviors co-evolve,  
344 selection can increasingly favor these traits, which were disfavored when alone.  
345 In that model, “reciprocal niche construction” occurred when the negative feed-  
346 back created by one strategy positively influenced selection on the other, cre-  
347 ating a perpetual cycle that maintained both forms of cooperation. Arguably,  
348 this can be viewed as an instance of hitchhiking: the currently-maladaptive  
349 form of cooperation is maintained by association with the adaptive form.

350 When dispersal is limited, competition among kin can undermine cooperation.

351 To separate kin competition from kin selection, Lehmann (2007) developed  
352 a model in which the selective feedbacks produced by a cooperative, niche-  
353 constructing behavior only benefitted future generations. Kin competition  
354 thereby was reduced, and cooperation instead benefitted descendants. This  
355 work highlights an important aspect of niche construction—often, the rate of  
356 selective feedback from niche construction is different from the rate at which  
357 populations grow.

## 358 **Evolution at Multiple Timescales**

359 In our work, the niche is modeled implicitly by the composition of the popula-  
360 tion. Any changes, therefore, in the population produce immediate effects on  
361 the constructed environment and the resulting feedbacks. However, timescales  
362 in our model could be de-coupled in two ways. First, cooperators modify their  
363 niche by enabling their population to reach larger density (Equation 4). These  
364 increased population sizes play a critical role by effectively increasing the rate  
365 of evolution in these populations. Because of the importance of this process, it  
366 would be very informative to explore how sensitive our results are to changes  
367 in how long the increases in population size are upheld. Similarly, changes  
368 in the timescale at which the niche at a patch change also have potential to  
369 dramatically alter our results. Not only would changes in timescale affect the  
370 selective values of alleles as the population changed, but they could also in-  
371 fluence whether or not populations were able to evolve adapted types and if  
372 so, how well those adapted types can propagate through the population to

373 address the threat of a defector.

374 Other studies, while not focused on cooperation, have similarly shown that the  
375 timescales at which niche construction feedbacks occur can strongly influence  
376 evolutionary outcomes (Laland *et al.*, 1996, 1999). This perspective is likely to  
377 be crucial for understanding the evolution of cooperative behaviors like the pro-  
378 duction of public goods. In these instances, environmental changes are likely  
379 to occur on different timescales than growth, which can have profound effects.  
380 For example, a multitude of factors including protein durability (Brown and  
381 Taddei, 2007; Kümmerli and Brown, 2010), diffusion (Allison, 2005; Driscoll  
382 and Pepper, 2010), and resource availability (Zhang and Rainey, 2013; Ghoul  
383 *et al.*, 2014) influence both the rate and the degree to which public goods alter  
384 the environment. While Lehmann (2007) showed that cooperation was favored  
385 when selective feedbacks act over longer timescales, niche construction may in  
386 fact hinder cooperation when selection is more quickly altered. For example,  
387 when public goods accumulate in the environment, cooperators must decrease  
388 production to remain competitive (Kümmerli and Brown, 2010; Dumas and  
389 Kümmerli, 2012). This favors cooperation that occurs facultatively, perhaps  
390 by sensing the abiotic (Bernier *et al.*, 2011; Koestler and Waters, 2014) or  
391 biotic environment (Brown and Johnstone, 2001; Darch *et al.*, 2012). To allow  
392 our model to address how traits such as these evolve, we would first need to  
393 de-couple the niche from the composition of the population by representing  
394 the niche explicitly.

## Cooperation and Niche Construction in Host-Symbiont Co-Evolution

As niche construction becomes more independent, it develops its own state and dynamics. A logical next step, then, could be to treat the environment as a biological entity itself, which could introduce additional evolutionary feedbacks. As the host population changes, so too does selection on their symbiont populations. Here, evolutionary outcomes depend greatly on the degree of shared interest between the host and symbiont. Future models could explicitly capture the environment as a biological entity to explore the rich coevolutionary dynamics that these systems might offer.

For example, the cooperative production of virulence factors by the human pathogen *P. aeruginosa* in lung infections is harmful to hosts with cystic fibrosis (Harrison, 2007). Similarly to what we have shown in this work, these antagonistic, negative niche constructing behaviors might actually work to maintain these infections. If this is the case, however, perhaps the case could be made for developing treatments that target the selective feedback loop that is created by niche construction. If these populations do indeed perpetually benefit from adaptations that are created by niche construction, as we have shown, case could perhaps be made for developing treatments that target the selective feedback loop that provides adaptive opportunities in these spatial environments. While the idea of removing negative selective feedbacks and supporting stability may seem counterintuitive, if it makes the population more susceptible, then perhaps pairing such a treatment with ones in which mutants

are introduced (e.g., Rumbaugh et al. (2009)), could significantly improve host fitness. Expanding models such as ours to address the additional dynamics present in host-symbiont systems such as these could be quite productive.

Or conversely, cooperative light production by *A. fischeri* is vital for the survival of its host, the Hawaiian bobtail squid (Ruby, 1996). While our current model and that of Van Dyken and Wade (2012) have showed that negative niche construction can play a decisive role in the evolution of cooperation, this instance of positive niche construction is a textbook example of cooperation and mutualism. Therefore, a greater understanding of the additional feedbacks created in symbioses such as these could be gained from modeling. Similar to our model, these host-symbiont systems likely have many other traits that are orthogonal to cooperation. Perhaps combinations of certain types of behaviors are important for maintaining cooperation, similar to what was shown by Van Dyken and Wade (2012).

It was recently argued that incorporating the effects of niche construction is critical for improving our understanding of viral evolution (Hamblin *et al.*, 2014) and evolution in co-infecting parasites (Hafer and Milinski, 2015). Incorporating host dynamics, co-evolution, and the feedbacks that they produce into models is likely to be equally important for gaining an understanding of how cooperative behaviors, both positive and negative, evolve in these host-symbiont settings.

## 439 **Acknowledgments**

440 We are grateful to Peter Conlin, Sylvie Estrela, Carrie Glenney, Martha Kor-  
441 nelius, and Luis Zaman for helpful comments on the manuscript, and to Anu-  
442 raag Pakanati for assistance with simulations. This material is based upon  
443 work supported by the National Science Foundation Postdoctoral Research  
444 Fellowship in Biology under Grant No. DBI-1309318 (to BDC) and under  
445 Cooperative Agreement No. DBI-0939454 (BEACON STC). Computational  
446 resources were provided by an award from Google Inc. (to BDC and BK).

## 447 **Box 1: Description of niche construction in our** 448 **model**

449 **(A) Individuals.** The genome of each individual consists of a single *coop-*  
450 *eration locus* and  $L$  *adaptive loci* (here,  $L = 5$ ). At the cooperation locus  
451 (labeled 0), this individual has allele 1, making it a cooperator. The adaptive  
452 loci (labeled 1-5) are arranged as a circular chromosome, where each locus has  
453 an integer allele between 0 and  $A$ , inclusive. In the description that follows,  
454 we focus exclusively on these adaptive loci. Genotypes are given by their al-  
455 lelic states starting with locus 1. For instance, the genotype shown here is  
456  $[2,0,5,2,1]$ . Because of their circular structure, allele 2 at the first locus follows  
457 allele 1 at the fifth locus.

458 **(B) Niche Construction.** Consider a subpopulation fixed for genotype  
459  $[1,2,0,0,0]$ . This subpopulation constructs environment  $E_{[1,2,0,0,0]}$ . Every non-  
460 zero allele influences selection at the next locus, favoring sequential allelic  
461 states. In this constructed environment, allele 3 at locus 3 would be fa-  
462 vored. If genotype  $[1,2,3,0,0]$  arises via mutation, it is expected to fix. How-  
463 ever, genotype  $[1,2,3,0,0]$  affects the environment differently than its ancestor.  
464 As  $[1,2,3,0,0]$  rises in abundance, the constructed environment changes from  
465  $E_{[1,2,0,0,0]}$  to  $E_{[1,2,3,0,0]}$ , which favors  $[1,2,3,4,0]$ .

466 **(C) Niche Construction and Adaptation.** The evolutionary transition  
467 shown in Part B is indicated in the dashed box. Here, we depict entire sub-  
468 populations fixed for a genotype using a single instance of that genotype. Simi-



469 larly, an arrow represents niche construction and adaptation to the constructed  
 470 environment. We start with a case in which there are five alleles ( $A = 5$ ). Sub-  
 471 populations begin with the non-adapted genotype  $[0,0,0,0,0]$ , shown on the far  
 472 left. A non-zero allele is introduced via mutation, which represents an adapta-  
 473 tion to external aspects of the environment. Here, allele 1 arises and fixes at  
 474 locus 1. The remainder of this figure focuses on adaptation to the constructed  
 475 aspects of the environment. This genotype has a “mismatch” (shown by the  
 476 red sector), because  $E_{[1,0,0,0,0]}$  favors  $[1,2,0,0,0]$ . Assuming allele 2 arises at the  
 477 second locus, it will be selected, creating a “match” at the first and second  
 478 loci (green sector). Now there is a mismatch between the second and third  
 479 loci in the resulting environment, which a new round of mutation and selection  
 480 corrects, and so on. The green sector grows as the red sector shifts clockwise.  
 481 Since allele 1 follows allele 5 when  $A = 5$ , there is no mismatch as we cycle  
 482 through the circular genotype  $[1,2,3,4,5]$ . This genotype will now remain in  
 483 the subpopulation.

484 **(D) Negative Niche Construction.** A different case emerges when a mis-  
 485 match exists as the genotype cycles back to where we started. Here, we change  
 486 the number of alleles to six ( $A = 6$ ). As shown on the far left, we begin with  
 487 a subpopulation fixed for genotype  $[1,2,3,4,5]$ . This genotype has a mismatch,  
 488 because the niche constructed by allele 5 favors allele 6 (not 1) at the next  
 489 locus (in this case locus 1). A mutant with genotype  $[6,2,3,4,5]$  has a fitness ad-  
 490 vantage and can fix in  $E_{[1,2,3,4,5]}$ . However, as this type constructs  $E_{[6,2,3,4,5]}$ , a  
 491 new mismatch appears. In this instance of negative niche construction, adapt-  
 492 ing to correct one mismatch generates a new mismatch. This system can never

493 escape its mismatches—the red sector just shifts clockwise around the genome  
494 perpetually.

## 495 Figures

### 496 Figure 1






Figure 1: **Adaptation, Hitchhiking, and the Evolution of Cooperation.** Curves show the average cooperator proportion among replicate populations for the duration of simulations, and shaded areas indicate 95% confidence intervals. Unless otherwise noted, parameter values are listed in [Table 1](#). **(A)** Without any opportunity to adapt ( $L = 0$ ), cooperation is quickly lost. **(B)** When adaptation can occur ( $L = 5$ ), but niche construction does not affect selection ( $\epsilon = 0$ ), cooperators rise in abundance by hitchhiking along with adaptations to the external environment. Nevertheless, this effect is transient, and cooperators eventually become extinct. **(C)** Niche construction enables cooperation to be maintained indefinitely. In the majority of populations, cooperation remained the dominant strategy. The trajectories of individual populations are shown in Figure 2A.

497 **Figure 2**



**Figure 2: Niche Construction and the Evolution of Cooperation.** The proportion of cooperators present in each replicate population is shown for the duration of simulations. **(A)** Despite some oscillations, cooperation dominates in 13 of 18 populations when niche construction affects selection. **(B)** When the selective effects of niche construction ( $\epsilon$ ) are removed, and the selective benefit of adaptation to the external environment ( $\delta$ ) is increased to compensate, cooperators are driven to extinction by isogenic defectors that arise by mutation ( $\epsilon = 0$ ,  $\delta = 0.6$ ). Note that cooperation was not present after initialization in one replicate population. **(C)** Cooperators are also driven to extinction without negative niche construction ( $A = 5$ ).

498 **Figure 3**



**Figure 3: Niche Construction and Invasion.** Curves trace the proportion of cooperators present in each replicate population for the duration of simulations ( $T = 1000$ ). In each simulation, a rare type was initiated at a single patch in the center of the population lattice ( $N^2 = 121$ ). Unless otherwise noted, mutations are disabled in these ecological simulations to highlight the dynamics of invasion ( $\mu_a = 0, \mu_c = 0$ ). **(A)** When cooperators and defectors are isogenic (i.e., both types have stress alleles [1,2,3,4,5]), rare defectors quickly invade and drive cooperators to extinction due to the cost of cooperation. Defectors were stochastically eliminated in 2 replicate populations. **(B)** However, negative niche construction creates adaptive opportunities that enable cooperators to resist invasion by isogenic defectors. Here, cooperation remained the dominant in 91 of 160 populations ( $\mu_a = 0.00005$ ). Results from simulations where mutations also occurred at the cooperation locus are shown in Figure S1. **(C)** In fact, an adapted cooperator type (stress alleles [6,2,3,4,5], see Box 1) can swiftly displace defectors when isogenic defectors cannot arise or adapt via mutation.

499 **Figure 4**




Figure 4: **Cooperator Adaptation Prevents Defector Invasion.** Here we depict the distribution of dominant types among subpopulations over time for one representative simulation in which isogenic defectors arise. To highlight the effects of adaptation, mutations did not occur at the cooperation locus ( $\mu_c = 0$ ). At time  $t = 0$  (upper left panel), a single isogenic defector population (red) is placed among cooperator populations (light blue). Because these defectors do not bear the costs of cooperation, they spread ( $t = 272$ ). However, cooperators in one population gain an adaptation that gives them a fitness advantage over defectors (second panel, dark blue, lower left). At  $t = 325$ , defectors continue to invade cooperator populations. However, the adapted cooperator type spreads more quickly due to its fitness advantage, invading both defector populations and ancestral cooperator populations ( $t = 390$ ), until it eventually fixes in the population ( $t = 500$ ). At  $t = 690$ , a new cooperator type emerges that is favored due to negative niche construction (orange). This new type spreads rapidly ( $t = 812$ ) until reaching fixation ( $t = 900$ ). At this point, it becomes susceptible to invasion by the next “adapted” cooperator type, and the cycle continues.

500 **Box 1 Figures**

501 **Figure B1**




Figure B1: Figure for Box 1



502 **Supplemental Figures**

503 **Supplemental Figure 1**




Figure S1: **Defector Invasion with Mutations.** The proportion of co-operators present in each replicate population is shown for the duration of simulations ( $T = 1000$ ). When mutations occur both at the adaptive loci and the cooperation locus ( $\mu_a = \mu_c = 0.00005$ ), cooperation remains dominant in 58 of 160 replicate populations.

Table 1: Model parameters and their value

Parameter	Description	Base Value
$L$	Number of adaptive loci	5
$c$	Fitness cost of cooperation	0.1
$A$	Number of alleles	6
$\delta$	Fitness benefit, adaptation to external environment	0.3
$\epsilon$	Fitness benefit, adaptation to constructed environment	0.00015
$z$	Baseline fitness	1
$S_{min}$	Minimum subpopulation size	800
$S_{max}$	Maximum subpopulation size	2000
$\mu_a$	Mutation rate (adaptation)	$10^{-5}$
$\mu_c$	Mutation rate (cooperation)	$10^{-5}$
$N^2$	Number of patches	625
$m$	Migration rate	0.05
$p_0$	Initial cooperator proportion	0.5
$\sigma$	Survival rate at population initialization	$10^{-5}$
$T$	Number of simulation cycles	3000
$d$	Subpopulation dilution factor	0.1

## References

- Allison, S.D. 2005. Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. *Ecology Letters*, **8**: 626–635.
- Asfahl, K.L., Walsh, J., Gilbert, K. and Schuster, M. 2015. Non-social adaptation defers a tragedy of the commons in *Pseudomonas aeruginosa* quorum sensing. *The ISME Journal*, doi: [10.1038/ismej.2014.259](https://doi.org/10.1038/ismej.2014.259).
- Bernier, S.P., Ha, D.-G., Khan, W., Merritt, J.H.M. and O'Toole, G.A. 2011. Modulation of *Pseudomonas aeruginosa* surface-associated group behaviors by individual amino acids through c-di-GMP signaling. *Research in Microbiology*, **162**: 680–688.
- Brown, S.P. and Johnstone, R.A. 2001. Cooperation in the dark: Signalling and collective action in quorum-sensing bacteria. *Proceedings of the Royal Society of London B: Biological Sciences*, **268**: 961–965.
- Brown, S.P. and Taddei, F. 2007. The durability of public goods changes the dynamics and nature of social dilemmas. *PLoS ONE*, **2**: e593.
- Cosson, P., Zulianello, L., Join-Lambert, O., Faurisson, F., Gebbie, L. and Benghezal, M.*et al.* 2002. *Pseudomonas aeruginosa* virulence analyzed in a *Dictyostelium discoideum* host system. *Journal of Bacteriology*, **184**: 3027–3033.
- Dandekar, A.A., Chugani, S. and Greenberg, E.P. 2012. Bacterial quorum sensing and metabolic incentives to cooperate. *Science*, **338**: 264–266.

527 Darch, S.E., West, S.A., Winzer, K. and Diggle, S.P. 2012. Density-dependent  
528 fitness benefits in quorum-sensing bacterial populations. *Proceedings of the*  
529 *National Academy of Sciences*, **109**: 8259–8263.

530 Diggle, S.P., Griffin, A.S., Campbell, G.S. and West, S.A. 2007. Cooperation  
531 and conflict in quorum-sensing bacterial populations. *Nature*, **450**: 411–414.

532 Driscoll, W.W. and Pepper, J.W. 2010. Theory for the evolution of diffusible  
533 external goods. *Evolution*, **64**: 2682–2687.

534 Dumas, Z. and Kümmerli, R. 2012. Cost of cooperation rules selection for  
535 cheats in bacterial metapopulations. *Journal of Evolutionary Biology*, **25**:  
536 473–484.

537 Fletcher, J.A. and Doebeli, M. 2009. A simple and general explanation for the  
538 evolution of altruism. *Proceedings of the Royal Society B: Biological Sciences*,  
539 **276**: 13–19.

540 Foster, K., Shaulsky, G., Strassmann, J., Queller, D. and Thompson, C. 2004.  
541 Pleiotropy as a mechanism to stabilize cooperation. *Nature*, **431**: 693–696.

542 Gardner, A. and West, S.A. 2010. Greenbeards. *Evolution*, **64**: 25–38.

543 Ghoul, M., West, S.A., Diggle, S.P. and Griffin, A.S. 2014. An experimental  
544 test of whether cheating is context dependent. *Journal of Evolutionary Biology*,  
545 **27**: 551–556.

546 Griffin, A.S., West, S.A. and Buckling, A. 2004. Cooperation and competition  
547 in pathogenic bacteria. *Nature*, **430**: 1024–1027.

548 Hafer, N. and Milinski, M. 2015. When parasites disagree: Evidence for

549 parasite-induced sabotage of host manipulation. *Evolution*, **69**: 611–620.

550 Hagberg, A.A., Schult, D.A. and Swart, P.J. 2008. Exploring network struc-  
551 ture, dynamics, and function using NetworkX. In: *Proceedings of the 7th*  
552 *Python in Science Conference (SciPy2008)*, pp. 11–15.

553 Hamblin, S.R., White, P.A. and Tanaka, M.M. 2014. Viral niche construction  
554 alters hosts and ecosystems at multiple scales. *Trends in Ecology & Evolution*,  
555 **29**: 594–599.

556 Hamilton, W.D. 1964. The genetical evolution of social behaviour I & II.  
557 *Journal of Theoretical Biology*, **7**: 1–52.

558 Hammarlund, S.P., Connelly, B.D., Dickinson, K.J. and Kerr, B. 2015. The  
559 evolution of cooperation by the Hankshaw effect. *bioRxiv*, doi: [10.1101/016667](https://doi.org/10.1101/016667).  
560 Cold Spring Harbor Labs Journals.

561 Harrison, F. 2007. Microbial ecology of the cystic fibrosis lung. *Microbiology*,  
562 **153**: 917–923.

563 Koestler, B.J. and Waters, C.M. 2014. Bile acids and bicarbonate inversely  
564 regulate intracellular cyclic di-GMP in *Vibrio cholerae*. *Infection and Immu-*  
565 *nity*, **82**: 3002–3014.

566 Kuzdzal-Fick, J.J., Fox, S.A., Strassmann, J.E. and Queller, D.C. 2011. High  
567 relatedness is necessary and sufficient to maintain multicellularity in *Dic-*  
568 *tyostelium*. *Science*, **334**: 1548–1551.

569 Kümmerli, R. and Brown, S.P. 2010. Molecular and regulatory properties of  
570 a public good shape the evolution of cooperation. *Proceedings of the National*

571 *Academy of Sciences*, **107**: 18921–18926.

572 Laland, K.N., Odling-Smee, F.J. and Feldman, M.W. 1999. Evolutionary con-  
573 sequences of niche construction and their implications for ecology. *Proceedings*  
574 *of the National Academy of Sciences*, **96**: 10242–10247.

575 Laland, K.N., Odling-Smee, F.J. and Feldman, M.W. 1996. The evolutionary  
576 consequences of niche construction: A theoretical investigation using two-locus  
577 theory. *Journal of Evolutionary Biology*, **9**: 293–316.

578 Lehmann, L. 2007. The evolution of trans-generational altruism: Kin selection  
579 meets niche construction. *Journal of Evolutionary Biology*, **20**: 181–189.

580 Maynard Smith, J. and Haigh, J. 1974. The hitch-hiking effect of a favourable  
581 gene. *Genetics Research*, **23**: 23–35.

582 McKinney, W. 2010. Data structures for statistical computing in Python. In:  
583 *Proceedings of the 9th Python in Science Conference* (S. van der Walt and J.  
584 Millman, eds), pp. 51–56.

585 Morgan, A.D., Quigley, B.J.Z., Brown, S.P. and Buckling, A. 2012. Selection  
586 on non-social traits limits the invasion of social cheats. *Ecology Letters*, **15**:  
587 841–846.

588 Nadell, C.D., Foster, K.R. and Xavier, J.B. 2010. Emergence of spatial struc-  
589 ture in cell groups and the evolution of cooperation. *PLoS Computational*  
590 *Biology*, **6**: e1000716.

591 Nowak, M.A. 2006. Five rules for the evolution of cooperation. *Science*, **314**:  
592 1560–1563.

- 593 Odling-Smee, F.J., Laland, K.N. and Feldman, M.W. 2003. *Niche construc-*  
594 *tion: The neglected process in evolution*. Princeton University Press.
- 595 R Core Team. 2015. *R: A language and environment for statistical computing*.  
596 Vienna, Austria: R Foundation for Statistical Computing.
- 597 Ruby, E.G. 1996. Lessons from a cooperative, bacterial-animal association:  
598 The *Vibrio fischeri*–*Euprymna scolopes* light organ symbiosis. *Annual Review*  
599 *of Microbiology*, **50**: 591–624.
- 600 Rumbaugh, K.P., Diggle, S.P., Watters, C.M., Ross-Gillespie, A., Griffin, A.S.  
601 and West, S.A. 2009. Quorum sensing and the social evolution of bacterial  
602 virulence. *Current Biology*, **19**: 341–345.
- 603 Sinervo, B., Chaine, A., Clobert, J., Calsbeek, R., Hazard, L. and Lancaster,  
604 L.*et al.* 2006. Self-recognition, color signals, and cycles of greenbeard mutu-  
605 alism and altruism. *Proceedings of the National Academy of Sciences*, **103**:  
606 7372–7377.
- 607 Van Dyken, J.D. and Wade, M.J. 2012. Origins of altruism diversity II: Run-  
608 away coevolution of altruistic strategies via “reciprocal niche construction”.  
609 *Evolution*, **66**: 2498–2513.
- 610 Veelders, M., Brückner, S., Ott, D., Unverzagt, C., Mösch, H.-U. and Essen,  
611 L.-O. 2010. Structural basis of flocculin-mediated social behavior in yeast.  
612 *Proceedings of the National Academy of Sciences*, **107**: 22511–22516.
- 613 Waite, A.J. and Shou, W. 2012. Adaptation to a new environment allows coop-  
614 erators to purge cheaters stochastically. *Proceedings of the National Academy*  
615 *of Sciences*, **109**: 19079–19086.



- 616 West, S.A., Griffin, A.S. and Gardner, A. 2007. Evolutionary explanations for  
617 cooperation. *Current Biology*, **17**: R661–R672.
- 618 Zhang, X.-X. and Rainey, P.B. 2013. Exploring the sociobiology of pyoverdin-  
619 producing *Pseudomonas*. *Evolution*, **67**: 3161–3174.