

TODO title

TODO

Abstract

Through their interactions, their activities, and even their mere presence, organisms change the environment for themselves and others. This “niche construction” process becomes particularly interesting when it creates evolutionary feedback, whereby selective pressures are altered in response to environmental change. Here we consider how niche construction influences the evolution of cooperation, which has been a long-standing challenge to evolutionary theory. We simulate populations of individuals that cooperatively produce a public good that permits increased growth in a stressful environment and investigate how local- and global-scale niche construction affects the ability of these populations to resist invasion by non-producing cheats. We find that niche construction profoundly impacts the evolution of cooperation by creating new opportunities for adaptation. Cooperators are able to escape subversion by cheats as long as niche construction clears these paths of adaptation. This work provides a crucial step towards understanding how evolution occurs in complex environments like those found in nature.

20 Introduction

21 Cooperative behaviors are common across all branches of the tree of life. In-
22 sects divide labor within their colonies, plants and soil bacteria exchange es-
23 sential nutrients, birds care for others' young, and the trillions of cells in the
24 human body restrain their growth and coordinate to provide vital functions.
25 Each instance of cooperation presents an evolutionary challenge: How can in-
26 dividuals that sacrifice their own well-being to help others avoid subversion by
27 those that do not? Over time, we would expect these *defectors* to rise in abun-
28 dance at the expense of others, eventually driving cooperators—and perhaps
29 the entire population—to extinction.

30 Several factors can defer this potential *tragedy of the commons* (Hamilton,
31 1964; Hardin, 1968; Nowak, 2006; West *et al.*, 2007b). For example, coopera-
32 tors must benefit more from the cooperative act than others. This can occur
33 when cooperators are clustered together in spatially-structured populations
34 (Fletcher and Doebeli, 2009; Nadell *et al.*, 2010; Kuzdzal-Fick *et al.*, 2011) or
35 when cooperators use communication (Brown and Johnstone, 2001; Darch *et*
36 *al.*, 2012) or other cues (Sinervo *et al.*, 2006; Gardner and West, 2010; Veelders
37 *et al.*, 2010) to cooperate conditionally with kin. Interestingly, cooperation can
38 also be bolstered by genetic linkage with self-benefitting traits (Foster *et al.*,
39 2004; Dandekar *et al.*, 2012; Asfahl *et al.*, 2015), setting the stage for an “adap-
40 tive race” in which cooperators and defectors vie for the first highly-beneficial
41 non-social adaptation (Waite and Shou, 2012; Morgan *et al.*, 2012).

42 Hankshaw and Kerr (2015) recently showed that in spatially-structured popu-

lations, cooperators can gain a substantial leg up on defectors in an adaptive race when the cooperative behavior increases local population density, thus increasing the likelihood of acquiring beneficial non-social mutations. By hitchhiking along with these adaptations, the cooperative trait can rapidly rise in abundance. Nevertheless, this advantage is fleeting. As soon as the opportunities for adaptation are exhausted, cooperators are once again at a disadvantage against defectors, particularly those that arise from within via mutations that disable cooperation. However, Hankshaw and Kerr (2015) demonstrated that cooperation can be maintained indefinitely when frequent environmental changes produce a steady stream of non-social adaptive opportunities. Although organisms typically find themselves in dynamic environments, change might not occur at a rate that provides sufficient adaptive opportunities to ensure long-term cooperator survival.

In this work, we examine whether the changes in selection that arise as organisms modify their environment can create opportunities that allow cooperators to be maintained by hitchhiking. We expand upon the model presented in Hankshaw and Kerr (2015), by allowing populations to alter their local environment based on the the presence of different non-social adaptations. Frequency-dependent selection on these adaptations creates an eco-evolutionary feedback that increasingly favors the adaptations present in each population. We first examine how the intensity of these feedbacks affects the hitchhiking process. Because the production of public goods increases population density, populations containing cooperators will exert a greater influence on their environment. As a result, these environments will be more quickly brought to states where

67 fitness is higher.

68 As populations construct unique niches, they potentially decrease the threat
69 of invasion from neighboring patches. This occurs when the traits that were
70 advantageous in an immigrant’s home niche are maladaptive elsewhere. Be-
71 cause environmental change is influenced solely by non-social phenotypes in
72 this model, this change of invasibility affects cooperators and defectors equally.
73 Here again, however, populations containing a greater number of cooperators
74 may have an advantage. The greater number of individuals that emigrate from
75 these larger populations allow them to “export” their niche—and thus reduce
76 the fitness of neighboring competitors—at a higher rate. We explore whether
77 the range expansion that this process enables provides additional opportunities
78 for cooperation to hitchhike.

79 Finally, we demonstrate how *negative* niche construction, where populations
80 change their environment in ways that reduce fitness, can further support
81 cooperation. Even though the niche construction process creates selective
82 feedbacks, we would expect the magnitude of these feedbacks to decrease as
83 populations evolve. Once individuals can no longer gain adaptations that
84 compensate for the costs of cooperation, they are then outcompeted by non-
85 cooperators. However if populations construct their environment in a way
86 which decreases fitness, cooperation can still hitchhike when this change also
87 creates the opportunity to gain compensatory adaptations.

88 **Materials and Methods**

89 We build upon the model described in Hankshaw and Kerr (2015), in which co-
90 operators and defectors compete and evolve in a spatially-structured metapop-
91 ulation of populations. Each of these populations grows to carrying capacity,
92 mutates, and migrates to neighboring patches. During this process, popula-
93 tions adapt to their local environments. In our extended model, we allow
94 these individuals to modify their local environment, and these modifications
95 feed back to affect selection.

96 **Model Description**

97 Our simulated environment consists of N^2 patches arranged as an $N \times N$
98 lattice (see [Table 1](#) for model parameters and their values), where each patch
99 supports a population of zero or more individuals. Each individual in the
100 population has a genotype, which is an ordered list of $L + 1$ integers (loci).
101 The first L loci are *adaptive loci*, and are each occupied by a 0 or an integer
102 from the set $A = \{1, \dots, a_{max}\}$, where a_{max} is the number of potential alleles.
103 At each of these loci, the presence of a non-zero allele represents an adaptation
104 to the environment that confers fitness benefit δ . A binary allele at locus $L + 1$
105 determines whether or not that individual is a cooperator. Individuals with
106 allelic state 1 at this locus are cooperators, carrying a cost c , while individuals
107 with allelic state 0 are defectors. When $\delta \geq c$, an adapted cooperator recoups
108 the cost of cooperation.

109 Organisms also influence their environment, which can feed back to influence

110 selection. We model this as a form of frequency dependent selection. Specif-
 111 ically, the selective value of adaptive allele a at locus l increases with the
 112 number of individuals in the population that have allele $a - 1$ (modulo a_{max})
 113 at locus $l - 1$ (modulo L). The slope of this increase is ϵ , which specifies the
 114 intensity of niche construction. As a consequence of this form of frequency
 115 dependence, genotypes with sequentially-increasing allelic states will tend to
 116 evolve. Because mutations are random, as described later, each population
 117 will evolve sequences that start with different allelic states. These different
 118 sequences represent the unique niches constructed by populations. Under this
 119 model, the fitness of an individual with genotype g in population P is:

$$W_g = z + \delta \sum_{l=1}^L I_A(a_{g,l}) + \epsilon \sum_{l=1}^L n(a_{g,l}) + ca_{g,L+1} \quad (1)$$

120 where z is a baseline fitness, $a_{g,l}$ represents the allelic state of genotype g at
 121 locus l , L is the number of adaptive loci, and c is the cost of the cooperative
 122 allele. The function I_A indicates whether allelic state y is in A (i.e., it is non-
 123 zero). The function $n(a_{g,l})$ gives the number of individuals in the population
 124 with allelic state at the previous locus equal to one less than that at the focal
 125 locus $a_{g,l}$, or:

$$n(a_{g,l}) = \sum_{i \in P} I_{a_{g,l}}(1 + a_{\gamma(i), \{1+(l-2 \pmod L)\} \pmod{a_{max}}}) \quad (2)$$

126 Here, $I_x(y)$ indicates whether the allelic state y matches allelic state x (1) or
 127 not (0), and $\gamma(j)$ is the genotype of individual j . Note that the selective value

128 of an allele at the first locus is affected by the allelic state at locus L .
 129 Cooperators produce a public good that is equally accessible to all members
 130 of the population. This public good increases the carrying capacity at that
 131 patch, allowing the population to reach greater density. This benefit increases
 132 linearly with the proportion of cooperators. Thus, if p is the proportion of
 133 cooperators in a population at the beginning of a growth cycle, then that
 134 population reaches the following size during the growth phase:

$$S(p) = S_{min} + p(S_{max} - S_{min}) \quad (3)$$

135 The function $S(p)$ reflects the benefit of public good production. A population
 136 composed entirely of defectors reaches size S_{min} , while one composed entirely
 137 of cooperators reaches size S_{max} (with $S_{max} \geq S_{min}$). During growth, indi-
 138 viduals compete for inclusion in the resulting population. The composition of
 139 population P with cooperator proportion p after growth is multinomial with
 140 parameters and $S(p)$ and $\{\pi_1, \pi_2, \dots, \pi_{|P|}\}$, where:

$$\pi_i = \frac{W_{\gamma(i)}}{\sum_{j \in P} W_{\gamma(j)}} \quad (4)$$

141 Here, $\gamma(i)$ is the genotype of individual i , and $W_{\gamma(i)}$ is its fitness (see Equation
 142 1). π_i therefore reflects that an individual's ability to persist is proportional
 143 to its fitness relative to others'.

144 For simplicity, we apply mutations after population growth. Mutations occur
 145 independently at each locus and cause the allelic state to change. Mutations

146 occur at each adaptive locus at rate μ_a , and cause a new allelic state to be
 147 chosen at random from the set $\{0\} \cup A$. At the binary cooperation locus,
 148 mutations occur at rate μ_c . These mutations flip the allelic state, causing
 149 cooperators to become defectors and vice versa. Therefore, the probability
 150 that genotype g mutates into genotype g' is given by:

$$\tau_{g \rightarrow g'} = \mu_a^{H_a(g, g')} (1 - \mu_a)^{\{L - H_a(g, g')\}} \mu_c^{H_c(g, g')} (1 - \mu_c)^{\{1 - H_c(g, g')\}} \quad (5)$$

151 where $H_a(g, g')$ and $H_c(g, g')$ are the Hamming distances between genotypes g
 152 and g' at the cooperation locus and adaptive loci, respectively. The Hamming
 153 distance is the number of loci at which allelic states differ (Hamming, 1950).
 154 Because we define no inherent relationship among alleles, each of the $a_{max} + 1$
 155 allelic states is equally likely to arise via mutation at a given locus.

156 After mutation, individuals emigrate to an adjacent patch at rate m . The
 157 destination patch is randomly chosen with uniform probability from the source
 158 patch's Moore neighborhood, which is composed of the nearest 8 patches on the
 159 lattice. Because the metapopulation lattice has boundaries, patches located
 160 on an edge have smaller neighborhoods.

161 Metapopulations are initiated in a state that follows an environmental change.
 162 First, populations are seeded at all patches with cooperator proportion p_0 and
 163 grown to density $S(p_0)$. An environmental challenge is then introduced, which
 164 subjects the population to a bottleneck. For each individual, the probability
 165 of survival is μ_t , which represents the likelihood that a mutation occurs that
 166 confers tolerance. Survivors are chosen by binomial sampling. Because indi-

167 individuals have not yet adapted to this new environment, the allelic state of each
168 individual's genotype is set to 0 at each adaptive locus ($\forall i \in P, l \in \{1, \dots, L\} :$
169 $a_{\gamma(i),l} = 0$). Following initialization, simulations are run for T cycles, where
170 each discrete cycle consists of growth, mutation, and migration. At the end
171 of each cycle, populations are thinned to allow for growth in the next cycle.
172 The individuals that remain are chosen by binomial sampling, where each
173 individual persists with probability d , regardless of allelic state.

174 **Source Code and Software Environment**

175 The simulation software and configurations for the experiments reported are
176 available online (Us, 2015). Simulations used Python 3.4.0, NumPy 1.9.1,
177 Pandas 0.15.2 (McKinney, 2010), and NetworkX 1.9.1 (Hagberg *et al.*, 2008).
178 Data analyses were performed with R 3.1.3 (R Core Team, 2015).

179 **Results**

180 **Niche construction prolongs cooperator survival**

181 “cooperator presence” greater in NC runs than with no-NC (GNH)

182 **Negative Niche construction helps**

183 Runs with wraparound persist indefinitely, while those without do not.

184 **NC Prevents/Limits Defector invasion**

185 **NC Enables Cooperator Spread by “exporting” environ-**
186 **ment**

187 **Discussion**

- 188 • summary of results
- 189 • similarities/differences from previous work
 - 190 – Schwilk and Kerr (2002)
 - 191 – 10.1073/pnas.0812644106
- 192 • public goods as niche construction
- 193 • future QS or other environmental sensing
- 194 • Facultative cooperation
 - 195 – Rodrigues (2012)
 - 196 – Dumas and Kümmerli (2010)
 - 197 – Kümmerli and Brown (2010)
 - 198 – Darch/Diggle
 - 199 – QS?
 - 200 – Environmental Sensing? - (Koestler and Waters, 2014, Bernier et
 - 201 al. (2011))
- 202 • Negative Niche construction as a stragegy? - would those that create this
- 203 constant pressure (L=5, A=6) do better than those that do not (L=5,

204 A=5)?

205 Niche construction and selective feedbacks Niche construction and other social
206 interactions

207 **Public Goods**

208 TODO: merge this in with the “Cooperative Niche Construction” section
209 **TEST**

210 By their very nature, public goods benefit populations by making their environ-
211 ment more hospitable. For example, bacteria produce extracellular products
212 that find soluble iron (Griffin *et al.*, 2004), digest large proteins (Diggle *et al.*,
213 2007; Darch *et al.*, 2012), and reduce the risk of predation (Cosson *et al.*, 2002),
214 among many others (West *et al.*, 2007a). While many studies have explored
215 how the environment affects the evolution of cooperative behaviors, relatively
216 few have examined how those behaviors affect the environment and the result-
217 ing change in evolutionary trajectories. Lehmann (2007) demonstrated analyti-
218 cally that when niche construction act benefits future generations, cooperation
219 is favored due to reduced competition among kin. When rate-benefitting and
220 yield-benefitting altruistic acts co-evolve, Van Dyken and Wade (2012) showed
221 that “reciprocal niche construction”, where the selective feedbacks produced
222 by one act benefitted the other, can lead to increased selection for both traits.
223 While these studies have focused on the niche constructing effects of cooper-
224 ation, we instead focus our attention here on how niche construction enables
225 cooperators to escape defection by hitchhiking along with non-social traits.

226 Primacy/Recency

227 In our model, alterations to the environment were immediately echoed by
228 changes in selection. However, decoupling the timescales on which these pro-
229 cesses occur can have substantial effects (Laland *et al.*, 1996). By integrating
230 past allelic states into Equation 1, we can begin to explore how the cumulative
231 effects of niche construction affect the creation of non-social adaptive oppor-
232 tunities and the benefits that they offer cooperation. Here, how these past
233 allelic states are integrated will play an important role. For example, when
234 the effects of earlier generations are weighted more heavily, the influence of
235 migration may be diminished. While this will reduce the threat of emigration
236 by defectors, cooperator populations will also be less effective at exporting
237 their niche.

238 Cooperative Niche Construction

239 While our focus for this work has been on the eco-evolutionary feedbacks cre-
240 ated by non-social traits, it would also be interesting to explore how this system
241 is affected by the timescale at which carrying capacity at a given patch is in-
242 creased by public goods. In natural settings, a multitude of factors including
243 protein durability (Brown, 2007; Kümmerli and Brown, 2010), diffusion (Al-
244 lison, 2005; Driscoll and Pepper, 2010), and resource availability (Zhang and
245 Rainey, 2013; Ghoul *et al.*, 2014) influence both the rate and the degree to
246 which public goods alter the environment (and thereby selection). Lehmann
247 (2007) demonstrated that a cooperative, niche constructing behavior can be fa-

vored when it only affected selection for future generations, thus reducing the potential for competition among contemporary kin. The evolutionary inertia that this creates, however, may ultimately work against cooperators. When public good accumulates in the environment, cooperators must reduce their investment in production to remain competitive (Kümmerli and Brown, 2010).
TODO: wrap up. Facultative cooperation requires sensing.

Host-Symbiont

In many instances of cooperation, the environment is itself a biological entity, which can produce additional evolutionary feedbacks. As the host population changes, so too will selection on their symbiont populations. Here, evolutionary outcomes depend greatly on the degree of shared interest between the host and symbiont. For example, the cooperative production of virulence factors by the human pathogen *P. aeruginosa* in lung infections is harmful to those with cystic fibrosis (Harrison, 2007). Conversely, cooperative light production by *A. fischeri* is vital for the survival of its host, the Hawaiian bobtail squid (Ruby, 1996). It was recently argued that incorporating the effects of niche construction is critical for improving our understanding of viral evolution (Hamblin *et al.*, 2014) and evolution in co-infecting parasites (Hafer and Milinski, 2015). Incorporating host dynamics, co-evolution, and the feedbacks that they produce into models is likely to be equally important for gaining an understanding of how cooperative behaviors evolve in these host-symbiont settings.

269 Acknowledgments

- 270 • TODO: Organizers?
- 271 • TODO: lab comments

272 We thank Anuraag Pakanati for assistance with simulations. This material is
273 based upon work supported by the National Science Foundation Postdoctoral
274 Research Fellowship in Biology under Grant No. 1309318 (to BDC) and under
275 Cooperative Agreement No. DBI-0939454. Any opinions, findings, and conclu-
276 sions or recommendations expressed in this material are those of the authors
277 and do not necessarily reflect the views of the National Science Foundation.
278 Computational resources were provided by an award from Google (to BDC
279 and BK).

280 **Figures**

281 **Figure 1**

282 **Figure 1A**





Figure 1: Proportion of cooperators over time when there are no opportunities for non-social adaptation

283 **Figure 1B**

284 Proportion of cooperators over time with non-social adaptation

285 **Figure 1C**

286 Proportion of cooperators over time with non-social adaptation and niche con-



287 struction

288 **Figure 2**

289 Plot showing integral for baseline values, double adaptive benefit (no NC,
290 epsilon=0), no negative NC (L=5, A=5), Extreme negative niche construction
291 (L=1, A=6)

292 **Figure 3**

293 Mean fitness over time for the treatments shown in Figure 2

294 **Figure 3B - Fitness with double delta, no epsilon**




Figure 2: Grand mean Fitness of cooperators and defectors, double delta, no epsilon

295 **Figure 4**

296 Cooperators invade from single population

297 **Figure 5**

298 Defectors are kept at bay

299 **Figure 6**

300 **Figure 6A - Effect of Migration Rate (m)**

301 **Figure 6B - Effect of Public Good Benefit ($S_{\max}-S_{\min}$)**

Table 1: Model parameters and their value

Parameter	Description	Base Value
N^2	Number of metapopulation sites	625
L	Number of adaptive loci	5
a_{max}	Number of alleles	6
δ	Fitness benefit, nonzero alleles	0.3
c	Production cost	0.1
ϵ	Fitness benefit, sequential alleles	0.00015
z	Baseline fitness	1
S_{min}	Minimum population size	800
S_{max}	Maximum population size	2000
μ_c	Mutation rate (cooperation)	10^{-5}
μ_a	Mutation rate (adaptation)	10^{-5}
m	Migration rate	0.05
p_0	Initial cooperator proportion	0.5
μ_t	Mutation rate (tolerance to new stress)	10^{-5}
T	Number of simulation cycles	1000
d	Population dilution factor	0.1

References

- Allison, S.D. 2005. Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. *Ecology Letters*, **8**: 626–635.
- Asfahl, K.L., Walsh, J., Gilbert, K. and Schuster, M. 2015. Non-social adaptation defers a tragedy of the commons in *Pseudomonas aeruginosa* quorum sensing. *The ISME Journal*, doi: [10.1038/ismej.2014.259](https://doi.org/10.1038/ismej.2014.259).
- Bernier, S.P., Ha, D.-G., Khan, W., Merritt, J.H.M. and O'Toole, G.A. 2011. Modulation of *Pseudomonas aeruginosa* surface-associated group behaviors by individual amino acids through c-di-GMP signaling. *Research in Microbiology*, **162**: 680–688.
- Brown, F., Sam P. AND Taddei. 2007. The durability of public goods changes the dynamics and nature of social dilemmas. *PLoS ONE*, **2**: e593. Public Library of Science.
- Brown, S.P. and Johnstone, R.A. 2001. Cooperation in the dark: Signalling and collective action in quorum-sensing bacteria. *Proceedings of the Royal Society of London B: Biological Sciences*, **268**: 961–965.
- Cosson, P., Zulianello, L., Join-Lambert, O., Faurisson, F., Gebbie, L. and Benghezal, M.*et al.* 2002. *Pseudomonas aeruginosa* virulence analyzed in a dictyostelium discoideum host system. *Journal of Bacteriology*, **184**: 3027–3033.
- Dandekar, A.A., Chugani, S. and Greenberg, E.P. 2012. Bacterial quorum

325 sensing and metabolic incentives to cooperate. *Science*, **338**: 264–266.

326 Darch, S.E., West, S.A., Winzer, K. and Diggle, S.P. 2012. Density-dependent
327 fitness benefits in quorum-sensing bacterial populations. *Proceedings of the*
328 *National Academy of Sciences*, **109**: 8259–8263.

329 Diggle, S.P., Griffin, A.S., Campbell, G.S. and West, S.A. 2007. Cooperation
330 and conflict in quorum-sensing bacterial populations. *Nature*, **450**: 411–414.

331 Driscoll, W.W. and Pepper, J.W. 2010. Theory for the evolution of diffusible
332 external goods. *Evolution*, **64**: 2682–2687.

333 Fletcher, J.A. and Doebeli, M. 2009. A simple and general explanation for the
334 evolution of altruism. *Proceedings of the Royal Society B: Biological Sciences*,
335 **276**: 13–19.

336 Foster, K., Shaulsky, G., Strassmann, J., Queller, D. and Thompson, C. 2004.
337 Pleiotropy as a mechanism to stabilize cooperation. *Nature*, **431**: 693–696.
338 Nature Publishing Group.

339 Gardner, A. and West, S.A. 2010. Greenbeards. *Evolution*, **64**: 25–38.

340 Ghoul, M., West, S.A., Diggle, S.P. and Griffin, A.S. 2014. An experimental
341 test of whether cheating is context dependent. *Journal of Evolutionary Biology*,
342 **27**: 551–556.

343 Griffin, A.S., West, S.A. and Buckling, A. 2004. Cooperation and competition
344 in pathogenic bacteria. *Nature*, **430**: 1024–1027.

345 Hafer, N. and Milinski, M. 2015. When parasites disagree: Evidence
346 for parasite-induced sabotage of host manipulation. *Evolution*, doi:

347 [10.1111/evo.12612](https://doi.org/10.1111/evo.12612).

348 Hagberg, A.A., Schult, D.A. and Swart, P.J. 2008. Exploring network struc-
349 ture, dynamics, and function using NetworkX. In: *Proceedings of the 7th*
350 *Python in Science Conference (SciPy2008)*, pp. 11–15.

351 Hamblin, S.R., White, P.A. and Tanaka, M.M. 2014. Viral niche construction
352 alters hosts and ecosystems at multiple scales. *Trends in Ecology & Evolution*,
353 **29**: 594–599.

354 Hamilton, W.D. 1964. The genetical evolution of social behaviour I & II.
355 *Journal of Theoretical Biology*, **7**: 1–52.

356 Hamming, R.W. 1950. Error detecting and error correcting codes. *Bell System*
357 *Technical Journal*, **29**: 147–160.

358 Hankshaw, S. and Kerr, B. 2015. Thumbs up!: Hitching a ride on the evolu-
359 tionary superhighway. *Unknown Journal*, **0**: 0–0.

360 Hardin, G. 1968. The tragedy of the commons. *Science*, **162**: 1243–1248.

361 Harrison, F. 2007. Microbial ecology of the cystic fibrosis lung. *Microbiology*,
362 **153**: 917–923.

363 Koestler, B.J. and Waters, C.M. 2014. Bile acids and bicarbonate inversely
364 regulate intracellular cyclic di-GMP in *Vibrio cholerae*. *Infection and Immu-*
365 *nity*, **82**: 3002–3014.

366 Kuzdzal-Fick, J.J., Fox, S.A., Strassmann, J.E. and Queller, D.C. 2011. High
367 relatedness is necessary and sufficient to maintain multicellularity in *Dic-*
368 *tyostelium*. *Science*, **334**: 1548–1551.

369 Kümmerli, R. and Brown, S.P. 2010. Molecular and regulatory properties of
370 a public good shape the evolution of cooperation. *Proceedings of the National*
371 *Academy of Sciences*, **107**: 18921–18926.

372 Laland, K.N., Odling-Smee, F.J. and Feldman, M.W. 1996. The evolutionary
373 consequences of niche construction: A theoretical investigation using two-locus
374 theory. *Journal of Evolutionary Biology*, **9**: 293–316.

375 Lehmann, L. 2007. The evolution of trans-generational altruism: Kin selec-
376 tion meets niche construction. *Journal of Evolutionary Biology*, **20**: 181–189.
377 Blackwell Publishing Ltd.

378 McKinney, W. 2010. Data structures for statistical computing in Python. In:
379 *Proceedings of the 9th Python in Science Conference* (S. van der Walt and J.
380 Millman, eds), pp. 51–56.

381 Morgan, A.D., Quigley, B.J.Z., Brown, S.P. and Buckling, A. 2012. Selection
382 on non-social traits limits the invasion of social cheats. *Ecology Letters*, **15**:
383 841–846.

384 Nadell, C.D., Foster, K.R. and Xavier, J.B. 2010. Emergence of spatial struc-
385 ture in cell groups and the evolution of cooperation. *PLoS Computational*
386 *Biology*, **6**: e1000716.

387 Nowak, M.A. 2006. Five rules for the evolution of cooperation. *Science*, **314**:
388 1560–1563.

389 R Core Team. 2015. *R: A language and environment for statistical computing*.
390 Vienna, Austria: R Foundation for Statistical Computing.

391 Ruby, E.G. 1996. Lessons from a cooperative, bacterial-animal association:
 392 The *Vibrio fischeri*–*Euprymna scolopes* light organ symbiosis. *Annual Review*
 393 *of Microbiology*, **50**: 591–624.

394 Schwilk, D.W. and Kerr, B. 2002. Genetic niche-hiking: An alternative expla-
 395 nation for the evolution of flammability. *Oikos*, **99**: 431–442.

396 Sinervo, B., Chaine, A., Clobert, J., Calsbeek, R., Hazard, L. and Lancaster,
 397 L.*et al.* 2006. Self-recognition, color signals, and cycles of greenbeard mutu-
 398 alism and altruism. *Proceedings of the National Academy of Sciences*, **103**:
 399 7372–7377.

400 Us. 2015. Name for data and code.

401 Van Dyken, J.D. and Wade, M.J. 2012. Origins of altruism diversity II: Run-
 402 away coevolution of altruistic strategies via “reciprocal niche construction”.
 403 *Evolution*, **66**: 2498–2513.

404 Veelders, M., Brückner, S., Ott, D., Unverzagt, C., Mösch, H.-U. and Essen,
 405 L.-O. 2010. Structural basis of flocculin-mediated social behavior in yeast.
 406 *Proceedings of the National Academy of Sciences*, **107**: 22511–22516.

407 Waite, A.J. and Shou, W. 2012. Adaptation to a new environment allows coop-
 408 erators to purge cheaters stochastically. *Proceedings of the National Academy*
 409 *of Sciences*, **109**: 19079–19086.

410 West, S.A., Diggle, S.P., Buckling, A., Gardner, A. and Griffin, A.S. 2007a.
 411 The social lives of microbes. *Annual Review of Ecology, Evolution, and Sys-*
 412 *tematics*, **38**: 53–77.

- 413 West, S.A., Griffin, A.S. and Gardner, A. 2007b. Evolutionary explanations
414 for cooperation. *Current Biology*, **17**: R661–R672.
- 415 Zhang, X.-X. and Rainey, P.B. 2013. Exploring the sociobiology of pyoverdin-
416 producing *Pseudomonas*. *Evolution*, **67**: 3161–3174.