

1 Negative Niche Construction Favors the
2 Evolution of Cooperation

3
4 **Abstract**

5 TODO

6 **Introduction**

7 Cooperative behaviors are common across all branches of the tree of life. In-
8 sects divide labor within their colonies, plants and soil bacteria exchange es-
9 sential nutrients, birds care for others' young, and the trillions of cells in the
10 human body coordinate to provide vital functions. Each instance of cooper-
11 ation presents an evolutionary challenge: How can individuals that sacrifice
12 their own well-being to help others avoid subversion by those that do not? Over
13 time, we would expect these *defectors* to rise in abundance at the expense of
14 others, eventually driving cooperators—and perhaps the entire population—to
15 extinction.

16 Several factors can prevent this *tragedy of the commons* (Hamilton, 1964;
17 Nowak, 2006; West *et al.*, 2007). One such factor involves non-random so-
18 cial interaction, in which cooperators benefit more from the cooperative act
19 than defectors. This can occur when cooperators are clustered together in
20 spatially-structured populations (Fletcher and Doebeli, 2009; Nadell *et al.*,
21 2010; Kuzdzal-Fick *et al.*, 2011) or when cooperators use communication
22 (Brown and Johnstone, 2001; Darch *et al.*, 2012) or other cues (Sinervo *et al.*,
23 2006; Gardner and West, 2010; Veelders *et al.*, 2010) to cooperate condition-
24 ally with kin. Cooperation can also be bolstered by pleiotropic connections to
25 personal benefits (Foster *et al.*, 2004; Dandekar *et al.*, 2012) or through associ-
26 ation with alleles encoding self-benefitting traits (Asfahl *et al.*, 2015). In these
27 cases, the alleles may provide private benefits that are completely independent
28 from the public benefits of cooperation. In asexual populations of cooperators
29 and defectors, this sets the stage for an “adaptive race” in which both types
30 vie for the first highly beneficial adaptation (Waite and Shou, 2012; Morgan
31 *et al.*, 2012). The tragedy of the commons can be deferred if a cooperator, by
32 chance, wins the adaptive race.

33 Hammarlund *et al.* (2015) recently showed that in spatially-structured pop-
34 ulations, the “Hankshaw effect” can give cooperators a substantial leg up on
35 defectors in an adaptive race. This advantage is reminiscent of Sissy Han-
36 kshaw, a fictional character in Tom Robbins’ *Even Cowgirls Get the Blues*,
37 whose oversized thumbs—which were otherwise an impairment—made her a
38 prolific hitchhiker. Similarly, cooperation is costly, but it increases local pop-
39 ulation density. As a result, cooperators are more likely to acquire beneficial

40 mutations. By hitchhiking along with these adaptations, cooperation can then
41 rise in abundance. Nevertheless, this advantage is fleeting. As soon as the
42 opportunities for adaptation are exhausted, cooperators are once again at a
43 selective disadvantage against equally-adapted defectors that arise via muta-
44 tion. However, Hammarlund et al. (2015) also demonstrated that cooperation
45 can be maintained indefinitely when frequent environmental changes produce
46 a steady stream of new adaptive opportunities. Although organisms typically
47 find themselves in dynamic environments, the nature and frequency of these
48 changes might not ensure long-term cooperator survival.

49 Importantly, however, organisms do more than simply experience changing
50 environments passively. Through their activities, their interactions with oth-
51 ers, and even their deaths, organisms constantly modify their environment.
52 These changes can produce evolutionary feedback loops in which environmen-
53 tal change alters selection, which, in turn, alters the distribution of types and
54 their corresponding influence on the environment (Odling-Smee *et al.*, 2003).
55 The nature of this feedback can have dramatic evolutionary consequences. One
56 critical distinction is whether the constructing type or some other type is most
57 adapted in the resulting environment. Under positive niche construction, selec-
58 tion favors the constructor, and evolution stagnates as this type fixes. Under
59 negative niche construction, selection favors a type other than the constructor.
60 In this latter case, populations find themselves continually chasing beneficial
61 mutations as their adaptive landscape perpetually shifts.

62 Here, we show that the selective feedbacks that result from niche construction
63 can maintain cooperation indefinitely. We find that it is specifically negative

64 niche construction that is responsible for this result because of the adaptive
65 opportunities that it produces. Furthermore, we show that the rate at which
66 niche construction occurs is also crucial. These results indicate that cooper-
67 ators can ensure their survival when they play an active role in their own
68 evolution.

69 **Methods**

70 Building upon Hammarlund et al. (2015), we develop an individual-based
71 model in which cooperators and defectors evolve and compete in a population
72 of subpopulations (i.e., a metapopulation). Through mutations, individuals
73 gain adaptations to their environment, which increase reproductive fitness,
74 and allow those lineages to rise in abundance. Migration among neighboring
75 subpopulations allows more successful lineages to spread.

76 In our expanded model, subpopulations additionally modify their local envi-
77 ronment. As this process occurs, environmental changes feed back to affect
78 selection. We explore how niche construction affects this process of adaptation
79 and whether cooperation can be maintained because of selective feedbacks.

80 **Model Description**

81 **Individual Genotypes and Adaptation**

82 Each individual has a haploid genome with $L + 1$ loci (see [Table 1](#) for model
83 parameters and their values). Different alleles at each locus are represented by

84 different integers. A binary allele at the first locus (here, locus zero) determines
 85 whether that individual is a cooperator (1), which carries fitness cost c , or a
 86 defector (0). Cooperation is independent from adaptation to the environment.
 87 The remaining L loci are *adaptive loci*, and are each occupied by 0 or a value
 88 from the set $\{1, 2, \dots, A\}$. Allele 0 represents a lack of adaptation, while
 89 a non-zero allele represents one of the A possible adaptations at that locus.
 90 These non-zero alleles signify adaptations to the external environment that
 91 are not affected by other individuals or the local niche. Adaptations confer
 92 a fitness benefit δ , regardless of which non-zero allele is present. We assume
 93 $\delta > c$, which allows a minimally adapted cooperator to recoup the cost of
 94 cooperation and gain a fitness advantage.

95 Niche Construction and Selective Feedbacks

96 Individual fitness is also affected by the current state of the local environment.
 97 Here, we represent the “niche” implicitly based on the allelic states present in
 98 the subpopulation. As allelic states change, subpopulations alter aspects of
 99 their environment, creating a unique niche.

100 Niche construction takes the form of density dependent selection, and indi-
 101 viduals evolve to better match their niche by a second form of adaptation.
 102 Specifically, the selective value of adaptive allele a at locus l increases with
 103 the number of individuals in the subpopulation that have allele $a - 1$ at locus
 104 $l - 1$. As a consequence, genotypes with sequentially increasing allelic states
 105 will tend to evolve. We treat both adaptive loci and allelic states as “circular”:

106 the selective value of an allele at locus 1 is affected by the allelic composition
 107 of the subpopulation at locus L . Similarly, the selective value of allele 1 at
 108 any locus increases with the number of individuals carrying allele A at the
 109 previous locus. This circularity is represented by the function $\beta(x, X)$, which
 110 gives the integer that is below an arbitrary value x in the set $\{1, 2, \dots, X\}$:

$$\beta(x, X) = \text{mod}_X(x - 2 + X) + 1 \quad (1)$$

111 Here, $\text{mod}_X(x)$ is the integer remainder when dividing x by X . The selective
 112 value of adaptive allele a at locus l is increased by ϵ for each individual in the
 113 subpopulation that has allele $\beta(a, A)$ at locus $\beta(l, L)$. Thus, ϵ specifies the
 114 intensity of niche construction.

115 Consider a genotype g with the allelic state at locus l given by $a_{g,l}$; the fitness
 116 of an individual with this genotype is defined as:

$$W_g = z - \underbrace{ca_{g,0}}_{\text{cost of cooperation}} + \underbrace{\delta \sum_{l=1}^L I(a_{g,l})}_{\text{adaptation to external environment}} + \underbrace{\epsilon \sum_{l=1}^L n(\beta(a_{g,l}, A), \beta(l, L))}_{\text{adaptation to constructed environment}} \quad (2)$$

117 where z is a baseline fitness and $I(a)$ indicates whether a given adaptive allele
 118 is non-zero:

$$I(a) = \begin{cases} 1 & \text{if } a \in \{1, 2, \dots, A\} \\ 0 & \text{otherwise} \end{cases} \quad (3)$$

Thus, an individual's fitness is determined both by adaptations to the external environment (δ) and adaptations to its constructed environment (ϵ). [Figure 1](#) illustrates the effects of these two components.

Population Growth and the Benefit of Cooperation

Cooperation enables a subpopulation to reach a greater density. This benefit affects all individuals equally and accumulates linearly with the proportion of cooperators in the subpopulation. If p is the proportion of cooperators present at the beginning of a growth cycle, then that subpopulation reaches the following size:

$$S(p) = S_{min} + p(S_{max} - S_{min}) \quad (4)$$

During growth, individuals compete through differential reproduction. Each individual's probability of success is determined by its fitness. The composition of a subpopulation with size P and cooperator proportion p after growth is multinomial with parameters $S(p)$ and $\{\pi_1, \pi_2, \dots, \pi_P\}$, where:

$$\pi_i = \frac{W_{\gamma(i)}}{\sum_{j=1}^P W_{\gamma(j)}} \quad (5)$$

Here, $W_{\gamma(i)}$ is the fitness of an individual i with genotype $\gamma(i)$ (see [Equation 2](#)). The value π_i represents an individual's reproductive fitness relative to others in the subpopulation.

135 **Mutation**

136 For simplicity, we apply mutations after growth. Mutations occur indepen-
137 dently at each locus and cause an allelic state change. At the binary coopera-
138 tion locus, mutations occur at rate μ_c . These mutations flip the allelic state,
139 causing cooperators to become defectors and vice versa. Mutations occur at
140 rate μ_a at each adaptive locus. These mutations replace the existing allele
141 with a random selection from the set $\{0\} \cup \{1, 2, \dots, A\}$. Because mutations
142 are stochastic, the allelic sequences that evolve depend on which allele arises
143 first and at which locus.

144 **Migration**

145 Populations are composed by N^2 patches arranged as an $N \times N$ lattice, where
146 each patch can support a subpopulation. After mutation, individuals emigrate
147 to an adjacent patch with probability m . During each migration event, a
148 single destination patch is randomly chosen with uniform probability from
149 each source patch's Moore neighborhood, which is composed of the nearest 8
150 patches on the lattice. Because the population lattice has boundaries, patches
151 located on the periphery have smaller neighborhoods.

152 **Population Initialization and Simulation**

153 At the beginning of each simulation, subpopulations are seeded at all patches
154 with cooperator proportion p_0 and grown to density $S(p_0)$. An environmental
155 challenge is then introduced, which subjects all subpopulations to a bottleneck.

156 For each individual, the probability of survival is μ_t , which represents the
157 likelihood that tolerance arises via mutation. Because individuals have not yet
158 adapted to this new environment, the allelic state of each individual's genotype
159 is 0 at each adaptive locus. Following initialization, simulations are run for T
160 cycles, where each discrete cycle consists of subpopulation growth, mutation,
161 migration, and dilution. Dilution thins the population to support growth in
162 the next cycle. Each individual remains with probability d , regardless of allelic
163 state.

164 Simulation Source Code and Software Dependencies

165 The simulation software and configurations for the experiments reported are
166 available online.¹ Simulations used Python 3.4, NumPy 1.9.1, Pandas 0.15.2
167 (McKinney, 2010), and NetworkX 1.9.1 (Hagberg *et al.*, 2008). Data analy-
168 ses were performed with R 3.1.3 (R Core Team, 2015). Reported confidence
169 intervals were estimated by bootstrapping with 1000 resamples.

170 Results

171 Using the model described in the previous section, we perform simulations
172 that follow the evolution of cooperation in a population consisting of subpopu-
173 lations that are connected by spatially-limited migration. Individuals compete
174 in these subpopulations by gaining a limited number of adaptations that con-

¹These materials will be made public at the time of publication, and a reference will be placed here.

175 fer fitness benefits. While cooperation does not directly affect the selective
 176 value of these adaptations, cooperation can have indirect effects on the adap-
 177 tive process. Specifically, cooperation increases subpopulation density. As a
 178 result, larger subpopulations of cooperators experience more mutational op-
 179 portunities to gain adaptations. Cooperation can hitchhike along with these
 180 adaptations, which compensate for the cost of cooperation. During this process,
 181 subpopulations alter their local environments, which feeds back to influence
 182 selection. Here, we explore how niche construction affects the evolution of
 183 cooperation in the simulation environment defined by the parameter values
 184 listed in [Table 1](#).

185 **Cooperation Persists with Niche Construction**

186 Without any opportunity for adaptation ($L = 0$), cooperators are swiftly elim-
 187 inated in competition with defectors ([Figure 2A](#)). Despite an initial lift in
 188 cooperator abundance due to increased productivity, the cost of cooperation
 189 becomes disadvantageous as migration mixes the initially isolated subpopula-
 190 tions. When populations can adapt to the external environment ($L = 5$), but
 191 niche construction is absent ($\epsilon = 0$), cooperators are maintained only tran-
 192 siently ([Figure 2B](#)). Here, larger cooperator subpopulations can more quickly
 193 adapt to their external environment as before. As previously described by
 194 Hammarlund et al. (2015), however, cooperation is subsequently lost once
 195 populations become fully adapted to their environment. Once this has oc-
 196 curred, isogenic defectors (i.e., defectors with identical adaptive loci) arise via

197 mutation and displace cooperation due to their selective advantage. However,
198 when niche construction creates selective feedbacks, cooperation persists in
199 over 2/3 of the replicate populations (Figure 2C). We see in Figure 3A that
200 despite oscillations, cooperation is maintained at high levels in these popula-
201 tions.

202 **Fitness Increases Alone do not Support Persisting Coop-** 203 **eration**

204 In the model, both adaptation and niche construction contribute to an indi-
205 vidual's fitness. To determine whether cooperation is maintained solely due to
206 the larger selective values that result from the contributions of niche construc-
207 tion (ϵ), we performed simulations in which these contributions were removed
208 ($\epsilon = 0$), and we instead increased the fitness benefits conferred by adaptation to
209 the external, non-constructed environment ($\delta = 0.6$). In doing so, we conserva-
210 tively estimate the selective effects of niche construction by supplementing the
211 selective benefits of adaptations to the external environment by the maximum
212 possible selective benefit that results from niche construction. Nevertheless,
213 we find that simply increasing selective values does not enable cooperators to
214 persist (Figure 3B). Niche construction, therefore, plays a decisive role here.

215 Negative Niche Construction is Critical to Cooperator 216 Persistence

217 Negative niche construction can occur in our model due to the selection for
218 sequentially-increasing allelic states and the circular arrangement of these al-
219 leles (see [Figure 1](#)). This occurs when the number of adaptive alleles (A) does
220 not divide evenly into the number of adaptive loci (L). In such a case, any
221 sequence of integers on the circular genome will always contain a break in the
222 sequence; that is, one locus with an allele that is not one less than the allele at
223 the next locus. Given this unavoidable mismatch, any type that has fixed will
224 always favor selection for a new type. However, if this negative niche construc-
225 tion is removed (by setting $L = 5$, $A = 5$), cooperators are again driven extinct
226 after an initial lift in abundance ([Figure 3C](#)). These results indicate that the
227 type of niche construction matters. Specifically, negative niche construction
228 is crucial for maintaining cooperation.

229 Selective Feedbacks Limit Defector Invasion

230 The adaptation resulting from selective feedbacks can limit invasion by de-
231 fectors, which arise either through immigration from neighboring patches or
232 through mutation from a cooperator ancestor. The latter challenge is partic-
233 ularly threatening, as these isogenic defectors are equally adapted, yet do not
234 incur the cost of cooperation. As demonstrated in [Figure 4A](#), these isogenic
235 defectors rapidly spread when introduced at a single patch in the center of an
236 11×11 population of cooperators if mutations do not occur. However, when

237 resident cooperators can gain adaptations via mutation, cooperators evade de-
238 fector invasion in over half of the replicate populations (Figure 4B). Figure
239 5 depicts one such instance where cooperation survived. In that population,
240 defectors quickly began to spread. However, an adaptation arose at a neighbor-
241 ing cooperator population that was more fit. This type spread more quickly,
242 halting defectors and eventually driving them to extinction. Because this adap-
243 tion occurred in a cooperator population, cooperation was able to hitchhike to
244 safety. Figure 4C shows how quickly an adapted cooperator type can invade
245 a population of defectors.

246 The Rate of Niche Construction Matters

247 **TODO:** Sorry, results coming soon!

248 Discussion

249 Despite their negative effects, deleterious traits can rise in abundance due to
250 genetic linkage with other traits that are strongly favored by selection (May-
251 nard Smith and Haigh, 1974). In a process termed the “Hankshaw effect”,
252 Hammarlund et al. (2015) recently demonstrated that cooperation can pro-
253 long its existence by increasing the likelihood of hitchhiking with a beneficial
254 trait. While this process does favor cooperation in the short term, it eventually
255 reaches a dead end; when the opportunities for adaptation are exhausted, and
256 cooperators can no longer hitchhike, they face extinction. In this work, we

257 have considered whether niche construction might serve to perpetually gener-
 258 ate new adaptive opportunities, and thus favor cooperation indefinitely.

259 When niche construction occurs, cooperation can indeed persist (Figure 2C).
 260 In our model, niche construction introduces additional selective effects that
 261 could influence the evolutionary process, leading to a more pronounced Han-
 262 kshaw effect. However, simply raising the fitness benefits conferred by adap-
 263 tations does not prolong cooperation (Figure 3B), which indicates that niche
 264 construction and the selective feedbacks that it produces play a crucial role.

265 Further, we find that it is specifically negative niche construction that main-
 266 tains cooperation (Figure 3C). Here we observe another facet of the Hankshaw
 267 effect: because populations of cooperators are larger, they are better able to
 268 respond to the adaptive opportunities that result from negative niche construc-
 269 tion. Without these adaptive opportunities, adaptation eventually grinds to a
 270 halt. Once this occurs, cooperators face the threat of invasion by isogenic de-
 271 fectors that arise through mutation. Since these defectors are equally adapted
 272 but do not bear the cost of cooperation, they quickly drive cooperators to ex-
 273 tinction. Importantly, because every type constructs an environment in which
 274 a different type is more fit, negative niche construction creates continual adap-
 275 tive opportunities. These opportunities can allow cooperators to resist invasion
 276 by defectors, even when defectors are equally adapted (Figure 4B). It is these
 277 recurring cycles of invasion and adaptation that underlie the oscillations in
 278 cooperator populations that we see in Figure 3A. When stochastic mutations
 279 do not engender these adaptations, defectors invade, and the cycle is broken.

280 **TODO: the rate of niche construction is crucial**

281 In our model, cooperation is orthogonal to niche construction, which allows us
282 to focus on hitchhiking. However, by increasing the size of the local patch, this
283 form of cooperation could itself be seen as a niche constructing behavior. Pre-
284 vious studies have more directly explored how niche construction and coopera-
285 tion interact. Lehmann (2007) showed that cooperation can be favored when
286 niche construction decoupled kin competition from kin selection in spatially-
287 structured populations. Perhaps most similar our work, Van Dyken and Wade
288 (2012) demonstrated that when two negative niche constructing cooperative
289 behaviors co-evolve, selection can increasingly favor these traits, which were
290 disfavored when alone. In that model, “reciprocal niche construction” occurred
291 when the negative feedback created by one strategy positively influenced se-
292 lection on the other, creating a perpetual cycle that maintained both forms of
293 cooperation. Arguably, this can be viewed as an instance of hitchhiking: the
294 currently-maladaptive form of cooperation is maintained by association with
295 the adaptive form. Outside of the context of cooperation, studies have shown
296 that niche construction can allow deleterious alleles to be maintained (Laland
297 *et al.*, 1996, 1999). However, cooperation, especially in competition against
298 equally-adapted defectors, can be considered deleterious.

299 In the model described by Lehmann (2007), the selective feedbacks produced
300 by the cooperative, niche-constructing behavior only affected future genera-
301 tions. Because of this, kin competition was reduced, and cooperation instead
302 benefitted descendants. Other studies, while not focusing on cooperation, have
303 similarly shown that the timescales at which niche construction feedbacks oc-

304 cur can strongly influence evolutionary outcomes (Laland *et al.*, 1996, 1999).
 305 This perspective is likely to be crucial for understanding the evolution of co-
 306 operative behaviors like the production of public goods.

307 For example, bacteria produce a host of extracellular products that scavage
 308 soluble iron (Griffin *et al.*, 2004), digest large proteins (Diggle *et al.*, 2007;
 309 Darch *et al.*, 2012), and reduce the risk of predation (Cosson *et al.*, 2002).
 310 While many studies have focused on how the environment affects the evolu-
 311 tion of these cooperative public goods, relatively few have addressed how the
 312 environmental changes created by public goods feed back to influence evolu-
 313 tion. In these instances, environmental changes are likely to occur on different
 314 timescales than reproduction, which can have profound effects. For exam-
 315 ple, a multitude of factors including protein durability (Brown and Taddei,
 316 2007; Kümmerli and Brown, 2010), diffusion (Allison, 2005; Driscoll and Pep-
 317 per, 2010), and resource availability (Zhang and Rainey, 2013; Ghoul *et al.*,
 318 2014) influence both the rate and the degree to which public goods alter the
 319 environment. While Lehmann (2007) showed that cooperation was favored
 320 when selective feedbacks act over longer timescales, niche construction may in
 321 fact hinder cooperation when selection is more quickly altered. For example,
 322 when public goods accumulate in the environment, cooperators must decrease
 323 production to remain competitive (Kümmerli and Brown, 2010; Dumas and
 324 Kümmerli, 2012). This favors cooperation that occurs facultatively, perhaps
 325 by sensing the abiotic (Bernier *et al.*, 2011; Koestler and Waters, 2014) or
 326 biotic environment (Brown and Johnstone, 2001; Darch *et al.*, 2012).

327 In many instances where cooperation occurs, the environment is itself a biolog-

ical entity, which can introduce additional evolutionary feedbacks. As the host population changes, so too does selection on their symbiont populations. Here, evolutionary outcomes depend greatly on the degree of shared interest between the host and symbiont. For example, the cooperative production of virulence factors by the human pathogen *P. aeruginosa* in lung infections is harmful to hosts with cystic fibrosis (Harrison, 2007). Conversely, cooperative light production by *A. fischeri* is vital for the survival of its host, the Hawaiian bobtail squid (Ruby, 1996). It was recently argued that incorporating the effects of niche construction is critical for improving our understanding of viral evolution (Hamblin *et al.*, 2014) and evolution in co-infecting parasites (Hafer and Milinski, 2015). Incorporating host dynamics, co-evolution, and the feedbacks that they produce into models is likely to be equally important for gaining an understanding of how cooperative behaviors evolve in these host-symbiont settings.

Acknowledgments

We are grateful to Peter Conlin, Sylvie Estrela, Carrie Glenney and Martha Kornelius for helpful comments on the manuscript and to Anuraag Pakanati for assistance with simulations. This material is based upon work supported by the National Science Foundation Postdoctoral Research Fellowship in Biology under Grant No. DBI-1309318 (to BDC) and under Cooperative Agreement No. DBI-0939454 (BEACON STC). Computational resources were provided by an award from Google Inc. (to BDC and BK).

350 **Figures**

351 **Figure 1**




Figure 1: Adaptation to External and Constructed Environments. (A) We start with the case with five loci ($L = 5$) and five non-zero alleles ($A = 5$). All simulations are initialized with a non-adapted genotype with allele 0 at every locus—the genotype on the far left. Random mutation will introduce a non-zero allele, which is expected to increase in frequency. For simplicity, we assume that allele 1 arises at the first locus (in the “12 o’clock” position). The rest of this schematic focuses on niche construction. Every non-zero allele at any locus influences selection at the next locus in the clockwise direction. There is a “mismatch” in this genotype (highlighted by the red sector) because the niche constructed by allele 1 at the first locus favors allele 2 (not 0) at its immediate clockwise neighbor (the second locus). Once the appropriate allele arises, it will be selected. In this case, the genotype $[1,2,0,0,0]$ receives an epsilon effect in addition to the extra delta. The “match” at the first and second locus is highlighted as a green sector. However, now there is a new mismatch (between the second and third locus), which a new round of mutation and selection corrects, and so on. The green sector grows as the red sector ticks clockwise. Importantly, because A divides evenly into L , this genotype can evolve into a perfectly reinforcing sequence $[1,2,3,4,5]$, which enjoys an maximal epsilon increment of fitness of due to its niche construction. (B) The case of negative niche construction is illustrated for the case of five loci ($L = 5$) and six non-zero alleles ($A = 6$). Here we start with a population fixed for the genotype on the far left $[1,2,3,4,5]$. There is a single mismatch in this genotype (highlighted by the red sector) because the niche constructed by allele 5 favors allele 6 (not 1) at its immediate clockwise neighbor. If the fitter mutant $[6,2,3,4,5]$ arises (see next genotype to the right), it will fix. (We note that the strength of selection will drop as its frequency increases). However, now there is a new mismatch in the genotype (highlighted again with a red sector). Thus, we see that correcting one mismatch generates a new mismatch. Thus, this system will never escape its mismatches—the red sector just clicks clockwise around the genome. Indeed, after six (or A) rounds of mismatch correction/generation, we have ended back where we started with the original genotype turned clockwise by one locus. Here, the adaptation to previous niche construction generates further niche construction that leads to novel adaptation.

352 **Figure 2**




Figure 2: Adaptation, Hitchhiking, and the Evolution of Cooperation. The proportion of cooperators present in the population is shown for the duration of simulations. Curves show the average among replicate populations, and shaded areas indicate 95% confidence intervals. Unless otherwise noted, parameter values are listed in [Table 1](#). **(A)** Without any opportunity to adapt (L , the number of adaptive loci, is zero), cooperation is quickly lost. **(B)** When adaptation can occur ($L = 5$), but populations do not alter their environment (ϵ , the intensity of niche construction, is zero), cooperation hitchhikes along with adaptations, allowing cooperators to temporarily rise in abundance before eventually going extinct. **(C)** Niche construction enables cooperation to be maintained indefinitely. In the majority of populations (13/18), cooperation remained the dominant strategy. Individual populations are shown in Figure 3A.


Figure 3

Figure 3: **Niche Construction and the Evolution of Cooperation.** The proportion of cooperators present in each replicate population is shown for the duration of simulations. **(A)** Despite some oscillations, niche construction enables cooperation to be maintained indefinitely in 14 of 18 populations. **(B)** When niche construction is removed and the fitness benefit of adaptation is increased to compensate ($\epsilon = 0$, $\delta = 0.6$), adapted defectors arise and drive cooperators to extinction. **(C)** Without negative niche construction, cooperation is not maintained ($A = 5$).


Figure 4

Figure 4: Niche Construction and Invasion. Curves trace the proportion of cooperators present in the population for the duration of 160 replicate simulations ($T = 1000$). These experiments examine whether a rare cooperator or defector strategy can invade when initiated at a single patch in the center of the population lattice ($N^2 = 121$). Unless otherwise noted, mutations ($\mu_a = 0, \mu_c = 0$) are disabled in these ecological simulations to highlight the dynamics of invasion. The results from simulations where this limitation is removed are shown in Figure S1. **(A)** When cooperators and defectors are isogenic (i.e., both types have stress alleles [1,2,3,4,5]) and mutation cannot occur, rare defectors quickly invade and drive cooperators to extinction due to the cost of cooperation. Defectors were stochastically eliminated in 2 replicate populations. **(B)** However, the adaptive opportunities produced by negative niche construction can allow cooperators to resist invasion by isogenic defectors. Here, cooperation persisted in the majority of populations ($\mu_a = 0.00005$, the base mutation rate). **(C)** We demonstrate that adaptations such as these can enable an cooperator (stress alleles [6,2,3,4,5], see Figure 1) to displace a population of defectors when defectors cannot arise or adapt via mutation.

355 **Figure 5**




Figure 5: **Defector Invasion Stopped by Cooperator Adaptation.** Here we depict the distribution of dominant types among populations over time for one representative simulation in which isogenic defectors arise. For clarity, mutations occurred at the adaptive loci, but not at the cooperation locus ($\mu_c = 0$) during this ecological simulation. A time $t = 0$ (leftmost panel), a single matched defector population (red) is placed among cooperator populations (light blue). Because these defectors do not bear the costs of cooperation, they spread ($t = 272$, second panel). However, cooperators in a single population gain an adaptation that give them a fitness advantage over defectors (dark blue, lower left). At $t = 325$ (third panel), defectors continue to invade cooperator populations. However, the adapted cooperator type, which can invade both defector populations and ancestral cooperator populations, can spread more quickly due to its greater fitness. Eventually, this strategy spreads and fixes in all populations (rightmost panel) until this strategy itself is replaced by the next adaptation.

356 **Figure 6**

357 **TODO** Yep. Almost ready.

³⁵⁸ **Supplemental Figure 1**




Figure S1: **Defector Invasion with Mutations.** With mutations occurring both at the adaptive loci and the cooperation locus ($\mu_a = \mu_c = 0.00005$), cooperation remains the dominant strategy in 58 replicate simulations. Curves trace the proportion of cooperators present in the population for the duration of 160 replicate simulations ($T = 1000$)

Table 1: Model parameters and their value

| Parameter | Description | Base Value |
|------------|--|------------|
| L | Number of adaptive loci | 5 |
| c | Fitness cost of cooperation | 0.1 |
| A | Number of alleles | 6 |
| δ | Fitness benefit, nonzero alleles | 0.3 |
| ϵ | Fitness benefit, sequential alleles | 0.00015 |
| z | Baseline fitness | 1 |
| S_{min} | Minimum subpopulation size | 800 |
| S_{max} | Maximum subpopulation size | 2000 |
| μ_a | Mutation rate (adaptation) | 10^{-5} |
| μ_c | Mutation rate (cooperation) | 10^{-5} |
| N^2 | Number of patches | 625 |
| m | Migration rate | 0.05 |
| p_0 | Initial cooperator proportion | 0.5 |
| μ_t | Mutation rate (tolerance to new environment) | 10^{-5} |
| T | Number of simulation cycles | 3000 |
| d | Subpopulation dilution factor | 0.1 |

References

- Allison, S.D. 2005. Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. *Ecology Letters*, **8**: 626–635.
- Asfahl, K.L., Walsh, J., Gilbert, K. and Schuster, M. 2015. Non-social adaptation defers a tragedy of the commons in *Pseudomonas aeruginosa* quorum sensing. *The ISME Journal*, doi: [10.1038/ismej.2014.259](https://doi.org/10.1038/ismej.2014.259).
- Bernier, S.P., Ha, D.-G., Khan, W., Merritt, J.H.M. and O'Toole, G.A. 2011. Modulation of *Pseudomonas aeruginosa* surface-associated group behaviors by individual amino acids through c-di-GMP signaling. *Research in Microbiology*, **162**: 680–688.
- Brown, S.P. and Johnstone, R.A. 2001. Cooperation in the dark: Signalling and collective action in quorum-sensing bacteria. *Proceedings of the Royal Society of London B: Biological Sciences*, **268**: 961–965.
- Brown, S.P. and Taddei, F. 2007. The durability of public goods changes the dynamics and nature of social dilemmas. *PLoS ONE*, **2**: e593.
- Cosson, P., Zulianello, L., Join-Lambert, O., Faurisson, F., Gebbie, L. and Benghezal, M.*et al.* 2002. *Pseudomonas aeruginosa* virulence analyzed in a *Dictyostelium discoideum* host system. *Journal of Bacteriology*, **184**: 3027–3033.
- Dandekar, A.A., Chugani, S. and Greenberg, E.P. 2012. Bacterial quorum sensing and metabolic incentives to cooperate. *Science*, **338**: 264–266.

382 Darch, S.E., West, S.A., Winzer, K. and Diggle, S.P. 2012. Density-dependent
383 fitness benefits in quorum-sensing bacterial populations. *Proceedings of the*
384 *National Academy of Sciences*, **109**: 8259–8263.

385 Diggle, S.P., Griffin, A.S., Campbell, G.S. and West, S.A. 2007. Cooperation
386 and conflict in quorum-sensing bacterial populations. *Nature*, **450**: 411–414.

387 Driscoll, W.W. and Pepper, J.W. 2010. Theory for the evolution of diffusible
388 external goods. *Evolution*, **64**: 2682–2687.

389 Dumas, Z. and Kümmerli, R. 2012. Cost of cooperation rules selection for
390 cheats in bacterial metapopulations. *Journal of Evolutionary Biology*, **25**:
391 473–484.

392 Fletcher, J.A. and Doebeli, M. 2009. A simple and general explanation for the
393 evolution of altruism. *Proceedings of the Royal Society B: Biological Sciences*,
394 **276**: 13–19.

395 Foster, K., Shaulsky, G., Strassmann, J., Queller, D. and Thompson, C. 2004.
396 Pleiotropy as a mechanism to stabilize cooperation. *Nature*, **431**: 693–696.

397 Gardner, A. and West, S.A. 2010. Greenbeards. *Evolution*, **64**: 25–38.

398 Ghoul, M., West, S.A., Diggle, S.P. and Griffin, A.S. 2014. An experimental
399 test of whether cheating is context dependent. *Journal of Evolutionary Biology*,
400 **27**: 551–556.

401 Griffin, A.S., West, S.A. and Buckling, A. 2004. Cooperation and competition
402 in pathogenic bacteria. *Nature*, **430**: 1024–1027.

403 Hafer, N. and Milinski, M. 2015. When parasites disagree: Evidence for

404 parasite-induced sabotage of host manipulation. *Evolution*, **69**: 611–620.

405 Hagberg, A.A., Schult, D.A. and Swart, P.J. 2008. Exploring network struc-
 406 ture, dynamics, and function using NetworkX. In: *Proceedings of the 7th*
 407 *Python in Science Conference (SciPy2008)*, pp. 11–15.

408 Hamblin, S.R., White, P.A. and Tanaka, M.M. 2014. Viral niche construction
 409 alters hosts and ecosystems at multiple scales. *Trends in Ecology & Evolution*,
 410 **29**: 594–599.

411 Hamilton, W.D. 1964. The genetical evolution of social behaviour I & II.
 412 *Journal of Theoretical Biology*, **7**: 1–52.

413 Hammarlund, S.P., Connelly, B.D., Dickinson, K.J. and Kerr, B. 2015. The
 414 evolution of cooperation by the Hankshaw effect. *bioRxiv*, doi: [10.1101/016667](https://doi.org/10.1101/016667).
 415 Cold Spring Harbor Labs Journals.

416 Harrison, F. 2007. Microbial ecology of the cystic fibrosis lung. *Microbiology*,
 417 **153**: 917–923.

418 Koestler, B.J. and Waters, C.M. 2014. Bile acids and bicarbonate inversely
 419 regulate intracellular cyclic di-GMP in *Vibrio cholerae*. *Infection and Immu-*
 420 *nity*, **82**: 3002–3014.

421 Kuzdzal-Fick, J.J., Fox, S.A., Strassmann, J.E. and Queller, D.C. 2011. High
 422 relatedness is necessary and sufficient to maintain multicellularity in *Dic-*
 423 *tyostelium*. *Science*, **334**: 1548–1551.

424 Kümmerli, R. and Brown, S.P. 2010. Molecular and regulatory properties of
 425 a public good shape the evolution of cooperation. *Proceedings of the National*

426 *Academy of Sciences*, **107**: 18921–18926.

427 Laland, K.N., Odling-Smee, F.J. and Feldman, M.W. 1999. Evolutionary con-
428 sequences of niche construction and their implications for ecology. *Proceedings*
429 *of the National Academy of Sciences*, **96**: 10242–10247.

430 Laland, K.N., Odling-Smee, F.J. and Feldman, M.W. 1996. The evolutionary
431 consequences of niche construction: A theoretical investigation using two-locus
432 theory. *Journal of Evolutionary Biology*, **9**: 293–316.

433 Lehmann, L. 2007. The evolution of trans-generational altruism: Kin selection
434 meets niche construction. *Journal of Evolutionary Biology*, **20**: 181–189.

435 Maynard Smith, J. and Haigh, J. 1974. The hitch-hiking effect of a favourable
436 gene. *Genetics Research*, **23**: 23–35.

437 McKinney, W. 2010. Data structures for statistical computing in Python. In:
438 *Proceedings of the 9th Python in Science Conference* (S. van der Walt and J.
439 Millman, eds), pp. 51–56.

440 Morgan, A.D., Quigley, B.J.Z., Brown, S.P. and Buckling, A. 2012. Selection
441 on non-social traits limits the invasion of social cheats. *Ecology Letters*, **15**:
442 841–846.

443 Nadell, C.D., Foster, K.R. and Xavier, J.B. 2010. Emergence of spatial struc-
444 ture in cell groups and the evolution of cooperation. *PLoS Computational*
445 *Biology*, **6**: e1000716.

446 Nowak, M.A. 2006. Five rules for the evolution of cooperation. *Science*, **314**:
447 1560–1563.

- 448 Odling-Smee, F.J., Laland, K.N. and Feldman, M.W. 2003. *Niche construc-*
449 *tion: The neglected process in evolution*. Princeton University Press.
- 450 R Core Team. 2015. *R: A language and environment for statistical computing*.
451 Vienna, Austria: R Foundation for Statistical Computing.
- 452 Ruby, E.G. 1996. Lessons from a cooperative, bacterial-animal association:
453 The *Vibrio fischeri*–*Euprymna scolopes* light organ symbiosis. *Annual Review*
454 *of Microbiology*, **50**: 591–624.
- 455 Sinervo, B., Chaine, A., Clobert, J., Calsbeek, R., Hazard, L. and Lancaster,
456 L.*et al.* 2006. Self-recognition, color signals, and cycles of greenbeard mutu-
457 alism and altruism. *Proceedings of the National Academy of Sciences*, **103**:
458 7372–7377.
- 459 Van Dyken, J.D. and Wade, M.J. 2012. Origins of altruism diversity II: Run-
460 away coevolution of altruistic strategies via “reciprocal niche construction”.
461 *Evolution*, **66**: 2498–2513.
- 462 Veelders, M., Brückner, S., Ott, D., Unverzagt, C., Mösch, H.-U. and Essen,
463 L.-O. 2010. Structural basis of flocculin-mediated social behavior in yeast.
464 *Proceedings of the National Academy of Sciences*, **107**: 22511–22516.
- 465 Waite, A.J. and Shou, W. 2012. Adaptation to a new environment allows coop-
466 erators to purge cheaters stochastically. *Proceedings of the National Academy*
467 *of Sciences*, **109**: 19079–19086.
- 468 West, S.A., Griffin, A.S. and Gardner, A. 2007. Evolutionary explanations for
469 cooperation. *Current Biology*, **17**: R661–R672.

470 Zhang, X.-X. and Rainey, P.B. 2013. Exploring the sociobiology of pyoverdin-
471 producing *Pseudomonas*. *Evolution*, **67**: 3161–3174.