Grouping mechanisms for object-based vision and attention

by

Brian H. Hu

A dissertation submitted to The Johns Hopkins University in conformity with the requirements for the degree of Doctor of Philosophy.

Baltimore, Maryland

May, 2017

© Brian H. Hu 2017

All rights reserved

Abstract

The visual brain faces the difficult task of reconstructing a three-dimensional (3D) world from two-dimensional (2D) retinal images. In doing so, visual information is organized in terms of objects in 3D space, and this organization is the basis for selective attention, object recognition, and action planning. In complex visual scenes, both the foreground and the background are rich in features of different types, scales, etc. The brain must find a way to group together the features that belong to objects on the foreground, and distinguish them from features in the background.

The goal of this thesis is to understand how the neural circuits in primate cortex accomplish this task using feedback grouping mechanisms for object-based vision and attention. In Chapter 1, we introduce the background information needed to understand the physiology and previous modeling experiments. In Chapter 2, we propose a quantitative neural model of contour grouping constrained by recent physiological data. We validate the model by reproducing several experimental results, including the measure of contour-response d', as well as the magnitude and time course of neuronal responses to contours. In Chapter 3, we extend this model to natural images,

ABSTRACT

and the results are quantitatively compared with human-generated segmentations and

figure-ground labels (Berkeley Segmentation Dataset). Beginning with Chapter 4, we

shift our focus to the representation of 3D information in the visual system. First,

we show that 3D surfaces can be represented by a feedforward, linear combination of

basis functions whose response properties are similar to those of disparity-selective

neurons commonly found in early visual cortex. With our model, we are able to

reproduce results from a set of psychophysical experiments where attention has to

be directed to surfaces. In Chapter 5, we propose a model of 3D visual saliency

and show that the added depth information improves saliency prediction. Overall,

this work will investigate whether feedback grouping mechanisms are fundamental

for linking early feature representations to perceptual objects. The models developed

will address how visual features are grouped into 2D and 3D object representations.

Primary Reader: Ernst Niebur

Secondary Reader: Rüdiger von der Heydt

iii

Acknowledgments

As I was sitting on the boat, I was thinking about how the trip was really a metaphor for my PhD experience. I have had opportunities to steer the boat (sometimes in the wrong direction!), gone through both good and bad weather, and through it all, have had a good captain whom I can trust to guide me to the finish.

Dedication

This thesis is dedicated to my younger sister Joy, whose perseverance in the face of adversity has been my inspiration.

Contents

A	bstra	act	ii
A	ckno	wledgments	iv
Li	st of	Tables	ix
Li	st of	Figures	x
1	Intı	roduction	1
	1.1	Segmentation and figure-ground organization	1
	1.2	The role of cortical feedback	3
	1.3	The role of attention	4
	1.4	Overview of the grouping model	8
2	Cor	ntour Integration	12
	2.1	Introduction	12
	2.2	Materials and methods	15

CONTENTS

		2.2.1	Model Structure	15
		2.2.2	Model Implementation	19
		2.2.3	Contour integration experiments	21
		2.2.4	Figure-ground segregation experiments	23
		2.2.5	Quantitative assessment of border ownership selectivity:	
			Vector modulation index	25
	2.3	Result	s	26
		2.3.1	Contour enhancement in V1 and V4	26
		2.3.2	The role of feedback and attention in contour grouping \dots .	32
		2.3.3	Border ownership assignment and highlighting figures in noise	35
		2.3.4	Border ownership assignment in the presence of multiple objects	38
	2.4	Discus	sion	39
		2.4.1	Model predictions	39
		2.4.2	Comparison to other models	41
		2.4.3	Roles of V1 and V4 in visual processing	42
		2.4.4	Contour and object grouping neurons	43
		2.4.5	Scope and limitations of the model	44
3	3D	Surfac	e Representation	46
	3.1	Introd	uction	46
	3.2		ds	49
	3.3		s	51
	5.5	1 COUIT	~	O T

CONTENTS

	3.4	Conclusion	54
4	3D	Proto-object based Saliency	56
	4.1	Introduction	56
	4.2	Related Work	60
		4.2.1 Models of 3D visual attention	60
		4.2.2 3D eyetracking datasets	63
	4.3	Model	66
	4.4	Results	70
	4.5	Discussion	75

List of Tables

List of Figures

1.1	Grouping	model																															9
-----	----------	-------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---

2.1	Structure of the model network. Each circle stands for a population	
	of neurons with similar receptive fields and response properties. Ma-	
	genta, blue, and orange lines represent feedforward excitatory, lateral	
	inhibitory, and feedback excitatory projections, respectively. Edges	
	and other local features of a figure (black dashed parallelogram) acti-	
	vate edge cells (E), whose receptive fields are shown by green ellipses.	
	Edge cells project to border ownership cells (B) that have the same	
	preferred orientation and retinotopic position as the E cells they re-	
	ceive input from. However, for each location and preferred orientation	
	there are two B cell populations with opposite side-of-figure prefer-	
	ences, in the example shown B_L whose neurons respond preferentially	
	when the foreground object is to the left of their receptive fields and	
	B_R whose members prefer the foreground to the right side of their re-	
	ceptive fields. E cells also excite other E cells with the same preferred	
	orientation (connections not shown), as well as a class of inhibitory	
	cells (IE) which, in turn, inhibit E cells of all preferred orientations at	
	a given location (only E cells of one preferred orientation are shown). B	
	cells have reciprocal, forward excitatory and feedback modulatory con-	
	nections with two types of grouping cells, G_c and G_o , which integrate	
	global context information about contours and objects, respectively. E	
	cells also receive positive modulatory feedback from these same group-	
	ing cells. Opposing border ownership cells compete directly via IB cells	
	and indirectly via grouping cells, which bias their activity and thus gen-	
	erate the response differences of opposing border ownership selective	
	neurons. G cell populations also directly excite inhibitory grouping	
	cells (IG; again with the indices L and R), which inhibit G_c cells non-	
	specifically and G_o cells in all orientations except the preferred one.	
	Top-down attention is modeled as input to the grouping cells and can	
	therefore either be directed towards objects (solid lines) or contours	
	(dashed lines) in the visual field (top)	16
2.2	Spatial distribution of border ownership cell to grouping cell connec-	10
2.2		
	tivity; darker pixels indicate stronger connection weights. A) Contour	
	grouping neurons integrate features along oriented contours (horizontal	
	line shown in black), emphasizing the Gestalt principle of good contin-	
	uation. B) Object grouping neurons integrate features in a co-circular	
	pattern (square figure shown in black), emphasizing the Gestalt prin-	

хi

ciples of convexity and proximity.

18

2.3	Normalized V1 E cell and V4 G_c cell population responses to contours	
	of varying lengths in either the contour (A) or the background (B)	
	condition. In (A), the "recorded" neuron is on the contour; in (B), it	
	is offset from the contour. The top row shows stimuli, the second and	
	third rows show activity in model areas V1 and V4, respectively. Yel-	
	low circles mark the RFs of the V1 neurons whose activity is shown in	
	Figure 2.4. Activity from model area V2 is not shown because a single	
	contour does not produce clear border ownership selectivity, and the	
	- · · · · · · · · · · · · · · · · · · ·	
	activity in V2 is essentially the same as that in V1, but with reduced	
	spatial resolution due to the lower number of neurons. Columns in	
	each condition show, from left to right, increasing contour length, with	
	the right-most column showing a jittered stimulus configuration (see	
	text). Neural activity is color coded and normalized to the 7-bar stim-	
	ulus in both contour and background conditions, with warmer colors	
	representing higher activity (see color bar at right)	22
2.4	Normalized V1 E cell (contour and background sites) and V4 G_c	
	cell neuronal activity and contour-response d' to contours of varying	
	lengths. (A) V1 contour (top) and background (middle) sites and V4	
	sites (bottom) showed facilitation followed by saturation with increas-	
	ing contour length (see legend). V1 background sites showed greater	
	suppression with longer contours. The jitter condition involved a 7-bar	
	pattern where each bar was laterally offset to disrupt collinearity. (B)	
	Corresponding experimental observations showing normalized and av-	
	eraged PSTHs from the Chen et al. (2014) study. (C) Contour-response	
	d' was higher for the V4 sites compared to the V1 contour sites, and	
	was facilitated by increasing contour length. V1 background sites had	
	increasingly negative d' with longer contours, indicating background	
	suppression. The jitter condition reduced the absolute value of the d'	
	values to close to zero, making it similar to the baseline noise condi-	
	tion. (D) Corresponding experimental observations, showing the mean	
	contour-response d' from the Chen et al. (2014) study. Panels B and	
	D are modified from Figure 2 of Chen et al. (2014). All model results	
	(neural responses and contour-response d') are averages for a single	
	neuron over 100 simulations	27
2.5	Contour-response d' in V1 E cells (A) and V4 G_c cells (B) for the model	
	with (green) and without attention (black), and for the model with	
	feedback removed (magenta). Attention strongly increased contour-	
	response d' in V4 (B), while the lack of feedback strongly decreased	
	contour-response d' in V1 (A)	29

2.6	Figure-ground segregation of a square object with (bottom row) and without (top row) noise. Shown are (left to right) the input stimulus, the edge cell activity (E), the border ownership assignment along edges (shown as the vector modulation index \vec{v} , section 2.2.5), the object grouping neuron activity (G_o) and the contour grouping activity (G_c) . Activities are normalized within each map, and warmer colors indicate	
2.7	higher activity (see color bar at right)	31
2.8	background. Panels A and B are modified from Figure 3 of Qiu et al. (2007)	34 37
3.1 3.2	Network structure (adapted from ref. Marshall et al. (1996)) Psychophysical and model results (adapted from ref He and Nakayama (1995)). For each trial type (A or B), the top row shows the different stimuli, the middle row shows representative reaction times, and the bottom row shows the degree of attentional modulation of disparity-selective cells on the attended plane. Increase in activity is assumed	49
	to be inversely proportion to reaction times	52

3.3	Spread of attention across surfaces. When attention is directed to the center slanted plane (as in the experiment in Figure 2B-a), attention enhances the activity of all cells along the surface (red), while suppressing the activity of cells belonging to other surfaces (blue)	53
4.1	Examples of data used and results obtained. Columns (left to right) show one example each of the original image with its corresponding depth map, fixation map, our saliency model without (S) and with (S_d) depth information for the three 3D eye tracking datasets: a) NUS-3D.	
	b) Gaze-3D. c) NCTU-3D	64
4.2	Proto-object saliency model with added depth information. The depth	
	map is represented by the image at the top, far right, and the 2D image	
	is to its left. Based on Figure 5 of Russell et al. (2014)	67

Bibliography

- D. Ardila, S. Mihalas, R. von der Heydt, and E. Niebur. Medial axis generation in a model of perceptual organization. In *IEEE CISS-2012 46th Annual Conference* on Information Sciences and Systems, pages 1–4, Princeton University, NJ, 2012. IEEE.
- Y Benjamini and Y Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society*.

 Series B, 1995. URL http://www.jstor.org/stable/2346101.
- Ali Borji and Laurent Itti. State-of-the-art in visual attention modeling. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 2013.
- Ali Borji, Dicky N Sihite, and Laurent Itti. Objects do not predict fixations better than early saliency: A re-analysis of Einhäuser et al.'s data. *Journal of vision*, 13 (10):18, 2013.
- William H Bosking, Ying Zhang, Brett Schofield, and David Fitzpatrick. Orienta-

- tion selectivity and the arrangement of horizontal connections in tree shrew striate cortex. The Journal of Neuroscience, 17(6):2112–2127, 1997.
- S.L. Brincat and C.E. Connor. Underlying principles of visual shape selectivity in posterior inferotemporal cortex. *Nature Neuroscience*, 7:880–886, 2004.
- Neil DB Bruce and John K Tsotsos. An attentional framework for stereo vision. In Computer and Robot Vision, 2005. Proceedings. The 2nd Canadian Conference on, pages 88–95. IEEE, 2005.
- M. Cerf, J. Harel, W. Einhäuser, and C. Koch. Predicting human gaze using low-level saliency combined with face detection. *Advances in Neural Information Processing Systems*, 20:241–248, 2008.
- Garga Chatterjee, Daw-An Wu, and Bhavin R Sheth. Phantom flashes caused by interactions across visual space. *Journal of vision*, 11(2):14, 2011.
- Minggui Chen, Yin Yan, Xiajing Gong, Charles D Gilbert, Hualou Liang, and Wu Li. Incremental integration of global contours through interplay between visual cortical areas. *Neuron*, 82(3):682–694, 2014.
- Michele A Cox, Michael C Schmid, Andrew J Peters, Richard C Saunders, David A Leopold, and Alexander Maier. Receptive field focus of visual area V4 neurons determines responses to illusory surfaces. *Proceedings of the National Academy of Sciences*, 110(42):17095–17100, 2013.

- E. Craft, H. Schütze, E. Niebur, and R. von der Heydt. A neural model of figure-ground organization. *Journal of Neurophysiology*, 97(6):4310–26, 2007. PMID: 17442769.
- Y. Dong, S. Mihalas, F. Qiu, R. von der Heydt, and E. Niebur. Synchrony and the binding problem in macaque visual cortex. *Journal of Vision*, 8(7):1–16, 2008. URL http://journalofvision.org/8/7/30/,doi:10.1167/8.7.30. PMC2647779.
- J. Driver and G.C. Baylis. Edge-assignment and figure-ground segmentation in short-term visual matching. *Cognit Psychol*, 31(3):248–306, 1996.
- J. Duncan. Selective attention and the organization of visual information. J Exp Psychol Gen, 113:501–517, Dec 1984.
- R. Egly, J. Driver, and R. Rafal. Shifting visual attention between objects and locations: evidence for normal and parietal lesion subjects. *Journal of Experimental Psychology: General*, 123:161–77, 1994.
- W. Einhäuser, M. Spain, and P. Perona. Objects predict fixations better than early saliency. *J. Vision*, 8(14):1–26, 2008.
- Boris Epshtein, Ita Lifshitz, and Shimon Ullman. Image interpretation by a single bottom-up top-down cycle. *Proceedings of the National Academy of Sciences*, 105 (38):14298–14303, 2008.
- D. J. Field, A. Hayes, and R. F. Hess. Contour integration by the human visual

system: evidence for a local association field. Vision Research, 33(2):173–193, 1993.

- Josselin Gautier and Olivier Le Meur. A time-dependent saliency model combining center and depth biases for 2D and 3D viewing conditions. *Cognitive Computation*, 4(2):141–156, 2012.
- Ariel Gilad, Elhanan Meirovithz, and Hamutal Slovin. Population responses to contour integration: early encoding of discrete elements and late perceptual grouping.

 Neuron, 78(2):389–402, 2013.
- C.D. Gilbert and T.N. Wiesel. Columnar specificity of intrinsic horizontal and corticocortical connections in cat visual cortex. J. Neurosci., 9:2432–2442, 1989.
- P. Girard, J.M. Hupé, and J. Bullier. Feedforward and feedback connections between areas V1 and V2 of the monkey have similar rapid conduction velocities. J. Neurophysiol., 85:1328–1331, 2001.
- D. M. Green and J. A. Swets. Signal Detection Theory and Psychophysics. Wiley, New York, N.Y., 1966.
- S. Grossberg. 3-D vision and figure-ground separation by visual cortex. *Perception & Psychophysics*, 55:48–120, 1994.
- S. Grossberg. Cortical dynamics of three-dimensional figure—ground perception of two-dimensional pictures. *Psychological review*, 104(3):618, 1997. PMID: 9243966.

- Z. J. He and K. Nakayama. Surfaces versus features in visual search. Nature, 359: 231–233, 1992. PMID: 1528263.
- Z. J. He and K. Nakayama. Visual attention to surfaces in three-dimensional space.
 Proc. Natl. Acad. Sci. U. S. A., 9(24):11155–11159, 1995. PMID: 7479956.
- Jay Hegdé and David C Van Essen. A comparative study of shape representation in macaque visual areas V2 and V4. *Cerebral Cortex*, 17(5):1100–1116, 2007.
- F. Heitger and R. von der Heydt. A computational model of neural contour processing: figure-ground segregation and illusory contours. In *Proc. 4th Int. Conf. Computer Vision*, Proc. 4th Int. Conf. Computer Vision, pages 32–40. IEEE Computer Society Press, 1993.
- Ming-Chou Ho and Su-Ling Yeh. Effects of instantaneous object input and past experience on object-based attention. *Acta psychologica*, 132(1):31–39, 2009.
- Shaul Hochstein and Merav Ahissar. View from the top: Hierarchies and reverse hierarchies in the visual system. *Neuron*, 36(5):791–804, 2002.
- B. Hu, R. von der Heydt, and E. Niebur. A neural model for perceptual organization of 3D surfaces. In *IEEE CISS-2015 49th Annual Conference on Information Sciences* and Systems, pages 1–6, Baltimore, MD, 2015. IEEE Information Theory Society. doi: 10.1109/CISS.2015.7086906.

- D.H. Hubel and T.N. Wiesel. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *J. Physiol.*, 160:106–154, 1962.
- Heinz Hügli, Timothée Jost, and Nabil Ouerhani. Model performance for visual attention in real 3D color scenes. In Artificial Intelligence and Knowledge Engineering Applications: A Bioinspired Approach, pages 469–478. Springer, 2005.
- Quan Huynh-Thu and Luca Schiatti. Examination of 3D visual attention in stereoscopic video content. In *IS&T/SPIE Electronic Imaging*, pages 78650J–78650J. International Society for Optics and Photonics, 2011.
- J. Intriligator and P. Cavanagh. The spatial resolution of visual attention. Cognit. Psychol., 43:171–216, 2001.
- L. Itti, C. Koch, and E. Niebur. A model of saliency-based fast visual attention for rapid scene analysis. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 20(11):1254–1259, November 1998.
- Lina Jansen, Selim Onat, and Peter König. Influence of disparity on fixation and saccades in free viewing of natural scenes. *Journal of Vision*, 9(1):29, 2009.
- Ana Karla Jansen-Amorim, Mario Fiorani, and Ricardo Gattass. GABA inactivation of area V4 changes receptive-field properties of V2 neurons in Cebus monkeys. Experimental Neurology, 235(2):553–562, 2012.

Timothée Jost, Nabil Ouerhani, Roman von Wartburg, René Müri, and Heinz Hügli.

- Contribution of depth to visual attention: comparison of a computer model and human. In *Proceedings. Early cognitive vision workshop*, pages 1–4, 2004.
- Tilke Judd, Frédo Durand, and Antonio Torralba. A Benchmark of Computational Models of Saliency to Predict Human Fixations. In *MIT Technical Report*, 2012.
- R. Kimchi, Y. Yeshurun, and A. Cohen-Savransky. Automatic, stimulus-driven attentional capture by objecthood. *Psychon Bull Rev*, 14(1):166–172, Feb 2007.
- C. Koch and S. Ullman. Shifts in selective visual attention: towards the underlying neural circuitry. *Human Neurobiol.*, 4:219–227, 1985.
- K. Koffka. Principles of Gestalt psychology. Harcourt-Brace, New York, 1935.
- V. A. F. Lamme. The neurophysiology of figure-ground segregation in primary visual cortex. J Neurosci, 15:1605–1615, 1995.
- V. A. F. Lamme, K. Zipser, and H. Spekreijse. Figure-ground activity in primary visual cortex is suppressed by anesthesia. *Proc. Natl. Acad. Sci. U. S. A.*, 9(6): 3263–3268, 1998.
- Congyan Lang, Tam V Nguyen, Harish Katti, Karthik Yadati, Mohan Kankanhalli, and Shuicheng Yan. Depth matters: Influence of depth cues on visual saliency. In Computer Vision–ECCV 2012, pages 101–115. Springer, 2012.
- Oliver W Layton, Ennio Mingolla, and Arash Yazdanbakhsh. Dynamic coding of border-ownership in visual cortex. *Journal of vision*, 12(13):8, 2012.

- P. Le Callet and E. Niebur. Visual Attention and Applications in Multimedia Technologies. *IEEE Proceedings*, 101(9):2058–67, 2013. NIHMS539064.
- Tai Sing Lee, David Mumford, Richard Romero, and Victor A. F. Lamme. The Role of the Primary Visual Cortex in Higher Level Vision. Vision Research, 38:2429–2452, 1998.
- Wu Li, Valentin Piëch, and Charles D Gilbert. Perceptual learning and top-down influences in primary visual cortex. *Nature neuroscience*, 7(6):651–657, 2004.
- Wu Li, Valentin Piëch, and Charles D Gilbert. Contour saliency in primary visual cortex. *Neuron*, 50(6):951–962, 2006.
- Wu Li, Valentin Piëch, and Charles D Gilbert. Learning to link visual contours.

 Neuron, 57(3):442–451, 2008.
- Z. Li. A Neural Model of Contour Integration in the Primary Visual Cortex. Neural Computation, 10(903-940), 1998.
- Chih-Yao Ma and Hsueh-Ming Hang. Learning-based saliency model with depth information. *Journal of vision*, 15(6):19–19, 2015.
- D. Marr and T. Poggio. Cooperative Computation of Stereo Disparity. Science, 194, 1976. PMID: 968482.
- John C Marshall and Peter W Halligan. Visuo-spatial neglect: A new copying test to assess perceptual parsing. *Journal of neurology*, 240(1):37–40, 1993.

- Jonathan A Marshall, George J Kalarickal, and Elizabeth B Graves. Neural model of visual stereomatching: slant, transparency and clouds. *Network: Computation in Neural Systems*, 7(4):635–669, 1996.
- Anne B Martin and Rüdiger von der Heydt. Spike Synchrony Reveals Emergence of Proto-Objects in Visual Cortex. *The Journal of Neuroscience*, 35(17):6860–6870, 2015.
- C. J. McAdams and J. H. R. Maunsell. Effects of attention on orientation-tuning functions of single neurons in macaque cortical area V4. J. Neurosci., 19:431–441, 1999.
- S. Mihalas, Y. Dong, R. von der Heydt, and E. Niebur. Mechanisms of perceptual organization provide auto-zoom and auto-localization for attention to objects. *Proceedings of the National Academy of Sciences*, 108(18):7583–8, 2011. PMC3088583.
- Ajay K Mishra, Ashish Shrivastava, and Yiannis Aloimonos. Segmenting "simple"??? objects using RGB-D. In *Robotics and Automation (ICRA)*, 2012 IEEE International Conference on, pages 4406–4413. IEEE, 2012.
- B. C. Motter. Neural correlates of attentive selection for color or luminance in extrastriate area V4. *J. Neurosci.*, 14:2178–2189, 1994.
- B.C. Motter. Focal attention produces spatially selective processing in visual cortical

- areas V1, V2, and V4 in the presence of competing stimuli. *J. Neurophysiology*, 70 (3):909–919, 1993.
- K. Nakayama and G. H. Silverman. Serial and parallel processing of visual feature conjunctions. *Nature*, 320:264–265, 1986. PMID: 3960106.
- K. Nakayama, Z. J. He, and S. Shimojo. Visual surface representation: a critical link between lower-level and higher-level vision. In S. Kosslyn and D. Osherson, editors, Visual Cognition: An Invitation to Cognitive Science, volume 2, chapter 1, pages 1–70. The MIT Press, 2nd edition, 1995.
- E. Niebur. Separate but equal: Different kinds of information require different neural representations. In H. Bothe, editor, *Proceedings of the International Congress on Intelligent Systems and Applications (ISA-BIS)*, pages 1544–9, Wetaskiwin, Canada, December 2000. ICSC Academic Press. ISBN 3-906454-24-X.
- E. Niebur and C. Koch. Control of Selective Visual Attention: Modeling the "Where" Pathway. In D. S Touretzky, M. C. Mozer, and M. E. Hasselmo, editors, Advances in Neural Information Processing Systems, volume 8, pages 802–808. MIT Press, Cambridge, MA, 1996.
- P.J. O'Herron and R. von der Heydt. Short-term memory for figure-ground organization in the visual cortex. *Neuron*, 61(5):801–809, 2009. PMC2707495.
- P.J. O'Herron and R. von der Heydt. Remapping of Border Ownership in the Visual

- Cortex. The Journal of Neuroscience, 33(5):1964–1974, 2013. PMID: 23365235 [PubMed in process].
- I. Ohzawa, G. C. DeAngelis, and R. D. Freeman. Stereoscopic depth discrimination in the visual cortex: neurons ideally suited as disparity detectors. *Science*, 249: 1037–1041, 1990.
- Nabil Ouerhani and Heinz Hügli. Computing visual attention from scene depth.

 In Pattern Recognition, 2000. Proceedings. 15th International Conference on, volume 1, pages 375–378. IEEE, 2000.
- Lucy M Palmer, Adam S Shai, James E Reeve, Harry L Anderson, Ole Paulsen, and Matthew E Larkum. NMDA spikes enhance action potential generation during sensory input. *Nature neuroscience*, 17(3):383–390, 2014.
- S.E. Palmer and T. Ghose. Extremal Edge—A Powerful Cue to Depth Perception and Figure-Ground Organization. *Psychological Science*, 19(1):77, 2008. ISSN 0956-7976.
- D. Parkhurst, K. Law, and E. Niebur. Modelling the role of salience in the allocation of visual selective attention. *Vision Research*, 42(1):107–123, 2002.
- Anitha Pasupathy and Charles E Connor. Population coding of shape in area V4.

 Nature neuroscience, 5(12):1332–1338, 2002.
- Valentin Piëch, Wu Li, George N Reeke, and Charles D Gilbert. Network model

of top-down influences on local gain and contextual interactions in visual cortex. *Proceedings of the National Academy of Sciences*, 110(43):E4108–E4117, 2013. PMC3808648.

- G. F. Poggio and B. Fischer. Binocular interaction and depth sensitivity in striate and prestriate cortex of behaving rhesus monkey. *J. Neurophysiol.*, 40:1392–1405, Nov 1977. PMID: 411898.
- G.F. Poggio and T. Poggio. The analysis of stereopsis. Ann. Rev. Neurosci., 7: 379–412, 1984.
- Gian F Poggio, Francisco Gonzalez, and F Krause. Stereoscopic mechanisms in monkey visual cortex: binocular correlation and disparity selectivity. *The Journal of* neuroscience, 8(12):4531–4550, 1988.
- Uri Polat, Keiko Mizobe, Mark W. Pettet, Takuji Kasamatsu, and Anthony M. Norcia. Collinear Stimuli Regulate Visual Responses Depending on Cell's Contrast Threshold. *Nature*, 391:580–584, February 1998.
- Jasper Poort, Florian Raudies, Aurel Wannig, Victor AF Lamme, Heiko Neumann, and Pieter R Roelfsema. The role of attention in figure-ground segregation in areas V1 and V4 of the visual cortex. *Neuron*, 75(1):143–156, 2012.
- F. T. Qiu and R. von der Heydt. Figure and ground in the visual cortex: V2 combines stereoscopic cues with Gestalt rules. Neuron, 47:155–166, 2005.

- F. T. Qiu and R. von der Heydt. Neural representation of transparent overlay. Nat. Neurosci., 10(3):283–284, 2007.
- F. T. Qiu, T. Sugihara, and R. von der Heydt. Figure-ground mechanisms provide structure for selective attention. *Nat. Neurosci.*, 10(11):1492–9, October 2007.
- S. Ramenahalli and E. Niebur. Computing 3D saliency from a 2D image. In *Information Sciences and Systems (CISS)*, 2013 47th Annual Conference on, pages 1-5, 2013. doi: 10.1109/CISS.2013.6552297. URL http://ieeexplore.ieee.org/stamp.jsp?arnumber=6552297.
- S. Ramenahalli, S. Mihalas, and E. Niebur. Extremal Edges: Evidence in Natural Images. In IEEE CISS-2011 45th Annual Conference on Information Sciences and Systems, pages 1–6, Baltimore, MD, 2011. IEEE Information Theory Society.
- S. Ramenahalli, S. Mihalas, and E. Niebur. Figure-ground classification based on spectral properties of boundary image patches. In *IEEE CISS-2012 46th Annual Conference on Information Sciences and Systems*, pages 1–6, Princeton, NJ, 2012. IEEE Information Theory Society.
- Sudarshan Ramenahalli, Stefan Mihalas, and Ernst Niebur. Local spectral anisotropy is a valid cue for figure-ground organization in natural scenes. *Vision Research*, 103: 116–126, Oct 2014. doi: 10.1016/j.visres.2014.08.012. URL http://dx.doi.org/10.1016/j.visres.2014.08.012. NIHMSID 631573.

- R. A. Rensink. The dynamic representation of scenes. Visual Cognition, 7(1/2/3): 17–42, 2000.
- Nicolas Riche, Matthieu Duvinage, Matei Mancas, Bernard Gosselin, and Thierry Dutoit. Saliency and human fixations: state-of-the-art and study of comparison metrics. In *Computer Vision (ICCV)*, 2013 IEEE International Conference on, pages 1153–1160. IEEE, 2013.
- P. R. Roelfsema, V. A. F. Lamme, and H. Spekreijse. Object-based attention in the primary visual cortex of the macaque monkey. *Nature*, 395(6700):376–381, 1998.
- Pieter R Roelfsema. Cortical algorithms for perceptual grouping. Annu. Rev. Neurosci., 29:203–227, 2006.
- Pieter R Roelfsema, Victor A F Lamme, and Henk Spekreijse. Synchrony and covariation of firing rates in the primary visual cortex during contour grouping. *Nat Neurosci*, 7(9):982–991, Sep 2004. doi: 10.1038/nn1304. URL http://dx.doi.org/10.1038/nn1304.
- Ari Rosenberg, Noah J Cowan, and Dora E Angelaki. The Visual Representation of 3D Object Orientation in Parietal Cortex. *The Journal of Neuroscience*, 33(49): 19352–19361, 2013.
- A. F. Russell, S Mihalas, R. von der Heydt, E. Niebur, and R. Etienne-Cummings.
 A model of proto-object based saliency. Vision Research, 94:1–15, 2014.

- P Sajda and L.H. Finkel. Intermediate-Level Visual Representations and the Construction of Surface Perception. *J Cogn Neurosci*, 7:267–291, 1995.
- K. Sakai and H. Nishimura. Surrounding suppression and facilitation in the determination of border ownership. *Journal of Cognitive Neuroscience*, 18(4):562–579, 2006.
- Ashutosh Saxena, Min Sun, and Andrew Y Ng. Make3d: Learning 3d scene structure from a single still image. *Pattern Analysis and Machine Intelligence*, *IEEE Transactions on*, 31(5):824–840, 2009.
- Daniel Scharstein and Chris Pal. Learning conditional random fields for stereo. In Computer Vision and Pattern Recognition, 2007. CVPR'07. IEEE Conference on, pages 1–8. IEEE, 2007.
- Mircea A Schoenfeld, Jens-Max Hopf, Christian Merkel, Hans-Jochen Heinze, and Steven A Hillyard. Object-based attention involves the sequential activation of feature-specific cortical modules. *Nature neuroscience*, 17(4):619–624, 2014.
- B. J. Scholl. Objects and attention: the state of the art. Cognition, 80(1-2):1-46, 2001.
- M. Sigman, G. A. Cecchi, C. D. Gilbert, and M. O. Magnasco. On a common circle: natural scenes and Gestalt rules. *Proc Natl Acad Sci U S A*, 98(4):1935–1940, Feb

2001. doi: 10.1073/pnas.031571498. URL http://dx.doi.org/10.1073/pnas.031571498.

- Enrico Simonotto, Massimo Riani, Charles Seife, Mark Roberts, Jennifer Twitty, and Frank Moss. Visual perception of stochastic resonance. *Physical Review Letters*, 78(6):1186, 1997.
- W. Singer. Neuronal synchrony: a versatile code for the definition of relations?
 Neuron, 24:49–65, 1999.
- M. Stemmler, M. Usher, and E. Niebur. Lateral cortical connections may contribute to both contour completion and redundancy reduction in visual processing. Soc. Neurosci. Abstr., 21(1):510, 1995a.
- M. Stemmler, M. Usher, and E. Niebur. Lateral Interactions in Primary Visual Cortex: A Model Bridging Physiology and Psychophysics. Science, 269:1877–1880, 1995b.
- Dan D Stettler, Aniruddha Das, Jean Bennett, and Charles D Gilbert. Lateral connectivity and contextual interactions in macaque primary visual cortex. *Neuron*, 36(4):739–750, 2002.
- Tadashi Sugihara, Fangtu T Qiu, and Rüdiger von der Heydt. The speed of context integration in the visual cortex. *Journal of neurophysiology*, 106(1):374–385, 2011. PMC3129740.

- Hans Super and Victor AF Lamme. Altered figure-ground perception in monkeys with an extra-striate lesion. *Neuropsychologia*, 45(14):3329–3334, 2007.
- A. Thiele and G. Stoner. Neuronal synchrony does not correlate with motion coherence in cortical area MT. *Nature*, 421(6921):366–370, 2003.
- A. Treisman. The binding problem. Curr Opin Neurobiol, 6(2):171–178, April 1996.
- A. Treisman and G. Gelade. A feature-integration theory of attention. *Cognitive Psychology*, 12:97–136, 1980. PMID: 7351125.
- J. C. M. Treue, S.and Trujillo. Feature-based attention influences motion processing gain in macaque visual cortex. *Nature*, 399:575–9, 1999.
- S. Treue. Neural correlates of attention in primate visual cortex. *Trends in Neuro-sciences*, 24:295–300, May 2001.
- J. K. Tsotsos. Analyzing vision at the complexity level. Behav. Brain Sci., 13: 423–469, 1990.
- S. Ullman. Visual routines. Cognition, 18:97–159, 1984.
- Shimon Ullman, RL Gregory, and J Atkinson. Low-Level Aspects of Segmentation and Recognition [and Discussion]. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 337(1281):371–379, 1992.

- Matthieu Urvoy, Marcus Barkowsky, Romain Cousseau, Yao Koudota, Vincent Ricorde, Patrick Le Callet, Jesus Gutierrez, and Narciso Garcia. NAMA3DS1-COSPAD1: Subjective video quality assessment database on coding conditions introducing freely available high quality 3D stereoscopic sequences. In *Quality of Multimedia Experience (QoMEX)*, 2012 Fourth International Workshop on, pages 109–114. IEEE, 2012.
- Shaun P Vecera, Anastasia V Flevaris, and Joseph C Filapek. Exogenous spatial attention influences figure-ground assignment. *Psychological Science*, 15(1):20–26, 2004.
- R. von der Heydt and R. Pierson. Dissociation of color and figure-ground effects in the watercolor illusion. *Spatial Vision*, 19(2-4):323–340, 2006.
- R. von der Heydt, H. Zhou, and H. S. Friedman. Representation of stereoscopic edges in monkey visual cortex. *Vision Res.*, 40(15):1955–1967, 2000. PMID: 10828464.
- R. von der Heydt, F. T. Qiu, and Z. J. He. Neural mechanisms in border ownership assignment: motion parallax and gestalt cues. *J. Vision*, 3(9):666a, 2003a.
- R. von der Heydt, H. Zhou, and H. S. Friedman. Neural coding of border ownership: Implications for the theory of figure-ground perception. In M. Behrmann, R. Kimchi, and C. R. Olson, editors, *Perceptual Organization in Vision: Behavioral and Neural Perspectives*, Carnegie Symposium Series, pages 281–304. Lawrence Erlbaum Associates, 2003b.

- R. von der Heydt, T. J. Macuda, and F. T. Qiu. Border-ownership dependent tilt aftereffect. Journal of the Optical Society of America A-Optics Image Science and Vision, 22(10):2222–2229, 2005.
- D. Walther and C. Koch. Modeling attention to salient proto-objects. Neural Networks, 19:1395–1407, Nov 2006.
- Junle Wang, Matthieu Perreira DaSilva, Patrick LeCallet, and Vincent Ricordel.
 Computational model of stereoscopic 3D visual saliency. *Image Processing, IEEE Transactions on*, 22(6):2151–2165, 2013.
- Manuel Werlberger, Werner Trobin, Thomas Pock, Andreas Wedel, Daniel Cremers, and Horst Bischof. Anisotropic Huber-L1 Optical Flow. In *Proceedings of the British Machine Vision Conference (BMVC)*, London, UK, September 2009.
- M. Wertheimer. Untersuchungen zur Lehre von der Gestalt II. *Psychol. Forsch.*, 4: 301–350, 1923.
- Jonathan R Williford and Rudiger von der Heydt. Border-ownership coding. Scholarpedia, 8(10):30040, 2013.
- Jonathan R Williford and Rudiger von der Heydt. Early Visual Cortex Assigns Border
 Ownership in Natural Scenes According to Image Context. *Journal of Vision*, 14
 (10):588–588, 2014.
- Jun Xie, Guanghua Xu, Jing Wang, Sicong Zhang, Feng Zhang, Yeping Li,

Chengcheng Han, and Lili Li. Addition of visual noise boosts evoked potential-based brain-computer interface. *Scientific reports*, 4, 2014.

Yin Yan, Malte J Rasch, Minggui Chen, Xiaoping Xiang, Min Huang, Si Wu, and Wu Li. Perceptual training continuously refines neuronal population codes in primary visual cortex. *Nature neuroscience*, 17(10):1380–1387, 2014.

A.L. Yarbus. Eye Movements and Vision. Plenum Press, New York, 1967.

Shih-Cheng Yen and Leif H Finkel. Extraction of perceptually salient contours by striate cortical networks. *Vision research*, 38(5):719–741, 1998.

- N.R. Zhang and R. von der Heydt. Analysis of the context integration mechanisms underlying figure—ground organization in the visual cortex. *The Journal of Neuro-science*, 30(19):6482–6496, 2010. PMC2910339.
- Li Zhaoping. Border ownership from intracortical interactions in visual area V2.

 Neuron, 47:143–153, 2005. PMID: 15996554.
- H. Zhou, H. S. Friedman, and R. von der Heydt. Coding of border ownership in monkey visual cortex. J. Neurosci., 20(17):6594–6611, 2000. PMID: 10964965.