

11/8 in Class – Euler’s Method and one Simulation

edited after class to fix time in Problem 2

1. Use Euler’s Method to find a numerical solution for $\dot{x} = 3t^2$ given that $x(0) = 0$. Let it run from $t = 0$ to $t = 2$ s. Start with at least 10 time steps, and try smaller intervals to see how your answer converges.
 - (a) Do the first few by hand on the white board.
 - (b) Write the equation you will use, $x(k+1) = \dots$
 - (c) Then go back and write a MatLab program to do it for you.
 - (d) Graph both your calculated x and the exact x as functions of time. Compare them.
 - (e) Play around. Decrease the time step and/or increase the total time. What happens?
2. Use Euler’s approximation to solve this DE: $\dot{N} = -\Gamma N$ where $\Gamma = 4.552 \times 10^5 \frac{1}{s}$. At $t = 0$, $N = 1000$. For the first pass, go from $t=0$ to 1×10^{-5} s (fixed this number) seconds with only 10 time steps. (That’s just for debugging. Up the number of time steps when you’re confident it’s working.)
 - (a) This one is separable and so also pretty easy to solve—find the expression for the exact $N(t)$.
 - (b) Write the equations you will use, $N(k+1) = \dots$ and $t(k+1) = \dots$
 - (c) Do the first few by hand on the white board.
 - (d) Then write a MatLab program to do it for you.
 - (e) Graph both your calculated, approximate N and the exact N as functions of time. Compare them.
 - (f) Play around. Decrease the time step by a lot. What happens?
3. Back to simulations (for the DE you just solved!) A muon is a particle that is often created during collisions between cosmic rays and the earth’s atmosphere. A muon is a bit like a heavy electron (about 200 times heavier, with the same charge). Unlike an electron, a muon is unstable, meaning it decays. The muon has a mean lifetime, τ , of 2.197×10^{-6} s. The decay rate per unit time, Γ , is equal to the inverse of the mean lifetime.

$$\Gamma = \frac{1}{\tau}$$

The probability (P) that any given muon will decay in some time interval, dt , is $P = \Gamma dt$. Imagine starting with 1000 muons. Your task is to run a simulation using the random number generator to determine if a given muon decays, and plot the number of muons as a function of time as detailed below.

- (a) Go to the board and work together to write a detailed structure plan for how you will code a simulation of the decay of 1000 muons. Be sure your structure plan answers:
 - i. What do you think is a reasonable size for the time interval, dt ?

- ii. What do you think is a reasonable time scale for the entire simulation to run?
 - iii. What constants do you need?
 - iv. Will you use for loops or vectorize? Explain how you will implement either in words.
- (b) Code it. This could be iterative. Go back to your group and your structure plan if you need a new one! Call it `muonDecay.m`
- (c) Run the simulation of this case and plot the number of particles as a function of time. Use blue circles for the simulated data.
- (d) Write out your data for t and N as two columns in a data file called `muonDecay.txt`
- (e) From the last problem, you calculated the theoretical curve, plot this on top of your simulated data. This time make it a black line. (Which should be close to your blue circles.)