

# 機率與統計 HW3

許博翔

May 13, 2024

**Problem 1.**  $\Pr(K = k \wedge X = x)$

= the probability that "the first  $x$  circuits are acceptable, the  $x+1$ -th is reject, and the  $x+2$ -th to  $n$ -th circuits contain exactly  $k-1$  rejected circuits

$$= p^x \times (1-p) \times \binom{n-x-1}{k-1} p^{n-x-1-k+1} (1-p)^{k-1}$$

$$= \binom{n-x-1}{k-1} p^{n-k} (1-p)^k.$$

$$\therefore P_{K,X}(k, x) = \begin{cases} \binom{n-x-1}{k-1} p^{n-k} (1-p)^k, & \text{if } k+x \leq n, k \geq 1, x \geq 0 \\ 0, & \text{otherwise} \end{cases}.$$

**Problem 2.**  $P_N(n) = 0$  for  $n < 0$ .

$$\text{For } n \geq 0, P_N(n) = \sum_{k=0}^n \frac{100^k e^{-100}}{(n+1)!} = (n+1) \times \frac{100^n e^{-100}}{(n+1)!} = \frac{100^n e^{-100}}{n!}.$$

$$\therefore P_N(n) = \begin{cases} \frac{100^n e^{-100}}{n!}, & \text{if } n \in \mathbb{Z}^+ \cup \{0\} \\ 0, & \text{otherwise} \end{cases}.$$

$\therefore$  By the definition, if  $k > n$ , then  $P_{N,K}(n, k) = 0$ .

$$\begin{aligned} \therefore P_K(k) &= \sum_{n=0}^{\infty} P_{N,K}(n, k) = \sum_{n=k}^{\infty} P_{N,K}(n, k) = \sum_{n=k}^{\infty} \frac{100^n e^{-100}}{(n+1)!} = \sum_{n=k+1}^{\infty} \frac{100^{n-1} e^{-100}}{n!} = \\ &= \frac{1}{100} \sum_{n=k+1}^{\infty} \frac{100^n e^{-100}}{n!} = \frac{1}{100} \sum_{n=k+1}^{\infty} P_N(n) = \frac{1}{100} \Pr(n > k). \end{aligned}$$

**Problem 3.**

**Problem 4.**

(a) If  $x \geq 2$ , then  $F_X(x) = 1$ .

If  $x \leq 0$ , then  $F_X(x) = 0$ .

If  $0 \leq x \leq 2$ , then  $F_X(x) = \int_{-\infty}^x f_X(y)dy = \int_0^x \frac{y}{2}dy = \frac{x^2}{4}$ .

$$\therefore F_X(x) = \begin{cases} 0, & \text{if } x \leq 0 \\ \frac{x^2}{4}, & \text{if } 0 \leq x \leq 2 \\ 1, & \text{otherwise} \end{cases}.$$

(b) Since  $X_1, X_2$  are independent,  $\Pr[X_1 \leq 1, X_2 \leq 1] = \Pr[X_1 \leq 1]\Pr[X_2 \leq 1] = F_X(1)F_X(1) = \left(\frac{1}{4}\right)^2 = \frac{1}{16}$ .

(c)  $F_W(1) = \Pr[\max(X_1, X_2) \leq 1] = \Pr[X_1 \leq 1, X_2 \leq 1] = \frac{1}{16}$ .

(d)  $F_W(w) = \Pr[\max(X_1, X_2) \leq w] = \Pr[X_1 \leq w, X_2 \leq w]$

$$\because X_1, X_2 \text{ are independent } \Pr[X_1 \leq w]\Pr[X_2 \leq w] = F_X(w)^2 = \begin{cases} 0, & \text{if } w \leq 0 \\ \frac{w^4}{16}, & \text{if } 0 \leq w \leq 2 \\ 1, & \text{otherwise} \end{cases}.$$

**Problem 5.** First,  $X \sim \text{Unif}[0, \frac{d}{2}]$ ,  $\Theta \sim \text{Unif}[0, \frac{\pi}{2}]$ .

$$\Rightarrow f_X(x) = \begin{cases} \frac{2}{d}, & \text{if } 0 \leq x \leq \frac{d}{2} \\ 0, & \text{otherwise} \end{cases}, f_\Theta(\theta) = \begin{cases} \frac{2}{\pi}, & \text{if } 0 \leq \theta \leq \frac{\pi}{2} \\ 0, & \text{otherwise} \end{cases}.$$

The needle intersects one of the lines  $\iff X \leq \frac{l}{2} \sin \theta$ .

$\therefore$  the probability that the needle will intersect one of the lines  $= \Pr[X \leq \frac{l}{2} \sin \theta]$ .

Note that  $l < d$ , so the upperbound of  $X$  in the following integral is  $\min(\frac{l}{2} \sin \theta, \frac{d}{2}) = \frac{l}{2} \sin \theta$ .

And  $X, \Theta$  are independent, so their joint pdf  $f_{X,\Theta}(x, \theta) = f_X(x)f_\Theta(\theta)$ .

$$\Rightarrow \Pr[X \leq \frac{l}{2} \sin \theta] = \int_0^{\frac{\pi}{2}} \int_0^{\frac{l}{2} \sin \theta} f_X(x)f_\Theta(\theta)dx d\theta = \int_0^{\frac{\pi}{2}} \frac{l}{2} \sin \theta \frac{2}{d} \frac{2}{\pi} d\theta = \frac{2l}{d\pi}.$$

This experiment can get  $p$ : the approximated value of the probability that the needle will intersect one of the lines when the needle is dropped for sufficient large number of times.

And one can approximate  $\pi \approx \frac{2l}{dp}$ .

**Problem 6.**  $r_{X,Y} = E[XY] = \int_{-1}^1 \int_{-1}^y f_{X,Y}(x, y)xy dx dy = \frac{1}{2} \int_{-1}^1 \frac{y^2 - 1}{2} y dy =$

$$\frac{1}{4} \left( \frac{y^4}{4} - \frac{y^2}{2} \right) \Big|_{-1}^1 = 0.$$

$$E[e^{X+Y}] = \int_{-1}^1 \int_{-1}^y f_{X,Y}(x,y) e^{x+y} dx dy = \frac{1}{2} \int_{-1}^1 (e^{2y} - e^{y-1}) dy = \left( \frac{1}{4} e^{2y} - \frac{1}{2} e^{y-1} \right) \Big|_{-1}^1 = \frac{1}{4} (e^2 - e^{-2}) - \frac{1}{2} (1 - e^{-2}) = \frac{1}{4} (e^2 - 2 + e^{-2}) = \left( \frac{e - \frac{1}{e}}{2} \right)^2.$$

**Problem 7.** First, if  $W < 1$ , then both  $\frac{X}{Y}, \frac{Y}{X}$  are less than 1.

Since  $\Pr[X \leq 0 \vee Y \leq 0] = 0$  by the definition of  $f_{X,Y}(x,y)$ .

$\therefore \frac{X}{Y} < 1, \frac{Y}{X} < 1 \Rightarrow X < Y, Y < X$ , which is impossible.

$\therefore$  there must be  $W \geq 1$ .

$$F_W(w) = \Pr[W \leq w] = \Pr\left[\max\left(\frac{X}{Y}, \frac{Y}{X}\right) \leq w\right] = \Pr\left[\frac{X}{Y} \leq w \wedge \frac{Y}{X} \leq w\right] = \Pr\left[Y \geq \frac{X}{w} \wedge X \geq \frac{Y}{w}\right] = 1 - \Pr\left[X < \frac{Y}{w}\right] - \Pr\left[Y < \frac{X}{w}\right].$$

Note that  $\frac{Y}{w} \leq \frac{a}{w} \leq a, \frac{X}{w} \leq \frac{a}{w} \leq a$ .

$$\begin{aligned} \therefore 1 - \Pr\left[X < \frac{Y}{w}\right] - \Pr\left[Y < \frac{X}{w}\right] &= 1 - \int_0^a \int_0^{\frac{y}{w}} f_{X,Y}(x,y) dx dy - \int_0^a \int_0^{\frac{x}{w}} f_{X,Y} dy dx = \\ &= 1 - 2 \int_0^a \int_0^{\frac{y}{w}} \frac{1}{a^2} dx dy = 1 - \frac{2}{a^2} \int_0^a \frac{y}{w} dy = 1 - \frac{1}{a^2 w} a^2 = 1 - \frac{1}{w}. \\ \Rightarrow f_W(w) &= F'_W(w) = \frac{1}{w^2}. \end{aligned}$$

**Problem 8.**

(a) Since at least one bus arrive, there is  $n \geq 1$ .

Since at most one bus arrives in a minute, there is  $t \geq n$ .

$\therefore$  the set is  $\{(n, t) : t \geq n \geq 1, n, t \in \mathbb{Z}\}$ .

(b) If  $n > t$ , then by (a),  $P_{N,T}(n, t) = 0$ .

If  $n \leq t$ , it means that exactly  $n - 1$  buses passed through in the first  $t - 1$  minutes, and a bus passed through at the  $t$ -th minute, so the probability is  $\binom{t-1}{n-1} p^n (1-p)^{t-n}$ .

The probability that I didn't board the first  $n-1$  buses but the  $n$ -th is  $(1-q)^{n-1} q$ .

$$\therefore P_{N,T}(n, t) = \begin{cases} \binom{t-1}{n-1} p^n (1-p)^{t-n} (1-q)^{n-1} q, & \text{if } t \geq n \geq 1, n, t \in \mathbb{Z} \\ 0, & \text{otherwise} \end{cases}.$$

(c)  $P_N(n)$  = the probability that I didn't board the first  $n - 1$  buses but the  $n$ -th

$$= (1 - q)^{n-1}q.$$

$P_T(t)$  = the probability that in  $t$  minutes:

(1) the first  $t - 1$  minutes either the bus didn't come, or I didn't board the bus, which has probability  $(1 - pq)$ .

(2) the  $t$ -th minute the bus came and I boarded the bus, which has probability  $pq$ .

$$\therefore P_T(t) = (1 - pq)^{t-1}pq.$$

$$(d) \quad P_{N|T}(n|t) = \frac{P_{N,T}(n,t)}{P_T(t)} = \begin{cases} \frac{\binom{t-1}{n-1} p^n (1-p)^{t-n} (1-q)^{n-1} q}{(1-pq)^{t-1} pq}, & \text{if } n \leq t \\ 0, & \text{otherwise} \end{cases}$$

$$P_{T|N}(t|n) = \frac{P_{N,T}(n,t)}{P_N(n)} = \begin{cases} \frac{\binom{t-1}{n-1} p^n (1-p)^{t-n} (1-q)^{n-1} q}{(1-q)^{n-1} q} = \binom{t-1}{n-1} p^n (1-p)^{t-n}, & \text{if } n \leq t \\ 0, & \text{otherwise} \end{cases}$$

### Problem 9.

(a)  $P_N(n) = 0$  for  $n < 0$ .

$$\text{For } n \geq 0, P_N(n) = \sum_{k=0}^n \frac{100^k e^{-100}}{(k+1)!} = (n+1) \times \frac{100^n e^{-100}}{(n+1)!} = \frac{100^n e^{-100}}{n!}.$$

$$\therefore P_N(n) = \begin{cases} \frac{100^n e^{-100}}{n!}, & \text{if } n \in \mathbb{Z}^+ \cup \{0\} \\ 0, & \text{otherwise} \end{cases}.$$

$$P_{K|N}(k|n) = \frac{P_{N,K}(n,k)}{P_N(n)} = \begin{cases} \frac{\frac{100^n e^{-100}}{n!}}{\frac{100^n e^{-100}}{(n+1)!}} = \frac{1}{n+1}, & \text{if } k = 0, 1, \dots, n; n = 0, 1, \dots \\ 0, & \text{otherwise} \end{cases}.$$

$$(b) \quad E[K|N = n] = \sum_{k=0}^n k P_{K|N}(k|n) = \sum_{k=0}^n k \times \frac{1}{n+1} = \frac{n}{2}.$$

$$(c) \quad E[K|N] = \frac{N}{2}.$$

$$E[K] = E[E[K|N]] = E\left[\frac{N}{2}\right] = \sum_{n=0}^{\infty} \frac{n}{2} \frac{100^n e^{-100}}{n!} = \sum_{n=1}^{\infty} \frac{100^n e^{-100}}{(n-1)! \times 2} = 50 \sum_{n=1}^{\infty} \frac{100^{n-1} e^{-100}}{(n-1)!} =$$

$$50 \sum_{n=0}^{\infty} \frac{100^n e^{-100}}{n!} = 50 \sum_{n=0}^{\infty} P_N(n) = 50.$$

### Problem 10.