JUNE · IAS 2026 · CLASS 14

Talking the Discussion

The Discussion is new this week. Tonight we don't edit it — we say it out loud until it sounds like you.



FROM THE PAPER

Balance no time-by-group interaction, static or dynamic

Trunk rotation FCG −2.6° at contact, −4.0° at peak

Direction (+) toward the standing leg = the compensation

PART A · 20 MIN

The Skeleton

No numbers. Four sentences that carry the whole result.

BRIAN ASKS

- In one sentence — what happened to balance?
- And what did change?
- Which direction did the trunk move?
- Why does the direction matter?
- Say the whole thing again, no numbers, thirty seconds.

JUNE GETS COMFORTABLE WITH

Neither group improved balance more than the control.

What did change was trunk rotation — only in the foot core group.

Less rotation toward the standing leg.

Rotating toward the standing leg is how CAI patients compensate.

WE SAY



PART B · 25 MIN

Defending the Nulls

Brian is a curious colleague, not an examiner. June explains why nothing else moved.

BRIAN ASKS

- Isn't four weeks just too short?
- So foot core training does nothing for balance?
- The stability score didn't move but the trunk did — what does that actually mean?
- Why test standing still if you expected it to be too easy?
- Would six weeks have changed the picture?

JUNE GETS COMFORTABLE WITH

That's fair — four weeks is short. What we can say is …

It's not that it didn't work; the static task may not be sensitive enough to show it.

The movement strategy changed before the stability score did.

Other studies with the same exercises found the same thing.

WE SAY

FROM THE PAPER

Static Drewes (4-wk SFE) and Lynn (4-wk SFE/towel curl) — no change either

Dynamic earlier studies used reach tests, not landing

Lower limb 4 weeks + exercises that don't look like landing



PART C · 15 MIN

The Awkward One

The trunk core group rotated more after training. Nobody will let that go.

BRIAN ASKS

- Your trunk core group rotated more after training — did trunk training make it worse?
- If a reviewer asks that on stage, what do you say?
- Is it noise, or is it real?
- Could trunk training make people use the trunk more on purpose?
- How do you say "I don't know" and still sound like the expert?

JUNE GETS COMFORTABLE WITH

We don't have a clean explanation for that.

One possibility is … but with ten per group, I'd be careful reading too much into it.

The clear difference was between the two training groups.

That's something I'd want to test with a bigger sample.


WE SAY

FROM THE PAPER

TCG change +4.0° at contact, +4.6° at peak

CG change +1.4° / +2.0°

Paper offers no explanation for the TCG increase



PART D · 10 MIN

One Line

If the room remembers one thing — which one?

BRIAN ASKS

- If the audience remembers one sentence, what is it?
- Option 1: Train the foot, and the trunk changes.
- Option 2: One component isn't enough for CAI.
- Which one does your data actually support?
- Finish this: "If you remember one thing from this talk …"

JUNE GETS COMFORTABLE WITH

We trained the foot, and the trunk changed.

People with CAI rotate the trunk to stay upright — foot core training seemed to reduce the need for that.

We think this is a bottom-up effect along the kinetic chain.

If you remember one thing from this talk, …

WE SAY

FROM THE PAPER

Conclusion foot core training may influence proximal movement control

Discussion ending single-component training may be insufficient

Both are in the paper only one fits in ten minutes



PART E · 15 MIN

Numbers Without Stumbling

Methods in forty seconds, said smoothly. Then once more, faster.

BRIAN ASKS

- How many people, how many groups?
- How long, how often?
- What did they do — foot group first, then trunk group.
- What did you measure, in three words each?
- Say all of it in forty seconds.

JUNE GETS COMFORTABLE WITH

Thirty people with CAI, ten per group, stratified by sex.

Twelve sessions — thirty minutes, three times a week, four weeks.

Five foot exercises, sitting to standing to single-leg. Five trunk exercises, longer holds each week.

Single-leg balance, a hurdle landing, and motion capture — 35 markers.

WE SAY

FROM THE PAPER

Hurdle 30 cm, start at 40% of height, land on the involved leg, hold 10 s

Window initial contact + first 200 ms

Stats mixed ANOVA + SPM across the curve