# Copper Rockfish (Sebastes caurinus) along the California North of Pt. Conception US West coast in 2020

by Chantel R. Wetzel<sup>1</sup> Brian J. Langseth<sup>1</sup> Jason M. Cope<sup>1</sup> John E. Budrick<sup>2</sup>

<sup>&</sup>lt;sup>1</sup>Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake Boulevard East, Seattle, Washington 98112

 $<sup>^2</sup>$  California Department of Fish and Wildlife, 350 Harbor Boulevard, Belmont, California 94002

© Pacific Fisheries Management Council, 2020

Wetzel, C.R., B.J. Langseth, J.M. Cope, J.E. Budrick. 2020. Copper Rockfish (*Sebastes caurinus*) along the California North of Pt. Conception US West coast in 2020. Pacific Fisheries Management Council, Portland, Oregon. 9 p.

Correct citation for this publication:

# Contents

$\mathbf{E}_{2}$	cecut	ive Su	mmary	i
	Stoc	k		i
	Lane	dings		i
	Data	a and A	Assessment	i
	Stoc	k Biom	ass	i
	Reci	ruitmen	ıt	i
	Exp	loitatio	n Status	i
	Refe	erence F	Points	i
	Man	agemer	nt Performance	ii
	Unr	esolved	Problems and Major Uncertainties	ii
	Deci	ision Ta	able	ii
	Rese	earch ar	nd Data Needs	ii
1	Intr	oducti	ion	1
	1.1	Basic	Information	1
	1.2	Life H	istory	1
	1.3	Ecosys	stem Considerations	1
	1.4	Histor	ical and Current Fishery Information	1
	1.5	Summ	nary of Management History and Performance	1
	1.6	Foreig	n Fisheries	2
<b>2</b>	Dat	a		2
	2.1	Fisher	y-Dependent Data	2
	2.2	Fisher	y-Independent Data	2
		2.2.1	AFSC Slope Survey	2
		2.2.2	AFSC/NWFSC West Coast Triennial Shelf Survey	2
		2.2.3	NWFSC West Coast Groundfish Bottom Trawl Survey	3
	2.3	Biolog	cical Data	3
		2.3.1	Natural Mortality	3
		2.3.2	Maturation and Fecundity	3
		2.3.3	Sex Ratio	4
		2.3.4	Length-Weight Relationship	4
		2.3.5	Growth (Length-at-Age)	4

2020 iii

		2.3.6	Ageing Precision and Bias	4
	2.4	Enviro	onmental and Ecosystem Data	4
3	Ass	essmei	nt Model	4
	3.1	Summ	nary of Previous Assessments and Reviews	4
		3.1.1	History of Modeling Approaches (not required for an update assessment)	4
		3.1.2	Most Recent STAR Panel and SSC Recommendations (not required for an update assessment)	4
		3.1.3	Response to Groundfish Subcommittee Requests (not required in draft)	4
	3.2	Model	Structure and Assumptions	4
		3.2.1	Model Changes from the Last Assessment (not required for an update assessment)	4
		3.2.2	Modeling Platform and Structure	Ē
		3.2.3	Model Parameters	Ę
		3.2.4	Key Assumptions and Structural Choices	Ę
	3.3	Base 1	Model Results	Ę
		3.3.1	Parameter Estimates	Ę
		3.3.2	Fits to the Data	Ę
		3.3.3	Population Trajectory	Ę
		3.3.4	Reference Points	Ę
	3.4	Model	Diagnostics	Ę
		3.4.1	Convergence	6
		3.4.2	Sensitivity Analyses	6
		3.4.3	Retrospective Analysis	6
		3.4.4	Likelihood Profiles	6
		3.4.5	Unresolved Problems and Major Uncertainties	6
4	Ma	nagem	ent	6
	4.1	Refere	ence Points	6
	4.2	Unrese	olved Problems and Major Uncertainties	6
	4.3	Harve	st Projections and Decision Tables	6
	4.4	Evalua	ation of Scientific Uncertainty	$\epsilon$
	4.5	Resear	rch and Data Needs	(
5	Ack	nowle	dgments	6

iv

6	References	7
7	Tables	8
8	Figures	9

v

TIGE OF FIGURES	TIGHT OF HART PO
LIST OF FIGURES	LIST OF TABLES

List of	${f Figures}$
---------	---------------

# List of Tables

2020 vi

# **Executive Summary**

Reference Points

Replace text.

Stock			
This assessment reports the status of Copper Rockfish (Sebastes caurinus) off the California North of Pt. Conception US West coast using data through xxxx.			
Landings			
Replace text.			
Data and Assessment			
Replace text.			
Stock Biomass			
Replace text.			
Recruitment			
Replace text.			
Exploitation Status			
Replace text.			

Management Performance		
Replace text.		
Unresolved Problems and Major Uncertainties		
Replace text.		
Decision Table		
Replace text.		
Research and Data Needs		
Replace text.		

### Introduction

1.1 Basic	Information
-----------	-------------

This assessment reports the status of Copper Rockfish ( $Sebastes\ caurinus$ ) off the	California
North of Pt. Conception US West coast using data through xxxx.	

This assessment reports the status of Copper Rockfish (Sebastes caurinus) off the Calife North of Pt. Conception US West coast using data through xxxx.		
1.2 Life History		
Replace text.		
1.3 Ecosystem Considerations		
Replace text.		
1.4 Historical and Current Fishery Information		
Replace text.		
1.5 Summary of Management History and Performance		

Replace text.

#### 1.6 Foreign Fisheries

Replace text.

#### 2 Data

A description of each data source is provided below (Figure 1).

#### 2.1 Fishery-Dependent Data

#### 2.2 Fishery-Independent Data

#### 2.2.1 AFSC Slope Survey

The AFSC Slope Survey (Slope Survey) operated during the months of October to November aboard the R/V  $Miller\ Freeman$ . Partial survey coverage of the US west coast occurred during the years 1988-1996 and complete coverage (north of  $34^{\circ}30'$ S) during the years 1997 and 1999-2001. Typically, only these four years that are seen as complete surveys are included in assessments.

#### 2.2.2 AFSC/NWFSC West Coast Triennial Shelf Survey

The AFSC/NWFSC West Coast Triennial Shelf Survey (Triennial Survey) was first conducted by the Alaska Fisheries Science Center (AFSC) in 1977, and the survey continued until 2004 (Weinberg et al. 2002). Its basic design was a series of equally-spaced east-to-west transects across the continential shelf from which searches for tows in a specific depth range were initiated. The survey design changed slightly over time. In general, all of the surveys were

conducted in the mid summer through early fall. The 1977 survey was conducted from early July through late September. The surveys from 1980 through 1989 were conducted from mid-July to late September. The 1992 survey was conducted from mid July through early October. The 1995 survey was conducted from early June through late August. The 1998 survey was conducted from early June through early August. Finally, the 2001 and 2004 surveys were conducted from May to July.

Haul depths ranged from 91-457 m during the 1977 survey with no hauls shallower than 91 m. Due to haul performance issues and truncated sampling with respect to depth, the data from 1977 were omitted from this analysis. The surveys in 1980, 1983, and 1986 covered the US West Coast south to 36.8°N latitude and a depth range of 55-366 m. The surveys in 1989 and 1992 covered the same depth range but extended the southern range to 34.5°N (near Point Conception). From 1995 through 2004, the surveys covered the depth range 55-500 m and surveyed south to 34.5°N. In 2004, the final year of the Triennial Survey series, the Northwest Fisheries Science Center (NWFSC) Fishery Resource and Monitoring division (FRAM) conducted the survey following similar protocols to earlier years.

#### 2.2.3 NWFSC West Coast Groundfish Bottom Trawl Survey

The NWFSC West Coast Groundfish Bottom Trawl Survey (WCGBTS) is based on a random-grid design; covering the coastal waters from a depth of 55-1,280 m (Bradburn, Keller, and Horness 2011). This design generally uses four industry-chartered vessels per year assigned to a roughly equal number of randomly selected grid cells and divided into two 'passes' of the coast. Two vessels fish from north to south during each pass between late May to early October. This design therefore incorporates both vessel-to-vessel differences in catchability, as well as variance associated with selecting a relatively small number (approximately 700) of possible cells from a very large set of possible cells spread from the Mexican to the Canadian borders.

#### 2.3 Biological Data

#### 2.3.1 Natural Mortality

#### 2.3.2 Maturation and Fecundity

2.3.3	Sex Ratio
2.3.4	Length-Weight Relationship
2.3.5	Growth (Length-at-Age)
2.3.6	Ageing Precision and Bias
2.4	Environmental and Ecosystem Data
<b>3</b> A	Assessment Model
3.1	Summary of Previous Assessments and Reviews
3.1.1	History of Modeling Approaches (not required for an update assessment)
3.1.2	Most Recent STAR Panel and SSC Recommendations (not required for an update assessment)
3.1.3	Response to Groundfish Subcommittee Requests (not required in draft)
3.2	Model Structure and Assumptions
3.2.1	Model Changes from the Last Assessment (not required for an update assessment)

#### 3.2.2 Modeling Platform and Structure

General model specifications (e.g., executable version, model structure, definition of fleets and areas)

#### 3.2.3 Model Parameters

Describe estimated vs. fixed parameters, priors

#### 3.2.4 Key Assumptions and Structural Choices

#### 3.3 Base Model Results

#### 3.3.1 Parameter Estimates

#### 3.3.2 Fits to the Data

#### 3.3.3 Population Trajectory

#### 3.3.4 Reference Points

#### 3.4 Model Diagnostics

Describe all diagnostics

3.4.1	Convergence	
3.4.2	Sensitivity Analyses	
3.4.3	Retrospective Analysis	
3.4.4	Likelihood Profiles	
3.4.5	Unresolved Problems and Major Uncertainties	
4 Management		
<b>4.1</b> ]	Reference Points	
4.2	Unresolved Problems and Major Uncertainties	
4.3	Harvest Projections and Decision Tables	
4.4	Evaluation of Scientific Uncertainty	
<b>4.5</b>	Research and Data Needs	

Acknowledgments

Here are all the mad props!

#### 6 References

Bradburn, M. J., A. A Keller, and B. H. Horness. 2011. "The 2003 to 2008 US West Coast Bottom Trawl Surveys of Groundfish Resources Off Washington, Oregon, and California: Estimates of Distribution, Abundance, Length, and Age Composition." US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine Fisheries Service.

Weinberg, K. L., M. E. Wilkins, F. R. Shaw, and M. Zimmermann. 2002. "The 2001 Pacific West Coast Bottom Trawl Survey of Groundfish Resources: Estimates of Distribution, Abundance and Length and Age Composition." NOAA Technical Memorandum NMFS-AFSC-128. U.S. Department of Commerce.

# 7 Tables

# 8 Figures




Figure 1: Summary of data sources used in the base model.