

CLINICAL PERFORMANCE : 30-DAY HOSPITAL READMISSION ANALYTICS

High Risk Patients

7K

Total Encounters

27K

Total Readmitted


3.16K

Readmission Rate (%)


11.73%

Readmission Burden by Diagnosis


Risk Category Low Medium High




Readmission Risk by Discharge Pathway



Average Length of Stay



Risk Category



Clinical Insights

- Cardiometabolic patients represent the largest readmission burden.
- Polypharmacy (8+ medications) is associated with elevated readmission risk.
- Home discharge pathways account for the majority of return encounters.
- Older adults demonstrate the highest concentration of medication-related readmission risk.

Readmission Risk Heatmap: Age vs Medication Burden

Age range	Moderate (5-7)	Polypharmacy (8+)	Standard (0-4)
0-18	8	10	15
19-30	138	145	253
31-50	244	225	389
51 +	448	493	796

CLINICAL STRATEGY & OPERATIONAL EFFICIENCY

REDUCING AVOIDABLE HOSPITAL READMISSIONS THROUGH PREDICTIVE HEALTHCARE ANALYTICS

High Risk Readmission (%)

27.59

Cost Per Readmission

\$10K

Financial Exposure

\$31.64M

Savings Opportunity


\$4.746M

Operational Burden

27.58


Financial Exposure by Diagnosis

Discharge Pathway ● Facility/ Rehab ● Home



Readmission Cost Drivers

Primary Diagnosis ● Diabetes ● Hypertension ● Cardiovascular ● General Medical Condition




High-Risk Intervention Priority

Risk Group	Recommended Action
High Risk	48hr post-discharge follow-up
Medium Risk	Medication reconciliation
Low Risk	Standard discharge monitoring

Medication Burden vs Readmission Risk

Risk Category ● Low ● Medium ● High



KEY ANALYTIC INSIGHTS

- Patients aged 51+ discharged home show elevated readmission risk.
- Polypharmacy patients demonstrate higher 30-day readmission probability.
- Early follow-up programs may reduce high-risk readmissions.