

Lactase persistence DNA variant enhances lactase promoter activity *in vitro*: functional role as a *cis* regulatory element

Lynne C. Olds and Eric Sibley*

Department of Pediatrics, Stanford University Medical Center, Stanford, CA 94305, USA

Received May 2, 2003; Revised and Accepted July 7, 2003

Lactase persistence is a heritable, autosomal dominant, condition that results in a sustained ability to digest the milk sugar lactose throughout adulthood. The majority of the world's human population experiences a decline in production of the digestive enzyme lactase-phlorizin hydrolase during maturation. However, individuals with lactase persistence continue to express high levels of the lactase gene into adulthood. Lactase persistence has been strongly correlated with single nucleotide genetic variants, C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈, located 13.9 and 22 kb upstream from the lactase structural gene. We aimed to characterize a functional role for the polymorphisms in regulating lactase gene transcription. DNA in the region of the C/T₋₁₃₉₁₀ or G/A₋₂₂₀₁₈ human lactase variants was cloned upstream of the 3.0 kb rat lactase gene promoter in a luciferase reporter construct. Human intestinal Caco-2 cells were transfected with the lactase variant/promoter-reporter constructs and assayed for promoter activity. A 200 bp region surrounding the C₋₁₃₉₁₀ variant, associated with lactase non-persistence, results in a 2.2-fold increase in lactase promoter activity. The T₋₁₃₉₁₀ variant, associated with lactase persistence, results in an even greater 2.8-fold increase. The DNA sequence of the C/T₋₁₃₉₁₀ variants differentially interacts with intestinal cell nuclear proteins on EMSAs. AP2 co-transfection results in a similar repression of the C/T₋₁₃₉₁₀ variant/promoter-reporter constructs. The DNA region of the C/T₋₁₃₉₁₀ lactase persistence/non-persistence variant functions *in vitro* as a *cis* element capable of enhancing differential transcriptional activation of the lactase promoter. Such differential regulation by the C and T variants is consistent with a causative role in the mechanism specifying the lactase persistence/non-persistence phenotypes in humans.

INTRODUCTION

Intestinal lactase-phlorizin hydrolase (LPH, lactase) is the absorptive enterocyte membrane glycoprotein essential for digestive hydrolysis of lactose in milk. Lactase is present predominantly along the brush border membrane of differentiated enterocytes lining the villi of the small intestine. Expression of the lactase gene is temporally regulated during gut development and maturation (1,2). Lactase enzyme activity is maximal in the small intestine of pre-weaned mammals and declines markedly during maturation. In laboratory rodents, the maturational decline in lactase activity has been shown to occur at the time of weaning and contrasts with a maturational increase in enzymatic activity of other intestinal hydrolases essential for digestion of solid foods (comprehensively reviewed in 3). In humans, the activities of lactase and most of the other digestive hydrolases are maximal at birth. The age of onset of the

post-weaning maturational decline of lactase activity in humans is variable, ranging from the toddler years to young adulthood.

Lactose is the predominant carbohydrate in breast milk. The lactose disaccharide consists of a glucose and a galactose molecule linked by a $\beta(1-4)$ glycosidic bond. The digestive enzyme lactase catalyzes the hydrolysis of lactose to yield glucose and galactose that can then be absorbed across the intestinal mucosa membrane. The maturational decline in lactase activity renders most of the world's adult human population intolerant to milk and other dairy products (4,5). Lactose present in dairy products cannot be digested in the small intestine, due to the reduced lactase level, and instead is fermented by bacteria in the distal ileum and colon. The fermentative products result in symptoms of diarrhea, gas bloat, flatulence and abdominal pain.

In a minority of adults, high levels of lactase activity persist in adulthood. This hereditary persistence of lactase is common

*To whom correspondence should be addressed at: Department of Pediatrics, Stanford University School of Medicine, 750 Welch Road, Suite 116, Palo Alto, CA 943404, USA. Tel: +1 6507235070; Fax: +1 6504985608; Email: erc@stanford.edu

primarily in people of northern European descent and is attributed to inheritance of an unidentified autosomal dominant mutation that prevents the normal maturational decline in lactase expression (6). Enattah *et al.* (7), using linkage disequilibrium and haplotype analysis in humans, have recently reported the identification of genetic variants located -13 910 and -22 018 bp upstream of the human lactase gene that are associated with hereditary lactase persistence/non-persistence in Finnish families (7). The -13 910 bp C/T₋₁₃₉₁₀ variant is located 5' to the lactase gene within intron 13 of the adjacent *MCM6* gene (8) on chromosome 2. The intron 13 variant is a single nucleotide polymorphism, C to T. The authors report complete correlation between the lactase non-persistence phenotype and homozygosity for the C variant. Similarly, complete correlation is reported between the lactase persistence phenotype and the presence of the T variant allele. In the same study, a second variant (G/A₋₂₂₀₁₈) was associated with the lactase persistence trait. Although not completely associated, this second variant may be involved in regulating the lactase activity. The C/T₋₁₃₉₁₀ variant, however, is a strong candidate for the regulatory element involved in mediating lactase persistence. In a recent report, the lactase persistence allelic genotype T₋₁₃₉₁₀/A₋₂₂₀₁₈ has been shown to be associated with increased levels of lactase mRNA transcripts compared with the non-persistence allelic genotype C₋₁₃₉₁₀/G₋₂₂₀₁₈ (9). While strongly correlated with the lactase persistence/non-persistence phenotypes, the C/T₋₁₃₉₁₀ variant has not yet been shown to be causative of the traits and a functional mechanism has not been defined (10). Enattah *et al.* (7) have postulated that the C/T₋₁₃₉₁₀ variant affects an AP2 site, noting that the C allele associated with lactase non-persistence is in the consensus binding motif whereas the T variant disrupts this motif. In the present paper, we report that the C/T₋₁₃₉₁₀ lactase persistence/non-persistence variant functions as a *cis* element capable of directing differential transcriptional activation of the lactase promoter.

RESULTS

The lactase persistence C/T₋₁₃₉₁₀ variant region enhances lactase promoter activity in intestinal cell culture

We have previously demonstrated that a 3.0 kb rat lactase promoter fragment (11) is capable of driving transcription in Caco-2 cells, a human adenocarcinoma-derived cell-line that mimics a small intestinal enterocyte phenotype with respect to expression of several digestive hydrolases including lactase (12). In order to determine whether the DNA sequence surrounding the C/T₋₁₃₉₁₀ human lactase persistence variant is capable of regulating lactase promoter activity, Caco-2 cells were transfected with lactase promoter-reporter constructs harboring the C or T variant nucleotides. The constructs were generated by cloning the C/T₋₁₃₉₁₀ region of the human lactase promoter upstream of the 3.0 kb rat lactase promoter in the luciferase reporter plasmid pGL3Basic. The C/T₋₁₃₉₁₀ region (-14 017 to -13 800) was PCR-amplified from Caco-2 cell genomic DNA in order to generate the T variant reporter constructs p3kLac-T (forward orientation) and p3kLac-T'

(reverse orientation). Similarly the C/T₋₁₃₉₁₀ region was amplified from human RP11-34L23 BAC clone DNA to yield the C variant constructs p3kLac-C and p3kLac-C'. The C/T₋₁₃₉₁₀ region in each construct was sequenced to confirm differences only at the -13 910 single nucleotide variant. Transfected Caco-2 cell extracts were assayed for relative luciferase activity 48 h after transfection as shown in Figure 1.

Transfection with the lactase non-persistence associated C variant construct, p3kLac-C, results in a 2.2-fold induction of promoter activity driven by the 3 kb lactase promoter. Transfection with the lactase persistence-associated T variant construct, p3kLac-T, results in a 2.8-fold induction of promoter activity driven by the 3.0 kb lactase promoter. While both C/T₋₁₃₉₁₀ regions are capable of enhancing promoter activity, the lactase persistence-associated T variant possesses a significantly greater activity compared to the non-persistence-associated C variant ($P < 0.00001$, p3kLac-T versus p3kLac-C). Placement of the C/T₋₁₃₉₁₀ variant region in both forward and reverse orientations results in comparable activation of the lactase promoter. Differential enhancement of lactase promoter activity mediated by the C/T₋₁₃₉₁₀ variant region suggests a functional role for the DNA variant in specifying the lactase persistence/non-persistence phenotypes.

The G/A₋₂₂₀₁₈ variant region directs minimal enhancement of lactase promoter activity in intestinal cell culture

A second genomic variant, G/A₋₂₂₀₁₈, located upstream of the C/T₋₁₃₉₁₀ variant region has also been associated with the lactase persistence/non-persistence traits. Although not completely associated, this second variant may also have a role in regulating lactase promoter activity. To determine whether the G/A₋₂₂₀₁₈ variant region is involved in regulating the human lactase promoter, the G/A₋₂₂₀₁₈ region (22 133-21 910) was cloned in both orientations upstream of the 3.0 kb rat lactase promoter to generate p3kLac-G, p3kLac-G', p3kLac-A and p3kLac-A'. Transfection with the lactase non-persistence associated G variant constructs (p3kLac-G and p3kLac-G') or with the lactase persistence-associated A variant construct (p3kLac-A) results in minimal, ~1.2- to 1.4-fold, induction of promoter activity (Fig. 2). The A variant region cloned in the reverse orientation (p3kLac-A') results in an ~0.8-fold reduction in promoter activity.

To determine whether the G/A₋₂₂₀₁₈ variant region is capable of regulating the human lactase promoter in the context of the C/T₋₁₃₉₁₀ variant, the G/A₋₂₂₀₁₈ region was cloned upstream of the C/T₋₁₃₉₁₀ region to generate p3kLac-GC and p3kLac-AT. The G₋₂₂₀₁₈ variant and the C₋₁₃₉₁₀ variant are both associated with lactase non-persistence. The addition of the G₋₂₂₀₁₈ region upstream of the C₋₁₃₉₁₀ region does not result in a significant change in lactase promoter activity (Fig. 3). The A₋₂₂₀₁₈ variant and the T₋₁₃₉₁₀ variant are both associated with lactase persistence. Addition of the A₋₂₂₀₁₈ region upstream of the T₋₁₃₉₁₀ region results in a minimal, ~1.2-fold, increase in lactase promoter activity. Thus, the G/A₋₂₂₀₁₈ variant region appears to direct minimal enhancement of the lactase promoter in context of the C/T₋₁₃₉₁₀ variant region in the transfected reporter constructs.




Figure 1. Luciferase activity of intestinal cells transfected with C/T₋₁₃₉₁₀ variant region lactase promoter-reporter constructs. Caco-2 cells were transfected with pGL3, promoterless luciferase reporter construct, or with lactase promoter-luciferase reporter constructs and assayed for luciferase expression. p3kLac, 3.0 kb of the rat lactase promoter cloned in pGL3. p3kLac-C, p3kLac-C', p3kLac-T and p3kLac-T', human lactase gene C/T₋₁₃₉₁₀ variant region (open box) harboring a C or T corresponding to nucleotide -13 910 cloned in forward or reverse (arrow) orientation in p3kLac. Transfection efficiencies were normalized to renilla luciferase expression from a co-transfected pRL-CMV vector and expressed as relative luciferase activity (means ± SD, *n* = 3). The values for p3kLac, p3kLac-C and p3kLac-T represent data from three experiments with transfections performed in triplicate, *n* = 9. **P* < 0.00001 versus p3kLac-C.

AP2 overexpression represses lactase promoter activity in intestinal cell culture

It has been postulated that the C/T₋₁₃₉₁₀ variant may affect an AP2 transcription factor binding site. The C allele, associated with lactase non-persistence, is in the consensus binding motif whereas the T variant disrupts this motif (7). In order to determine whether the transcription factor AP2 is involved in mediating the C/T₋₁₃₉₁₀ region transcriptional activation of the lactase promoter, Caco-2 cells were co-transfected with the C/T₋₁₃₉₁₀ variant/promoter constructs and AP2 α and AP2 γ expression constructs (Fig. 4). Co-transfection with the AP2 α or AP2 γ construct results in a 1.7-fold or greater repression in promoter activity driven by both C/T₋₁₃₉₁₀ variant/promoter-reporter constructs as well as the 3.0 kb rat lactase promoter. This result suggests that AP2 regulatory *cis* element sequences may reside in the proximal rat lactase promoter but does not support a role for AP2 in mediating the lactase C/T₋₁₃₉₁₀ variant effect.

Differential interaction between the C/T₋₁₃₉₁₀ variant *cis* elements and intestinal cell nuclear proteins

In order to identify interaction between the C/T₋₁₃₉₁₀ variant *cis* element and intestinal cell nuclear proteins, we employed the electrophoretic mobility shift assay. C and T variant binding

region oligonucleotide probes were incubated in the presence of Caco-2 cell (~90% confluent) nuclear extract and then migrated through a non-denaturing acrylamide gel. The EMSA in Figure 5 reveals the position of the rapidly migrating unbound probe at the base of the gel and DNA-protein complexes of slower mobility formed after incubation with Caco-2 nuclear extract in the absence of competition (lanes 4 and 5). A slow migrating DNA-protein complex, indicated by *arrow*, is more abundant in incubation with the T₋₁₃₉₁₀ *cis* element probe (lane 4) compared with the C₋₁₃₉₁₀ probe (lane 5). The T₋₁₃₉₁₀ element DNA-protein complex is competed by 200-fold excess unlabeled T₋₁₃₉₁₀ oligonucleotide (lane 3), but is not competed away by 200-fold excess of an unrelated Sp1 oligonucleotide (lane 2). The nuclear protein bound to the T₋₁₃₉₁₀ *cis* element probe in Caco-2 cells therefore represents a specific DNA-protein interaction. In contrast, the C₋₁₃₉₁₀ element DNA-protein complex in the same region is much less abundant and is not competed away by 200-fold excess C₋₁₃₉₁₀ (lane 6) or Sp1 (lane 7) oligonucleotides. To confirm specificity of binding to the T₋₁₃₉₁₀ *cis* element, EMSAs reactions were also performed with the T₋₁₃₉₁₀ *cis* element probe in the presence of C₋₁₃₉₁₀ *cis* element competitor DNA. The T₋₁₃₉₁₀ *cis* element DNA-protein complex was not competed by 200-fold excess C₋₁₃₉₁₀ *cis* element oligonucleotide (Fig. 6, lane 1). The nuclear protein interacting with the T₋₁₃₉₁₀ *cis* element probe is thus a candidate factor that may be involved in differential transcriptional activation mediated by the C/T₋₁₃₉₁₀




Figure 2. Luciferase activity of intestinal cells transfected with G/A₋₂₂₀₁₈ variant region lactase promoter-reporter constructs. Caco-2 cells were transfected with pGL3, promoterless luciferase reporter construct, or with lactase promoter-luciferase reporter constructs and assayed for luciferase expression. p3kLac, 3.0 kb of the rat lactase promoter cloned in pGL3. p3kLac-G, p3kLac-G', p3kLac-A and p3kLac-A', human lactase gene G/A₋₂₂₀₁₈ variant region (gray box) harboring a G or A corresponding to nucleotide -22018 cloned in forward or reverse (arrow) orientation in p3kLac. Transfection efficiencies were normalized to renilla luciferase expression from a co-transfected pRL-CMV vector and expressed as relative luciferase activity (means \pm SD, $n = 3$).

variant region. EMSA reactions performed with nuclear extract isolated from 7 day post-confluent Caco-2 cells result in a comparable T₋₁₃₉₁₀ *cis* element specific DNA-protein complex and AP2 antisera does not detect proteins interacting with the C/T₋₁₃₉₁₀ elements on gel supershifts (data not shown).

DISCUSSION

Hereditary persistence of lactase is common primarily in people of northern European descent and has been attributed to inheritance of a hitherto unidentified autosomal dominant mutation that prevents the normal maturational decline in lactase expression (6). The recent identification by Enattah *et al.* (7) of genetic variants associated with lactase persistence has prompted interest in determining whether the single nucleotide variants are causative. The intron 13 variant is a single nucleotide polymorphism (C to T) located at nucleotide -13910 relative to the lactase start site and was completely correlated with lactase persistence/non-persistence. With respect to regulation of the lactase persistence/non-persistence phenotype, previous studies have suggested both transcriptional and post-transcriptional control mechanisms (6,13-18).

In the present study, we have focused on characterization of the role of the C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈ regions in regulating the lactase gene promoter in intestinal cell culture. Specifically, the DNA surrounding the C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈ polymorphisms

was assayed for the ability to function as a transcriptional activator or repressor in transfected Caco-2 cells. With respect to the C/T₋₁₃₉₁₀ region, DNA fragments harboring both the C and T variants are capable of enhancing lactase promoter (Fig. 1). Of note, however, the T variant results in greater promoter activation as compared to the C variant. Such an enhanced activity mediated by the T variant would be consistent with a functional role in the mechanism of lactase persistence. It is possible that a more dramatic effect was not seen due to the nature of transient transfection studies. Transfected Caco-2 cells contain multiple non-replicating episomal copies of the reporter constructs. Transiently transfected constructs are not always subject to regulatory mechanisms that involve complex chromatin structure, in contrast to the diploid chromosomal complement of the endogenous gene (19). In addition, the transfection constructs assayed likely lack other essential co-regulatory sequences located outside the 200 bp C/T₋₁₃₉₁₀ region. Caco-2 cells, while a model of intestinal cell differentiation, are not identical to adult enterocytes *in vivo* and as such may also lack essential co-regulatory proteins needed to specify maximal phenotypic expression patterns. Thus, the C/T₋₁₃₉₁₀ variants may exert a significantly greater differential regulatory effect *in vivo* in the context of the endogenous lactase gene. Compared with the C/T₋₁₃₉₁₀ region, the G/A₋₂₂₀₁₈ variant region results in minimal enhancement of the lactase promoter (Fig. 2). Unlike the C/T₋₁₃₉₁₀ variant, the G/A₋₂₂₀₁₈ variant was not completely correlated with the lactase




Figure 3. Luciferase activity of intestinal cells transfected with G/A₋₂₂₀₁₈ and C/T₋₁₃₉₁₀ variant region lactase promoter-reporter constructs. Caco-2 cells were transfected with lactase promoter-reporter constructs and assayed for luciferase expression. p3kLac-C, human lactase gene C₋₁₃₉₁₀ variant region (open box). p3kLac-G-C, human lactase G₋₂₂₀₁₈ variant region (gray box) cloned in p3kLac-C. p3kLac-T, human lactase gene T₋₁₃₉₁₀ variant region (open box). p3kLac-A-T, human lactase A₋₂₂₀₁₈ variant region (gray box) cloned in p3kLac-T. Transfection efficiencies were normalized to renilla luciferase expression from a co-transfected pRL-CMV vector and expressed as relative luciferase activity (means \pm SD, $n = 3$).

persistence/non-persistence phenotype in the original report by Enattah *et al.* (7). Therefore, the G/A₋₂₂₀₁₈ variant may be associated with but not causative of the lactase phenotypes.

Enattah *et al.* (7) have postulated that the C/T₋₁₃₉₁₀ variant affects an AP2 binding site, noting that the C allele, associated with lactase non-persistence, is in the consensus binding motif, whereas the T variant disrupts this motif. To determine whether AP2 overexpression differentially regulates the C/T₋₁₃₉₁₀ variant element, we co-transfected Caco-2 cells with the variant-promoter constructs and with AP2 α and AP2 γ . Interestingly, AP2 over-expression results in a repression of the lactase promoter reporter constructs (Fig. 4). However, the fold repression is comparable for both the C and T allelic constructs. In addition, AP2 antisera do not detect proteins interacting with the C/T₋₁₃₉₁₀ element on gel supershifts (data not shown). These results do not support a direct role for AP2 binding.

The DNA sequence surrounding the C/T₋₁₃₉₁₀ variants, however, does interact differentially with intestinal cell nuclear proteins on gel shift assays. A slow migrating DNA-protein complex is more abundant in gel shift reactions performed with the T₋₁₃₉₁₀ *cis* element probe than in reactions with the C₋₁₃₉₁₀ probe (Fig. 5, compare lanes 4 and 5). Such a differential gel shift binding pattern combined with the differential transactivation results suggests a possible mechanism to account for the lactase persistence/non-persistence phenotypes. Namely, the DNA sequence surrounding the T₋₁₃₉₁₀ variant may create a binding site for a transactivating

protein that is capable of enhancing lactase transcription in adults with lactase persistence, thereby preventing the maturational decline. Such a mechanism would be consistent with a dominant mode of inheritance for lactase persistence associated with the T₋₁₃₉₁₀ allelic polymorphism. In individuals with lactase non-persistence, i.e. homozygous for the C₋₁₃₉₁₀ variant, the C nucleotide may disrupt the transactivating protein binding site and thus not allow for sustained lactase transcription into adulthood. There are likely to be additional regulatory control regions that specify the maturational decline in lactase in these individuals. Studies looking at the effect of the variants in stably transfected cell lines or in living animals will be necessary to confirm a role for the C/T₋₁₃₉₁₀ variant region in specifying lactase persistence or non-persistence. In addition, to characterize a potential regulatory mechanism directed by the C/T₋₁₃₉₁₀ region, further studies must be directed at identification of nuclear proteins interacting with this DNA region.

MATERIALS AND METHODS

Materials and reagents

Restriction endonucleases were purchased from Life Technologies (Rockville, MD, USA). Reagents for PCR were obtained from Qiagen (Valencia, CA, USA). Oligonucleotides




Figure 4. AP2 repression of lactase promoter-reporter constructs in intestinal cells. Caco-2 cells were transfected with p3kLac, p3kLac-C or p3kLac-T in the presence of AP2 α (shaded bars, AP2 α /pcDNA3) or AP2 γ (open bars, AP2 γ /pcDNA3) expression plasmids. Cells transfected in the absence of AP2 α were co-transfected with the empty expression vector pcDNA3 as a control (black bars). Transfection efficiencies were normalized to renilla luciferase expression from a co-transfected pRL-CMV vector and expressed as relative luciferase activity (means \pm SD, $n = 3$).

were synthesized by the protein and nucleic acid (PAN) facility of the Stanford University Medical Center.

PCR amplification of the C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈ regions of the human lactase gene locus

A 218 bp fragment of the human lactase *LCT* gene C/T₋₁₃₉₁₀ variant region was PCR-amplified using a forward oligonucleotide corresponding to nt -14017 to -13994, 5'-AGACGTAAGTTACCATTAAATAC-3', and a reverse oligonucleotide corresponding to nt -13800 to -13821, 5'-CGTTAATACCCACTGACCTATC-3'. Both primers were synthesized with a 5' terminal *Mlu*I restriction site for subsequent cloning. The C/T₋₁₃₉₁₀ region (-14017 to -13800) was PCR-amplified from Caco-2 cell genomic DNA to generate the T variant region DNA fragment. Similarly the C/T₋₁₃₉₁₀ region was amplified from RP11-34L23, a human genomic DNA BAC clone (BACPAC Resources), to yield the C variant region DNA fragment. A 224 bp fragment of the lactase gene G/A₋₂₂₀₁₈ variant region was PCR-amplified using a forward oligonucleotide corresponding to nt -22133 to -22113, 5'-TAAGAACATTTTACTCTTC-3', and a reverse oligonucleotide corresponding to nt -21910 to -21930, 5'-AGAAAATGGGTTTTCGCCATG-3'. Both primers were synthesized with a 5' terminal *Kpn*I restriction site. The G/A₋₂₂₀₁₈ region (-22133 to -21910) was PCR-amplified from Caco-2 cell genomic DNA to generate the A variant region fragment. The G/A₋₂₂₀₁₈ region was amplified from RP11-34L23 BAC clone DNA to yield the G variant region fragment.

Subcloning of the lactase variant-promoter-reporter constructs

C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈ variant region promoter-reporter constructs were generated by cloning the PCR-amplified




Figure 5. Electrophoretic mobility shift assay with Caco-2 cell nuclear extract and C/T₋₁₃₉₁₀ probes. Caco-2 cell (~90% confluent) nuclear extract protein (2.0 μ g) was incubated with biotin-labeled T₋₁₃₉₁₀ or C₋₁₃₉₁₀ *cis* element oligonucleotide probe alone (lanes 4 and 5) or in the presence of 200 \times unlabeled specific (lanes 3 and 6) or non-specific Sp1 (lanes 2 and 7) oligonucleotides. Probe incubated in the absence of nuclear protein is shown in lanes 1 and 8. Samples were loaded on a 5% native acrylamide gel.

C/T₋₁₃₉₁₀ or G/A₋₂₂₀₁₈ variant region PCR products upstream of the lactase promoter in the previously described p3kLac construct (11). The p3kLac construct contains a 3.0 kb fragment of the rat lactase promoter cloned upstream of the luciferase cDNA in the reporter plasmid pGL3Basic (Promega). Specifically, the internal 218 bp *Mlu*I fragment of the C/T₋₁₃₉₁₀ region PCR products was cloned into p3kLac to generate p3kLac-C and p3kLac-T (forward orientation) and p3kLac-C' and p3kLac-T' (reverse orientation). Similarly, the 224 bp *Kpn*I fragment of the G/A₋₂₂₀₁₈ region PCR products was cloned into p3kLac to generate p3kLac-G, p3kLac-G', p3kLac-A and p3kLac-A'. The combination G/A₋₂₂₀₁₈ C/T₋₁₃₉₁₀ variant region reporter constructs were generated by cloning the C/T₋₁₃₉₁₀ region *Mlu*I fragments into p3kLac-G and p3kLac-A downstream of the G/A₋₂₂₀₁₈ region to yield p3kLac-G-C and p3kLac-A-T. Incorporation of the C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈ variant regions was confirmed by sequencing for all constructs generated.

Transient transfection assays

Caco-2 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) with 10% fetal bovine serum. Forty-eight hours prior to transfection the cells were split and 35 mm dishes were seeded with 2×10^5 cells. For each reporter construct, a DNA transfection mixture was prepared consisting of 0.4 pmol of the reporter construct, 0.1 μ g of pRL-CMV (Promega) as an internal control, and pBluescript KS+ II to adjust to a final of 2.1 μ g total DNA. The individual DNA mixtures were transfected in triplicate wells into cells (50–80% confluent) with Lipofectamine Reagent (BRL) according to the protocol of the manufacturer. For AP2 co-transfection experiments,




Figure 6. Electrophoretic mobility shift assay with T₋₁₃₉₁₀ element probe and C₋₁₃₉₁₀ element competition. Caco-2 cell (~90% confluent) nuclear extract protein (8.0 µg) was incubated with ³²P-labeled T₋₁₃₉₁₀ or C₋₁₃₉₁₀ *cis* element oligonucleotide probe alone (lanes 3 and 4) or in the presence of 100× unlabeled specific T₋₁₃₉₁₀ (lane 2) or C₋₁₃₉₁₀ (lane 1) oligonucleotides. Samples were loaded on a 4% native acrylamide gel.

0.2 pmol of AP2α/pcDNA3.1+ or AP2γ/pcDNA3.1+ (gift of R. J. Weigel, Stanford University) (20) was transfected along with 0.4 pmol of the luciferase reporter constructs. Cells were harvested 48 h following transfection (70–90% confluent) and luciferase activity was measured by the Dual-LuciferaseTM Reporter Assay System (Promega) as described by the manufacturer, in a Monolight 3010 luminometer. Transfection with the dual reporters (firefly luciferase for the lactase promoter–reporter plasmids and renilla luciferase for the pRL-CMV control) allowed for simultaneous expression and measurement of both reporter enzymes. Experimental lactase promoter–reporter activities were normalized to the activity of the pRL-CMV internal control and expressed as relative luciferase activity (means ± SD). Statistical significance (*P*-value) was determined using Student's unpaired *t*-test.

Electrophoretic mobility shift assays

Nuclear proteins from Caco-2 cells harvested at 90% confluency and 7 days post confluency were extracted using the Nuclear Extract Kit from Active Motif (Carlsbad, CA, USA). Caco-2 cell nuclear extract purchased from Active Motif was also assayed. Protein was quantified with the protein assay from BioRad

(Hercules, CA, USA). C/T₋₁₃₉₁₀ variant region oligonucleotides (21mers) were synthesized with either the T or C variant centrally located and annealed to produce double-stranded probes (5'-GAT AATGTAGC/TCCCTGGCCTC-3'). DNA–protein interactions were detected using the LightShift Chemiluminescent EMSA Kit (Pierce Biotechnology, Rockford, IL, USA) and 3' end-biotinylated probes. Specifically, nuclear extracts (2 µg) were incubated with 20 fmol double-stranded, biotinylated DNA probes and 500 ng of poly(deoxyinosinate-deoxycytidylate) in a buffer containing 10 mM Tris, 50 mM KCl, 1 mM DTT, 0.05% NP-40 and 5 mM MgCl₂ in a total volume of 20 µl for 25 min at room temperature. Complexes were separated in a 5% native polyacrylamide gel in 0.5× Tris-borate-EDTA buffer, transferred to a nylon membrane and cross-linked at 120 mJ/cm². Detection of the complexes was per the manufacturer's instructions. Competition EMSAs used 200-fold molar excess of unlabeled oligonucleotide or a non-specific competitor oligonucleotide, Sp1, 5'-GATC GGGGCGGGGCGGGGCGGGGCGATC-3'. EMSAs were also performed with corresponding ³²P labeled fill-in oligonucleotides generated as previously described (11) and visualized by autoradiography.

ACKNOWLEDGEMENTS

This work was supported by the National Institute of Diabetes and Digestive and Kidney Diseases grants DK60074 and DK60715.

REFERENCES

- Montgomery, R.K., Buller, H.A., Rings, E.H. and Grand, R.J. (1991) Lactose intolerance and the genetic regulation of intestinal lactase-phlorizin hydrolase. *FASEB J.*, **5**, 2824–2832.
- Rings, E.H., de Boer, P.A., Moorman, A.F., van Beers, E.H., Dekker, J., Montgomery, R.K., Grand, R.J. and Buller, H.A. (1992) Lactase gene expression during early development of rat small intestine. *Gastroenterology*, **103**, 1154–1161.
- Henning, S., Rubin, D.C. and Shulman, R.J. (1992) Ontogeny of the intestinal mucosa. In Johnson, L.R. (ed.), *Physiology of the Gastrointestinal Tract*. Raven, New York, pp. 571–610.
- Buller, H.A. and Grand, R.J. (1990) Lactose intolerance. *A. Rev. Med.*, **41**, 141–148.
- Sahi, T. (1994) Genetics and epidemiology of adult-type hypolactasia. *Scand. J. Gastroenterol. Suppl.*, **202**, 7–20.
- Wang, Y., Harvey, C.B., Pratt, W.S., Sams, V.R., Sarner, M., Rossi, M., Auricchio, S. and Swallow, D.M. (1995) The lactase persistence/non-persistence polymorphism is controlled by a cis-acting element. *Hum. Mol. Genet.*, **4**, 657–662.
- Enattah, N.S., Sahi, T., Savilahti, E., Terwilliger, J.D., Peltonen, L. and Jarvela, I. (2002) Identification of a variant associated with adult-type hypolactasia. *Nat. Genet.*, **30**, 233–237.
- Harvey, C.B., Wang, Y., Darmoul, D., Phillips, A., Mantei, N. and Swallow, D.M. (1996) Characterisation of a human homologue of a yeast cell division cycle gene, MCM6, located adjacent to the 5' end of the lactase gene on chromosome 2q21. *FEBS Lett.*, **398**, 135–140.
- Kuokkanen, M., Enattah, N.S., Oksanen, A., Savilahti, E., Orpana, A. and Jarvela, I.I. (2003) Transcriptional regulation of the lactase-phlorizin hydrolase gene by polymorphisms associated with adult-type hypolactasia. *Gut*, **52**, 647–652.
- Grand, R.J., Montgomery, R.K., Chitkara, D.K. and Hirschhorn, J.N. (2003) Changing genes; losing lactase. *Gut*, **52**, 617–619.
- Fang, R., Santiago, N.A., Olds, L.C. and Sibley, E. (2000) The homeodomain protein Cdx2 regulates lactase gene promoter activity during enterocyte differentiation. *Gastroenterology*, **118**, 115–127.
- Hauri, H.P., Sander, B. and Naim, H. (1994) Induction of lactase biosynthesis in the human intestinal epithelial cell line Caco-2. *Eur. J. Biochem.*, **219**, 539–546.

13. Escher, J.C., de Koning, N.D., van Engen, C.G., Arora, S., Buller, H.A., Montgomery, R.K. and Grand, R.J. (1992) Molecular basis of lactase levels in adult humans. *J. Clin. Invest.*, **89**, 480–483.
14. Fajardo, O., Naim, H.Y. and Lacey, S.W. (1994) The polymorphic expression of lactase in adults is regulated at the messenger RNA level. *Gastroenterology*, **106**, 1233–1241.
15. Lloyd, M., Mevissen, G., Fischer, M., Olsen, W., Goodspeed, D., Genini, M., Boll, W., Semenza, G. and Mantei, N. (1992) Regulation of intestinal lactase in adult hypolactasia. *J. Clin. Invest.*, **89**, 524–529.
16. Rossi, M., Maiuri, L., Fusco, M.I., Salvati, V.M., Fuccio, A., Auricchio, S., Mantei, N., Zecca, L., Gloor, S.M. and Semenza, G. (1997) Lactase persistence versus decline in human adults: multifactorial events are involved in down-regulation after weaning. *Gastroenterology*, **112**, 1506–1514.
17. Sebastio, G., Villa, M., Sartorio, R., Guzzetta, V., Poggi, V., Auricchio, S., Boll, W., Mantei, N. and Semenza, G. (1989) Control of lactase in human adult-type hypolactasia and in weaning rabbits and rats. *Am. J. Hum. Genet.*, **45**, 489–497.
18. Wang, Y., Harvey, C.B., Hollox, E.J., Phillips, A.D., Poulter, M., Clay, P., Walker-Smith, J.A. and Swallow, D.M. (1998) The genetically programmed down-regulation of lactase in children. *Gastroenterology*, **114**, 1230–1236.
19. Smith, C.L. and Hager, G.L. (1997) Transcriptional regulation of mammalian genes in vivo. A tale of two templates. *J. Biol. Chem.*, **272**, 27493–27496.
20. McPherson, L.A. and Weigel, R.J. (1999) AP2alpha and AP2gamma: a comparison of binding site specificity and trans-activation of the estrogen receptor promoter and single site promoter constructs. *Nucl. Acids Res.*, **27**, 4040–4049.