

# Twisted $M$ -theory and its perturbative quantization.

Surya and Brian

October 14, 2020

## Contents

<b>1</b>	<b>Definition of twisted supergravity</b>	<b>2</b>
1.1	The classical BV theory . . . . .	2
1.2	Moduli theoretic interpretation . . . . .	2
<b>2</b>	<b>Quantization of holomorphic-topological theories</b>	<b>2</b>
2.1	Effective renormalization . . . . .	2
2.2	The $11$ -dimensional theory . . . . .	2
2.3	The moduli space of quantizations . . . . .	2
<b>3</b>	<b>Relationship to the Type IIA string</b>	<b>2</b>
3.1	Kodaira–Spencer theory with potentials . . . . .	2
3.1.1	Warm-up: four-dimensional Kodaira–Spencer theory .	2
3.1.2	Eight-dimensional Kodaira–Spencer theory . . . . .	2
3.2	The Type IIA topological string . . . . .	2
3.3	Reduction of twisted supergravity . . . . .	2
3.3.1	Calabi–Yau compactifications . . . . .	2
3.4	Twisted supergravity on a three-fold . . . . .	2
<b>4</b>	<b>The non-minimal <math>G_2</math> twist</b>	<b>2</b>

- 1 Definition of twisted supergravity
  - 1.1 The classical BV theory
  - 1.2 Moduli theoretic interpretation
- 2 Quantization of holomorphic-topological theories
  - 2.1 Effective renormalization
  - 2.2 The 11-dimensional theory
  - 2.3 The moduli space of quantizations
- 3 Relationship to the Type IIA string
  - 3.1 Kodaira–Spencer theory with potentials
    - 3.1.1 Warm-up: four-dimensional Kodaira–Spencer theory
    - 3.1.2 Eight-dimensional Kodaira–Spencer theory
  - 3.2 The Type IIA topological string
  - 3.3 Reduction of twisted supergravity
    - 3.3.1 Calabi–Yau compactifications
  - 3.4 Twisted supergravity on a three-fold
- 4 The non-minimal G2 twist