

1 The Woods Hole Assessment Model (WHAM): a general state-space
2 assessment framework that incorporates time- and age-varying
3 processes via random effects and links to environmental covariates

4 Brian C. Stock¹, Timothy J. Miller¹

5 ¹brian.stock@noaa.gov, timothy.j.miller@noaa.gov, Northeast Fisheries Science Center, National Marine
6 Fisheries Service, 166 Water Street, Woods Hole, MA 02543, USA

Abstract

The rapid changes observed in many marine ecosystems that support fisheries pose a challenge to stock assessment and management predicated on time-invariant productivity and considering species in isolation. In single-species assessments, two main approaches have been used to account for productivity changes: allowing biological parameters to vary stochastically over time (empirical), or explicitly linking population processes such as recruitment (R) or natural mortality (M) to environmental covariates (mechanistic). Here, we describe the Woods Hole Assessment Model (WHAM) framework and software package, which combines these two approaches. WHAM can estimate time- and age-varying random effects on annual transitions in numbers at age (NAA), M , and selectivity, as well as fit environmental time-series with process and observation errors, missing data, and nonlinear links to R and M . WHAM can also be configured as a traditional statistical catch-at-age (SCAA) model in order to easily bridge from status quo models and test them against models with state-space and environmental effects, all within a single framework.

We fit models with and without (independent or autocorrelated) random effects on NAA, M , and selectivity to data from five stocks with a broad range of life history, fishing pressure, number of ages, and time-series length. Models that included random effects performed well across stocks and processes, especially random effects models with a two dimensional (2D) first-order autoregressive, AR(1), covariance structure over age and year. We conducted simulation tests and found negligible or no bias in estimation of important assessment outputs (SSB, F , stock status, and catch) when the operating and estimation models matched. However, bias in SSB and F was often non-trivial when the estimation model was less complex than the operating model, especially when models without random effects were fit to data simulated from models with random effects. Bias of the variance and correlation parameters controlling random effects was also negligible or slightly negative as expected. Our results suggest that WHAM can be a useful tool for stock assessment when environmental effects on R or M , or stochastic variation in NAA transitions, M , or selectivity are of interest. In the U.S. Northeast, where the productivity of several groundfish stocks has declined, conducting assessments in WHAM with time-varying processes via random effects or environment-productivity links may account for these trends and potentially reduce retrospective bias.

Keywords

state-space; stock assessment; random effects; time-varying; environmental effects; recruitment; natural mortality; Template Model Builder (TMB)

1 Introduction

The last two decades have increasingly seen a push for more holistic, ecosystem-based fisheries management (Larkin, 1996; Link, 2002). In part, this is due to the view that considering single species in isolation produces riskier and less robust outcomes long-term (Patrick and Link, 2015). In several high-profile cases, fisheries management has failed to prevent collapses because they did not reduce fishing pressure in responses to changes in natural mortality (M), recruitment, or migration patterns caused by dynamics external to the stock in question (Northern cod: Shelton et al., 2006; Rose and Rowe, 2015; Pacific sardine: Zwolinski and Demer, 2012; Gulf of Maine cod: Pershing et al., 2015). This is particularly concerning in the context of climate change and the wide range of biological processes—often assumed to be constant—in stock assessments that are likely to be affected (Stock et al., 2011; Tommasi et al., 2017).

One approach to account for changing productivity is to explicitly link population processes to environmental covariates in single-species stock assessments, i.e. the mechanistic approach *sensu* Punt et al. (2014). Traditional single-species assessments assume that most population dynamics processes are constant in time, even though fisheries scientists have long known about important drivers of time-varying population processes, e.g. recruitment, mortality, growth, and movement (Garstang, 1900; Hjort, 1914). Effects of the environment or interactions with other species can be considered contextually, rather than explicitly as estimated parameters (although temporal variation in empirical weight and maturity at age can affect reference points). Despite how counterintuitive this may seem to ecologists and oceanographers who study such relationships, the evidence for direct linkages to specific environmental covariates is often weak and can break down over time (McClatchie et al., 2010; Myers, 1998). Additionally, the primary goal of most assessments is to provide management advice on near-term sustainable harvest levels—not to explain ecological relationships. Even if an environmental covariate directly affects fish productivity, including the effect in an assessment may not improve management advice if the effect is weak (De Oliveira and Butterworth, 2005). Moreover, including environmental effects in an assessment or management system has been shown to actually provide worse management in some cases (De Oliveira and Butterworth, 2005; Punt et al., 2014; Walters and Collie, 1988). This can be true even in cases of relatively well-understood mechanistic links between oceanic conditions and fish populations, as in the case of sea surface temperature and Pacific sardine (Hill et al., 2018; Zwolinski and Demer, 2012). Still, incorporating mechanistic environment-productivity links in assessments does have the potential to reduce residual variance, particularly in periods when few demographic data exist (Miller et al., 2016; Shotwell et al., 2014).

An alternative approach is to allow biological parameters to vary stochastically over time, i.e. the empirical

approach *sensu* Punt et al. (2014). In this case, the variation is caused by a range of sources that are not explicitly modeled. Statistical catch-at-age (SCAA) models typically only estimate year-specific fishing mortality (F_y) and recruitment (R_y), often as deviations from a mean, R_0 , that may or may not be a function of spawning biomass, e.g. $\log R_y = \log R_0 + \epsilon_y$. The main reason that other parameters are assumed constant is simply that there are not enough degrees of freedom to estimate many time-varying parameters. A common solution is to penalize the deviations, e.g. $\epsilon_y \sim \mathcal{N}(0, \sigma_\epsilon^2)$, although the penalty terms, σ_ϵ^2 , must be fixed or iteratively tuned and are therefore somewhat subjective (Aeberhard et al., 2018; Methot and Taylor, 2011; Methot and Wetzel, 2013; Xu et al., 2020). State-space models that treat parameters as unobserved states can, in principle, avoid such subjectivity by estimating the penalty terms as variance parameters constraining random effects and maximizing the marginal likelihood (Thorson, 2019). In this way, state-space models can allow processes to vary in time while simultaneously estimating fewer parameters.

Although state-space stock assessments have existed for some time (Gudmundsson, 1994; Mendelsohn, 1988; Sullivan, 1992), the recent development of Template Model Builder (TMB, Kristensen et al., 2016) software to perform efficient Laplace approximation has greatly expanded their use (Cadigan, 2016; Miller et al., 2016; Nielsen and Berg, 2014). In addition to the key advantage of objectively estimating variance, or “data weighting”, parameters, state-space models naturally predict unobserved states, and therefore handle missing data and short-term projections in a straightforward way (ICES, 2020). In comparisons with SCAA models, state-space models have been shown to have larger, more realistic, uncertainty and lower retrospective bias (Miller and Hyun, 2018; Stock et al., 2021).

Retrospective bias can occur when changing environmental conditions lead to changes in productivity that are unaccounted for in stock assessments, and this is a concern common to several groundfish stocks on the Northeast U.S. Shelf (Brooks and Legault, 2016; Tableau et al., 2018). The Northeast U.S. Shelf ecosystem is rapidly changing, and this has motivated managers to make the “continue[d] development of stock assessment models that include environmental terms” a top priority (Hare et al., 2016). Applications of state-space models with environmental effects on recruitment, growth, M , and maturity have proven promising (Miller and Hyun, 2018; Miller et al., 2018; O’Leary et al., 2019; Xu et al., 2018). In addition to providing short-term (1-3 years) catch advice with reduced retrospective bias, it is hoped that environment-linked assessments will help create realistic evaluation of sustainable stock and harvest levels in the medium-term (3-10 years) for stocks that have not rebounded in response to dramatic decreases in F .

To address the needs of fisheries management in a changing climate, we seek an assessment framework that combines both the empirical and mechanistic approaches. Namely, it should be able to 1) estimate time-varying parameters as random effects (i.e. a state-space model), and 2) include environmental effects

directly on biological parameters. The framework should also allow for easy testing against status quo SCAA models to ease gradual adoption through the “research track” or “benchmark” assessment process (Lynch et al., 2018). The objectives of this manuscript are to introduce the Woods Hole Assessment Model (WHAM) framework and demonstrate its ability to:

1. estimate time- and age-varying random effects on annual changes in abundance at age, M , and selectivity;
2. fit environmental time-series with process and observation error, missing data, and a link to a population process;
3. project processes treated as random effects in short-term population forecasts; and
4. simulate new data and random effects to conduct self- and cross-tests (*sensu* Deroba et al., 2015) to estimate bias in parameters and derived quantities.

Throughout, we describe how the above are implemented using the open-source WHAM software package (Miller and Stock, 2020).

2 Methods

2.1 Model description

WHAM is a generalization and extension of Miller et al. (2016) in TMB. It is in many respects similar to the Age-Structured Assessment Program (ASAP, Legault and Restrepo, 1998; Miller and Legault, 2015) and can be configured to fit statistical catch-at-age models nearly identically. There is functionality built into WHAM to migrate ASAP input files to R inputs needed for WHAM, and WHAM uses many of the same types of data inputs, such as empirical weight-at-age, so that existing assessments in the U.S. Northeast can be easily replicated and tested against models with state-space and environmental effects in a single framework.

2.1.1 Processes with random effects

WHAM primarily diverges from ASAP through the implementation of random effects on three processes: inter-annual transitions in numbers at age (NAA), natural mortality (M), and selectivity (s), as well as allowing effects of environmental covariates ($Ecov$) on recruitment and natural mortality (but see ASAP4; Miller and Legault, 2015). The environmental covariates and their observations are part of the state-space framework with true, unobserved values treated as random effects and observation on them having error. Other than environmental covariates, the processes are assumed to have a two dimensional (2D) first-order autoregressive,

AR(1), covariance structure over age and year, although correlation in either or both dimensions can be turned off. The 2D AR(1) structure has been widely used to model deviations by age and year in the parameters $F_{a,y}$ (Kumar et al., 2020; Nielsen and Berg, 2014; Perreault et al., 2020), $M_{a,y}$ (Cadigan, 2016; Stock et al., 2021), $s_{a,y}$ (Xu et al., 2019), and $N_{a,y}$ (Stock et al., 2021), as well as in the catch ($C_{a,y}$) and survey index ($I_{a,y}$) observations (Berg and Nielsen, 2016). Although other covariance structures are plausible (e.g. compound symmetry, unconstrained, or order- k autoregressive; Nielsen and Berg, 2014; Berg and Nielsen, 2016), the 2D AR(1) structure is a simple option which requires few parameters and allows for smooth deviations in time and by age; values for nearby years and ages are more correlated than distant ones. This is often a reasonable null hypothesis for biological or fishery processes, but there are also clear examples where change occurs abruptly, e.g. a disease outbreak increases M or changes in gear or area regulations affect selectivity (ICES, 2017a).

2.1.1.1 Numbers at age (NAA)

The stock equations in WHAM that describe the transitions between numbers at age are identical to Nielsen and Berg (2014) and Miller et al. (2016):

$$\log N_{a,y} = \begin{cases} \log(f(SSB_{y-1})) + \varepsilon_{1,y}, & \text{if } a = 1 \\ \log(N_{a-1,y-1} - Z_{a-1,y-1} + \varepsilon_{a,y}), & \text{if } 1 < a < A \\ \log(N_{A-1,y-1}e^{-Z_{A-1,y-1}} + N_{A,y-1}e^{-Z_{A,y-1}}) + \varepsilon_{A,y}, & \text{if } a = A \end{cases} \quad (1)$$

where $N_{a,y}$ are the numbers at age a in year y , Z is the total mortality rate ($F + M$), f is the stock-recruit function (for options see Appendix A), and A represents the plus-group. The $\varepsilon_{a,y}$ deviations are akin to survival, which is clearly related to $M_{a,y}$, but they can also be caused by mis-specification of selectivity of the fishery or indices, movement into or out of the population, or misreported catch (Gudmundsson and Gunnlaugsson, 2012; Perretti et al., 2020; Schnute and Richards, 1995; Stock et al., 2021). In this analysis we consider the SCAA model, “Base”, and then examine four possible models that treat the $\varepsilon_{a,y}$ deviations as random effects.

Base is a SCAA model, which estimates recruitment deviations, $\varepsilon_{1,y}$, in each year $y > 1$ as unconstrained fixed effect parameters. The NAA deviations for ages $a > 1$ are fixed at zero so that transitions are deterministic. NAA-1 is most similar to Base, where only recruitment deviations, $\varepsilon_{1,y}$, are estimated. However, in NAA-1 the recruitment deviations are assumed to be independent and identically distributed (IID) random effects,

$$\varepsilon_{1,y} \sim \mathcal{N}\left(-\frac{\sigma_R^2}{2}, \sigma_R^2\right),$$

with variance σ_R^2 as an estimated parameter. The $-\frac{\sigma_R^2}{2}$ bias correction term is included by default so that the expected recruitment, $E(N_{1,y} = R_y)$ equals the expected R_y from the stock-recruit function (Methot and Taylor, 2011; Thorson, 2019). This bias correction adjustment can also be turned off.

NAA-2 is the same as NAA-1, except that the recruitment deviations are stationary AR(1) with autocorrelation parameter $-1 < \rho_{year} < 1$:

$$\varepsilon_{1,y+1} \sim \mathcal{N}\left(\rho_{year}\varepsilon_{1,y} - \frac{\sigma_R^2}{2(1-\rho_{year}^2)}, \sigma_R^2\right)$$

NAA-3 is the “full state-space” model from Nielsen and Berg (2014) and Miller et al. (2016), where all numbers at age are independent random effects and:

$$\varepsilon_{a,y} \sim \begin{cases} \mathcal{N}\left(-\frac{\sigma_R^2}{2}, \sigma_R^2\right), & \text{if } a = 1 \\ \mathcal{N}\left(-\frac{\sigma_a^2}{2}, \sigma_a^2\right), & \text{if } a > 1 \end{cases} \quad (2)$$

where σ_a^2 for all ages $a > 1$ are assumed to be the same but different from age $a = 1$, i.e. recruitment. This assumption is sensible because variability of deviations between expected and realized recruitment are typically larger than deviations from expected abundance at older ages.

NAA-4 treats the numbers at all ages as random effects, as in NAA-3, but the *NAA* deviations, $\varepsilon_{a,y}$, have a 2D stationary AR(1) structure as in Stock et al. (2021):

$$\mathbf{E} \sim \mathcal{MVN}(0, \Sigma)$$

where $\mathbf{E} = (\varepsilon_{1,1}, \dots, \varepsilon_{1,Y-1}, \varepsilon_{2,1}, \dots, \varepsilon_{2,Y-1}, \dots, \varepsilon_{A,1}, \dots, \varepsilon_{A,Y-1})'$ is a vector of all *NAA* deviations, Y is the total number of observation and prediction years, and Σ is the covariance matrix of \mathbf{E} defined by:

$$\text{Cov}(\varepsilon_{a,y}, \varepsilon_{\tilde{a},\tilde{y}}) = \frac{\sigma_a \sigma_{\tilde{a}} \rho_{age}^{|a-\tilde{a}|} \rho_{year}^{|y-\tilde{y}|}}{(1-\rho_{age}^2)(1-\rho_{year}^2)}$$

and $-1 < \rho_{age} < 1$ and $-1 < \rho_{year} < 1$ are the AR(1) coefficients in age and year, respectively. As in NAA-3, σ_a^2 for all ages $a > 1$ are assumed to be the same but different from age $a = 1$, σ_R^2 . The bias correction term for age $a > 1$ in NAA-4 is $-\frac{\sigma_a^2}{2(1-\rho_{year}^2)(1-\rho_{age}^2)}$.

2.1.1.2 Natural mortality (M)

For natural mortality, there are mean parameters for each age, μ_{M_a} , each of which may be estimated freely or fixed at the initial values. The μ_{M_a} may also be estimated in sets of ages, e.g. estimate one mean M shared across ages 3-5, $\mu_{M_3} = \mu_{M_4} = \mu_{M_5}$. There is also an option for M to be specified as a function of weight-at-age, $M_{a,y} = \mu_M W_{a,y}^b$, as in Lorenzen (1996) and Miller and Hyun (2018). Regardless of whether μ_{M_a} are fixed or estimated, WHAM can also be configured to estimate deviations in M , $\delta_{a,y}$, as random effects analogous to the *NAA* deviations (Cadigan, 2016; Stock et al., 2021):

$$\begin{aligned} \log(M_{a,y}) &= \mu_{M_a} + \delta_{a,y} \\ \text{Cov}(\delta_{a,y}, \delta_{\tilde{a},\tilde{y}}) &= \frac{\sigma_M^2 \varphi_{age}^{|a-\tilde{a}|} \varphi_{year}^{|y-\tilde{y}|}}{(1-\varphi_{age}^2)(1-\varphi_{year}^2)} \end{aligned} \quad (3)$$

where σ_M^2 , φ_{age} , and φ_{year} are the AR(1) variance and correlation coefficients in age and year, respectively. In this analysis, we demonstrate three alternative M random effects models. For simplicity, all models treat μ_{M_a} as known, as in most of the original assessments. M-1 is identical to the *NAA*-1 model, with no random effects on M ($\delta_{a,y} = 0$ and not estimated). M-2 allows IID M deviations, estimating σ_M^2 but fixing $\varphi_{age} = \varphi_{year} = 0$. M-3 estimates the full 2D AR(1) structure for M deviations.

2.1.1.3 Selectivity (Sel)

As in ASAP and many other SCAA assessment frameworks, WHAM assumes separability in the fishing mortality rate by age and year, e.g. $F_{a,y} = F_y s_a$, where F_y is the “fully selected” fishing mortality rate in year y and s_a is the selectivity at age a . We note that this differs from the approach in SAM (Nielsen and Berg, 2014), where the $F_{a,y}$ are estimated directly as multivariate random effects without the separability assumption. Three parametric forms are available (logistic, double-logistic, and decreasing-logistic), as well as a non-parametric option to estimate each s_a individually (“age-specific”). WHAM estimates selectivity parameters on the logit scale to avoid boundary problems during estimation. To allow for abrupt temporal changes in selectivity as in ASAP, WHAM can estimate selectivity in user-specified time blocks. WHAM can also estimate smooth changes in selectivity using autoregressive random effects as described below.

WHAM estimates annual full F_y and mean selectivity parameters as fixed effects. Deviations in selectivity parameters can be estimated as random effects, $\zeta_{p,y}$, with autocorrelation by parameter (p), year (y), both, or neither. This is done similarly to Xu et al. (2019), except that the deviations are placed on the parameters instead of the mean $s_{a,y}$ in order to guarantee that $0 < s_{a,y} < 1$. For example, logistic selectivity with two parameters a_{50} and k is estimated as:

$$\begin{aligned}
s_{a,y} &= \frac{1}{1+e^{-(a-a_{50y})/k_y}} \\
a_{50y} &= l_{a_{50}} + \frac{u_{a_{50}} - l_{a_{50}}}{1+e^{-(\nu_1+\zeta_{1,y})}} \\
k_y &= l_k + \frac{u_k - l_k}{1+e^{-(\nu_2+\zeta_{2,y})}} \\
\text{Cov}(\zeta_{1,y}, \zeta_{2,\tilde{y}}) &= \frac{\sigma_s^2 \phi_{par} \phi_{year}^{|y-\tilde{y}|}}{(1-\phi_{par}^2)(1-\phi_{year}^2)}
\end{aligned} \tag{4}$$

where ν_1 is the logit-scale mean a_{50} parameter with lower and upper bounds $l_{a_{50}}$ and $u_{a_{50}}$, ν_2 is the logit-scale mean k parameter with lower and upper bounds l_k and u_k , σ_s^2 is the AR(1) variance, and ϕ_{par} , and ϕ_{year} are the AR(1) correlation coefficients by parameter and year.

Below, we demonstrate three models with random effect deviations on logistic selectivity, akin to those for M . Sel-1 treats all numbers at age as independent random effects (i.e. NAA-3) but with no random effects on s ($\zeta_{p,y} = 0$ and not estimated). Sel-2 allows IID s random effects, estimating σ_s^2 but fixing $\phi_{par} = \phi_{year} = 0$. Sel-3 estimates the full 2D AR(1) structure for s random effects.

2.1.1.4 Environmental covariates (*Ecov*)

WHAM models environmental covariate data using state-space models with process and observation components. The true, unobserved values (or “latent states”, X_y) are then linked to the population dynamics equations with user-specified lag, ψ . For example, recruitment in year y may be influenced by X in the previous year, X_{y-1} ($\psi = 1$), while natural mortality in year y may be influenced by X_y ($\psi = 0$). Multiple environmental covariates may be included, but currently only as independent processes. The environmental and population model years do not need to match, and missing years are allowed. In particular, including environmental data in the projection period can be useful.

2.1.1.4.1 *Ecov* process model

There are currently two options in WHAM for the *Ecov* process model: a normal random walk and AR(1). We model the random walk as in Miller et al. (2016):

$$X_{y+1}|X_y \sim \mathcal{N}(X_y, \sigma_X^2)$$

where σ_X^2 is the process variance and X_1 is estimated as a fixed effect parameter. One disadvantage of the random walk is that it is nonstationary. In short-term projections, \hat{X}_y will be equal to the last estimate with an observation and the uncertainty of \hat{X}_y will increase over time. If \hat{X}_y influences the dynamics of the

217 population and reference points, this leads to increasing uncertainty in projections of the stock and status
 218 over time as well (Miller et al., 2016). For this reason, we generally prefer to model X_y as a stationary AR(1)
 219 process as in Miller et al. (2018):

$$\begin{aligned} X_1 &\sim \mathcal{N}\left(\mu_X, \frac{\sigma_X^2}{1-\phi_X^2}\right) \\ X_y &\sim \mathcal{N}\left(\mu_X(1-\phi_X) + \phi_X X_{y-1}, \sigma_X^2\right) \end{aligned} \quad (5)$$

220 where μ_X , σ_X^2 , and $|\phi_X| < 1$ are the marginal mean, conditional variance, and autocorrelation parameters.
 221 In addition to stationarity, another important difference between the random walk and AR(1) in short-term
 222 population projections is that the AR(1) will gradually revert to the mean over time, unless environmental
 223 covariate observations are included in the projection period.

224 **2.1.1.4.2 *Ecov* observation model**

225 The environmental covariate observations, x_y , are assumed to be normally distributed with mean X_y and
 226 variance $\sigma_{x_y}^2$:

$$x_y | X_y \sim \mathcal{N}\left(X_y, \sigma_{x_y}^2\right)$$

227 The observation variance in each year, $\sigma_{x_y}^2$, can be treated as known with year-specific values (as in Miller
 228 et al., 2016) or one overall value shared among years. They can also be estimated as parameters, likewise
 229 either as yearly values or one overall value. If yearly $\sigma_{x_y}^2$ estimates are desired, WHAM estimates the
 230 hyperparameters μ_{σ_x} and σ_{σ_x} as fixed effects and treats the $\sigma_{x_y}^2$ as random effects:

$$\sigma_{x_y}^2 \sim \mathcal{N}\left(\mu_{\sigma_x}, \sigma_{\sigma_x}^2\right)$$

231 **2.1.1.4.3 Link to population**

232 WHAM currently provides options to link the modeled environmental covariate, X_y , to the population
 233 dynamics via recruitment or natural mortality. It is also sometimes useful to fit the *Ecov* model without a
 234 link to the population dynamics so that models with and without environmental effects have the same data
 235 in the likelihood and can be compared via AIC.

236 In the case of recruitment, the options follow the framework laid out by Fry (1971) and Iles and Beverton
 237 (1998): “controlling” (density-independent mortality), “limiting” (carrying capacity effect, e.g. X_y determines

the amount of suitable habitat), “lethal” (threshold effect, i.e. R_y goes to 0 at some X_y value), “masking” (X_y decreases $dR/dSSB$, as expected if X_y affects metabolism or growth), and “directive” (e.g. behavioral). Of these, WHAM currently allows controlling, limiting, or masking effects in the Beverton-Holt stock-recruit function, and controlling or masking effects in the Ricker function (see Appendix A for all equations).

For natural mortality, environmental effects are placed on μ_M , shared across ages by default. This is the simplest assumption; linking an environmental covariate to M affects M at all ages identically, i.e. $M_{a,y} = \mu_{M_a} e^{\beta X_y - \psi}$. This could be modified in the future to estimate different effects on different ages by changing β from a scalar to a vector of β_a .

Regardless of where the environment-population link is, the effect can be either linear or polynomial. Nonlinear effects of environmental covariates are common in ecology, and quadratic effects are to be expected in cases where intermediate values are optimal (Agostini et al., 2008; Brett, 1971). WHAM includes a function to calculate orthogonal polynomials in TMB, akin to the `poly()` function in R. Orthogonal polynomials remove correlation between covariates, e.g. X and X^2 , which improves estimability and allows for straightforward evaluation of the significance of adding X^2 to a model that includes X .

In this analysis, we compare five models with limiting effects on Beverton-Holt recruitment:

$$\hat{R}_{y+1} = \frac{\alpha SSB_y}{1 + e^{\beta_0 + \beta_1 X_y + \beta_2 X_y^2} SSB_y} \quad (6)$$

where SSB_y is spawning stock biomass in year y , α and β_0 are the standard parameters of the Beverton-Holt function, and β_1 and β_2 are polynomial effect terms that modify β_0 based on the value of the estimated environmental covariate, X_y . Eqn. 6 is one of several stock-recruit functions with environmental links implemented in WHAM (see Appendix A for complete list).

Ecov-1 treats X_y as a random walk ($\phi_X = 1$) but does not include an effect on recruitment ($\beta_1 = \beta_2 = 0$). We include Ecov-1 in order to compare AIC of the original model to those with environmental effects on recruitment. Ecov-2 and Ecov-3 also treat X_y as a random walk, but Ecov-2 estimates β_1 and Ecov-3 estimates both β_1 and β_2 . Ecov-4 and Ecov-5 estimate X_y as an AR(1) process instead of a random walk (estimate ϕ_X), and Ecov-4 estimates β_1 and Ecov-5 estimates both β_1 and β_2 .

2.1.2 Population observation model

Like ASAP, there are likelihood components for annual aggregate catch and relative abundance observations for each fleet and index, and annual age compositions for each fleet and index.

2.1.2.1 Aggregate catch and indices

The predicted catch at age for fleet i , $\hat{C}_{a,y,i}$, is a function of $N_{a,y}$, $M_{a,y}$, $F_{y,i}$, and $s_{a,y,i}$:

$$\hat{C}_{a,y,i} = N_{a,y} (1 - e^{-Z_{a,y}}) \frac{F_{a,y,i}}{Z_{a,y}}. \quad (7)$$

The log-aggregate catch $\hat{C}_{y,i} = \sum_a \hat{C}_{a,y,i} W_{a,y,i}$ observation, where $W_{a,y,i}$ is the empirical weight at age, is assumed to have a normal distribution

$$\log(C_{y,i}) \sim \mathcal{N} \left(\log(\hat{C}_{y,i}) - \frac{\sigma_{\hat{C}_{y,i}}^2}{2}, \sigma_{\hat{C}_{y,i}}^2 \right) \quad (8)$$

where the standard deviation

$$\sigma_{C_{y,i}} = e^{\eta_i} \sigma_{\hat{C}_{y,i}}$$

is a function of an input standard deviation $\sigma_{\hat{C}_{y,i}}$ and a fleet-specific parameter η_i that is fixed at 0 by default, but may be estimated. The bias correction term, $-\frac{\sigma_{\hat{C}_{y,i}}^2}{2}$, is included by default based on Aldrin et al. (2020) but can be turned off.

Observations of aggregate indices of abundance are handled identically to the aggregate catch as in Eqn. 8 except that for index i the predicted index at age is

$$\hat{I}_{a,y,i} = q_i s_{a,y,i} N_{a,y} W_{a,y,i} e^{-Z_{a,y} f_{y,i}}. \quad (9)$$

where q_i is the catchability and $f_{y,i}$ is the fraction of the annual time step elapsed when the index is observed. There are options for indices to be in terms of abundance (numbers) or biomass and $W_{a,y,i} = 1$ for the former.

2.1.2.2 Catch and index age composition

WHAM includes several options for the catch and index age compositions, including the multinomial (the default to match ASAP), Dirichlet, Dirichlet-multinomial (Thorson, 2019), and logistic normal (Aitchison and Shen, 1980; Schnute and Haigh, 2007) distributions (Appendix B). Although the multinomial has been commonly used in stock assessment, it is generally inferior to more flexible options that are self-weighting and allow for correlations, such as the Dirichlet and logistic normal (Francis, 2014). We therefore assumed a logistic normal distribution for age composition observations in all of the applications and simulation studies here.

2.1.3 Projections

The default settings for short-term projections follow common practice for stock assessments in the U.S. Northeast: the population is projected three years using the average selectivity, maturity, weight, and natural mortality at age from the last five model years to calculate reference points (NEFSC, 2020a). WHAM implements similar options as ASAP for specifying F_y in the projection years: terminal year F_y , average F over specified years, $F_{X\%}$ (F at $X\%$ SPR, where X is specified and 40 by default), user-specified F_y , or F derived from user-specified catch. For all options except user-specified F_y , the uncertainty in projected F is propagated into the uncertainty of projected population attributes. For models with random effects on NAA , M , or $Ecov$, the default is to continue the stochastic process into the projection years. WHAM does not currently do this for selectivity because, like F , it is a function of management. Instead, selectivity is taken as the average of recent model years.

The decision whether or not to continue autocorrelated processes into the projection years is important because it can substantially affect short-term SSB forecasts (Stock et al., 2021). This occurs because any non-zero NAA or M deviations near the end of the assessment are propagated into the projection period.

If the $Ecov$ data extend beyond the population model years, WHAM will fit the $Ecov$ model to all available data and use the estimated X_y in projections. This may often be the case because lags can exist in both the physical-biological mechanism and the assessment process. As an example, a model with an $Ecov$ effect on recruitment may link physical oceanographic conditions, e.g. surface or bottom temperature, in year t to recruitment in year $t + 1$, and the assessment conducted in year t may only use population data through year $t - 1$. In this case, 3-year population projections only need the $Ecov$ model to be projected one year. While the default handling of $Ecov$ projections is to continue the stochastic process, WHAM includes options to use terminal year x_y , x_y averaged over specified years, or specified x_y . The option to specify x_y allows users to investigate how alternative climate projections may affect the stock.

2.2 Fits to original datasets

We fit the models described above to data from five stocks with a broad range of life history, status, and model dimension (number of ages and years): Southern New England-Mid Atlantic (SNEMA) yellowtail flounder, butterfish, North Sea cod, Icelandic herring, and Georges Bank (GB) haddock (Tables 1 and 2). We assumed the same form of logistic normal for SNEMA yellowtail flounder age composition observations as Miller et al. (2016) where unobserved ages are pooled with adjacent ages, but for other stocks unobserved ages are treated as missing. The variance parameters associated with the logistic normal distributions were

estimated for all stocks.

Although WHAM can treat multiple processes as random effects simultaneously (e.g. *NAA* and *M*; Stock et al., 2021), here we only fit alternate models for one process at a time for simplicity. We fit the *NAA* random effects models to all five stocks since these represent core WHAM functionality. We chose to highlight one stock each for the *M*, *Sel*, and *Ecov* processes because all models did not converge for all stocks and processes: SNEMA yellowtail flounder (*M* and *Ecov*) and GB haddock (*Sel*). We fixed μ_{M_a} at the values used in the original assessments, except for butterfish where it estimated in the assessment. Aside from the GB haddock *Sel* models, we used the same selectivity parameterization and time blocks as in the original assessments. We used the NAA-3 model (all *NAA* deviations are random effects) in the *Sel* and *Ecov* demonstrations, but the NAA-1 model (only recruitment deviations are random effects) for the *M* demonstrations, because deviations in the *NAA* transitions can be caused by unmodeled deviations in *M*. For the SNEMA yellowtail flounder *Ecov* models, we used the Cold Pool Index (CPI) as calculated in Miller et al. (2016). We updated the CPI through 2018, except that the 2017 observation was missing because the NEFSC fall bottom trawl survey was not completed in the SNEMA region. Within each process, we compared the performance of the different random effects models using AIC.

We fit all models using the open-source statistical software R (R Core Team, 2020) and TMB (Kristensen et al., 2016), as implemented in the R package WHAM (Miller and Stock, 2020). Documentation and tutorials for how to specify additional random effect structures in WHAM are available at <https://timjmiller.github.io/wham/>. Code and data files to run the analysis presented here are available at <https://github.com/brianstock-NOAA/wham-sim>.

2.3 Simulation tests

After fitting each model to the original datasets, we used the simulation feature of TMB to conduct self- and cross-tests (*sensu* Deroba et al., 2015). For each model, we generated 100 sets of new data and random effects, keeping the fixed effect parameters constant at values estimated in original fits. We then re-fit all models to datasets simulated under each as an operating model. We calculated the relative error in parameters constraining random effects (Table 1) and quantities of interest, such as spawning stock biomass (SSB), F , $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R . We calculated relative error as $\frac{\hat{\theta}_i}{\theta_i} - 1$, where θ_i is the true value for simulated dataset i and $\hat{\theta}_i$ is the value estimated from fitting the model to the simulated data. To estimate bias of each estimation model for a given operating model, we calculated 95% confidence intervals of the median relative error using the binomial distribution (Thompson, 1936). To summarize the bias across simulations and years for each

model, we calculated the median quantity across years and took the mean of the medians across simulations. Finally, for each operating model we calculated the proportion of simulations in which each estimation model converged and had the lowest AIC.

3 Results

3.1 Original datasets

The 2D AR(1) covariance structure for random effects on NAA , M , and selectivity performed well across stocks and processes, evidenced by lower AIC than the IID random effects models (Fig. 1). This AIC difference was larger for selectivity than NAA or M , but the differences for NAA and M were also non-trivial, ranging from 11.1–53.2. Incorporating random effects on NAA , M or selectivity did not have consistently positive or negative effects on key assessment model output (SSB, F , or recruitment); these effects differed by stock and process (Figs. S1–S10). One pair of 2D AR(1) covariance parameters, σ and ρ_{year} , was consistently estimated with negative correlation across all stocks and processes, -0.43 on average. All other pairwise combinations of the estimated 2D AR(1) parameters had correlation values less than 0.2 in magnitude.

3.1.1 Numbers-at-age (NAA)

Treating numbers at all ages as random effects was strongly supported by AIC, compared to treating only recruitment deviations as random effects (Fig. 1). Including autoregressive NAA random effects was also generally supported by AIC, either the AR(1) when only recruitment deviations were random effects or the 2D AR(1) when all ages were random effects. Across the five stocks, models that treated all NAA as random effects (i.e., full state-space models) also had reduced retrospective pattern and increased uncertainty in SSB, F , and recruitment (Figs. 2-3, Table S1). Including 2D AR(1) autocorrelation on NAA deviations further reduced retrospective patterns and increased uncertainty on average.

The estimated AR(1) and IID random effects were similar, with the AR(1) random effects slightly smoothed compared to the IID random effects when the estimated correlations were positive (e.g. $\rho_{year} = 0.73$, 95% CI: 0.58-0.87, for Georges Bank haddock in Fig. 4).

3.1.2 Natural mortality (M)

As for the *NAA* models, including random effect deviations on M was supported by AIC (Fig. 1). Although the 2D AR(1) M model did not converge for North Sea cod, it had the lowest AIC for butterfish and SNEMA yellowtail flounder. The M-3 model with 2D AR(1) autocorrelation estimated similar M deviations as the M-2 model, with smoothing when estimated correlations were positive (e.g. $\varphi_{year} = 0.63$, 95% CI: 0.31-0.94, and $\varphi_{age} = 0.40$, 95% CI: 0.09-0.70, for SNEMA yellowtail flounder in Fig. 5). Both M-2 and M-3 estimated elevated M for fish aged 5-6+ and lower M for ages 2-3 for several years in the middle of the time-series, and then higher M for ages 1-3 at the end of the time-series.

3.1.3 Selectivity (Sel)

For Georges Bank haddock, including 2D AR(1) random effect deviations on selectivity was strongly supported by AIC (Fig. 1). Sel-3 estimated similar patterns in selectivity compared to Sel-2, except with smoother variations by age and year (Fig. 6). Compared to the model with constant selectivity, Sel-2 and Sel-3 estimated lower s for age 3 in recent years and higher s for age 3 before 1990. They also estimated higher s for age 2 before 1990, especially from 1973 to 1976.

3.1.4 Environmental covariate effect on recruitment ($Ecov$)

As in previous analyses (Miller et al., 2016; Xu et al., 2018), including an effect of the CPI on recruitment for SNEMA yellowtail flounder was clearly supported by AIC (ΔAIC of 20.3-33.0 between Ecov-1 and Ecov-2 through Ecov-5, Fig. 1). The AR(1)-linear model (Ecov-4) had the lowest AIC—fitting the CPI using an AR(1) model was preferred over the random walk (ΔAIC of 12.7 between Ecov-2 and Ecov-4), and the quadratic term, β_2 , was deemed unnecessary (ΔAIC of 1.3 between Ecov-5 and Ecov-4). As expected, the AR(1) process model estimated higher uncertainty in years with higher observation error (e.g. 1982–1992) and missing observations (2017, Fig. 7A). The CPI negatively influenced recruitment, i.e. \hat{R} was higher following years with lower CPI (lower fall bottom temperature, Fig. 7C). Including the CPI-recruitment link changed the estimates of R_y by up to 30%, but in most years the relative difference in R_y was less than 10% (Figs. 7B–C and S3). The relative differences in SSB and F were small, less than 4% in most years, compared to the influence from adding random effects on all *NAA* or M (Figs. S1–S3).

3.2 Simulation tests

Several findings were consistent across processes and stocks. When the estimation and operating model were consistent, bias of SSB, F , $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R was generally less than 2% and not significant based on confidence intervals (Figs. 8–10 and S11–S16). Bias was also generally less than 2% when more complex models were fitted to less complicated operating model simulations (Fig. S17). In contrast, the bias was often greater than 5% when the estimation model was less complex than the operating model, especially when models without random effects were fit to data simulated from models with random effects (Fig. S17). Bias in SSB and F were always opposite, i.e. when SSB was biased high, F was biased low, and vice versa. Predicted catch was never biased. Bias of the variance and correlation parameters controlling random effects was generally not significant or negative, as expected using maximum likelihood estimation (Figs. S18–S21). Restricted maximum likelihood (REML) should be used if more accurate estimation of these parameters is a priority.

In cross-tests, the percentage of simulations in which AIC selected the correct model varied between 62–99% by process, stock, and operating model (Fig. 11). Estimation models more complex than the operating model were more likely to be chosen for *NAA* models than for *M* or selectivity.

3.2.1 Numbers-at-age

Across the five stocks, all *NAA* models estimated SSB, F , $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R with minimal bias in self-tests ($< 2\%$, Fig. S17). An exception was the estimation of F for Icelandic herring, particularly for NAA-1 and NAA-2 (median relative error in F with 95% CI for NAA-1: -0.060 (-0.073, -0.047), NAA-2: -0.060 (-0.073, -0.047), NAA-3: 0.015 (-0.004, 0.034), and NAA-4: 0.049 (0.017, 0.080); Figs. S14 and S22–S24). The convergence rate for most models and stocks was above 95%, with the same exception of NAA-1 and NAA-2 for Icelandic herring (Fig. S25). In cross-tests, NAA-1 and NAA-2 exhibited non-trivial bias when estimating data simulated from models that treated numbers at all ages as random effects. The degree of bias varied by quantity and stock between -25% and 25% (Figs. 8 and S11–S14). The more complex *NAA* models estimated all quantities without bias regardless of operating model.

For four of the five stocks, σ_R^2 was estimated without bias in self-tests (Fig. S18). The exception was butterflyfish, for which σ_R^2 was negatively biased in NAA-1, NAA-2, and NAA-4 (but not NAA-3). In contrast, both NAA-3 and NAA-4 estimated σ_a^2 with negative bias for all five stocks. There was no consistent pattern in bias for the correlation parameters ρ_{age} and ρ_{year} .

3.2.2 Natural mortality

SSB, F , $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R were estimated without significant bias in self-tests, although less complex models exhibited bias when fit to data simulated from more complex models (Figs. 9 and S15–S17).

As for the *NAA* models, σ_R^2 was estimated without bias in the *M* model self-tests (Fig. S18). σ_M^2 was biased low, φ_{year} had no bias, and the direction of bias in φ_{age} was inconsistent.

3.2.3 Selectivity

Models with IID or 2D AR(1) s random effects estimated SSB, F , $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R with little to no bias across operating models (Fig. 10). The model without s random effects, Sel-1, showed substantial bias when fit to data simulated from Sel-2 or Sel-3. For Georges Bank haddock, the variance and correlation parameters σ_a^2 , σ_{Sel}^2 , ϕ_{year} , and ϕ_{par} were all estimated with slight negative bias in self-tests of models with s random effects (Fig. S20).

3.2.4 Ecov-Recruitment

The random walk CPI models estimated σ_X^2 with less bias than the AR(1) CPI models (Fig. S21). β_0 was biased high in all models, although this was not significant for the model with lowest AIC (Ecov-4, AR(1)-linear). All models estimated the parameters ϕ_X , α , β_1 , and β_2 without significant bias.

4 Discussion

4.1 Overview

Our results suggest that the WHAM package can be a useful tool for stock assessment when environmental effects on recruitment, or stochastic changes in the numbers at age transitions, selectivity, or natural mortality are of interest. The simulation tests showed negligible or no bias in estimation of important assessment outputs (SSB, F , stock and harvest status) when the operating and estimation models matched. The less complex models, without random effects or autoregressive structure, exhibited some bias in cross-tests, while the more complex models did not. In these cases, bias in SSB and F were opposite such that predicted catch was unbiased. WHAM models with independent or 2D AR(1) random effect deviations performed well across stocks and processes, which suggests that they warrant consideration in future stock-specific studies. The 2D

AR(1) covariance structure performed particularly well, generally having the lowest AIC and retrospective patterns (Figs. 1-2, Table S1) and outperforming the model with independent random effects in cross-tests (Figs. 8-11 and S11–S15).

4.2 Limitations

For simplicity, we only demonstrated alternative models for a single random effect process in each of our comparisons. WHAM can, however, treat multiple processes as random effects simultaneously. For SNEMA yellowtail flounder, Stock et al. (2021) found that models with process errors in both M and NAA converged and performed best by AIC and Mohn’s ρ . Models with 2D autocorrelation in either process were identifiable and estimated similar F and SSB trends. Preliminary work on other stocks in the region have also shown this is possible. In general, which combinations of process errors are expected to work jointly is difficult to say without testing different combinations. The answer is very likely to be stock-specific as well, depending on the information content in the data, number of ages and years, life history, and exploitation history.

Due to the number of models and stocks we tested, we did not thoroughly examine diagnostics, such as residual patterns, for each model fit that would be checked in a full assessment. It is possible that doing so might lead to the rejection of some model configurations for some stocks.

Further improvements could be made to the handling of age compositions in WHAM. The current Dirichlet, Dirichlet-multinomial, and logistic normal options are more flexible than the multinomial distribution used in ASAP (Francis, 2014). Yet, extensions to the logistic normal are possible; most notably, additional correlation structures as in Francis (2014) and Martell and Lima (2015).

Finally, some of the 2D AR(1) covariance parameters were estimated with negative bias, particularly σ_a (Figs. S18-S21). This is the expected direction of bias using maximum likelihood, and using REML instead would likely reduce it (Miller et al., 2018). This is relatively easy to do in WHAM, or more generally in TMB, by treating all parameters except the 2D AR(1) covariance parameters as random effects. This directs TMB to use the Laplace approximation to integrate over the other fixed effect parameters (treated as random effects) with flat priors (Harville, 1974). We did not do this because it requires constructing and fitting the model in TMB a second time for each simulation.

4.3 Relationships to other existing assessment model frameworks

WHAM assumes separability in $F_{a,y}$, i.e. $F_{a,y} = F_y s_a$, and estimates annual full F_y as fixed effects and s_a as constant or time-varying random effects with various autoregressive assumptions possible. Most other assessment frameworks in the U.S, e.g. ASAP, Stock Synthesis (Methot and Wetzel, 2013), an Assessment Model for Alaska (AMAK; Anon., 2015), and the Beaufort Assessment Model (BAM; Williams and Shertzer, 2015), make this same separability assumption which is useful for specifying a fully-selected F in projections to calculate reference points or generating catch advice. Estimating selectivity in time blocks is common practice in these frameworks. In contrast, SAM estimates $F_{a,y}$ directly as a multivariate random walk process (Nielsen and Berg, 2014). The autoregressive models for selectivity parameters under certain configurations of WHAM should allow for similar F at age patterns as in SAM. Stock Synthesis allows 2D AR(1) random effects on $s_{a,y}$ instead of the parameters, and then the variance parameter is estimated by an iterative tuning algorithm (Xu et al., 2019). It is unclear whether estimating time-varying selectivity as random effects produces better results than assuming time blocks, or if there are advantages to using any of the three approaches for estimating time-varying selectivity used by WHAM, SAM, or Stock Synthesis.

Likewise, WHAM and the other U.S.-based assessment models treat catch, index, and composition observations differently than SAM. The U.S.-based assessment models treat aggregate and composition observations separately for fisheries and indices whereas SAM treats observations of catch and indices at age, $C_{a,y}$ and $I_{a,y}$, as multivariate log-normal. Separate observation models for aggregate and composition observations is natural when sampling for total catch differs from that for length and age composition.

Like SAM, WHAM can estimate interannual transitions in NAA as a random walk processes, but WHAM can also be configured to treat these deviations as stationary autoregressive processes. Alternatively (or simultaneously, Stock et al., 2021) WHAM can estimate deviations in natural mortality as autoregressive processes like NCAM (Cadigan, 2016). Uniquely, WHAM can model multiple environmental covariate time series as state-space processes and include their effects, possibly nonlinearly, in various ways on recruitment or natural mortality. Although most applications thusfar investigate effects of physical processes on demographic parameters, indices of predation might also be considered (Marshall et al., 2019).

4.4 Future use of WHAM

The productivity of several groundfish stocks in the U.S. Northeast has declined in recent decades, and conducting assessments in WHAM with time-varying processes via random effects or environment-productivity links could account for these trends and potentially reduce retrospective bias (Perretti et al., 2017; Stock et al.,

2021; Tableau et al., 2018). WHAM can be configured to fit SCAA models very similar to ASAP and therefore bridge between the two frameworks. Adding random effects or environmental covariates to an assessment model is a large structural change, and simulation self- and cross-tests such as we have demonstrated here should be conducted through the research track process (Lynch et al., 2018). If comparisons against status quo SCAA models prove favorable, WHAM could transition to being used in operational assessments. However, because the details of how to include time-varying processes, as well as the effect on the assessment output, will vary by stock, this evaluation may need to be conducted on a stock-by-stock basis.

Random effects allow for changing productivity and, if they are considered as an autoregressive process, can propagate the effect of these changes on assessment output in short-term projections (Stock et al., 2021). However, the AR(1) process demonstrated here trends to the mean in projections and will not predict values beyond extremes in observed time-series. This is an issue worth examining because many marine ecosystems are changing to such extent that recent conditions are time-series extremes, and conditions in the near future may continue to expand the range of observations (e.g. sea surface temperatures over the U.S. Northeast Shelf in the last decade; Chen et al., 2020). More complex nonstationary time-series models such as autoregressive integrated moving average (ARIMA) that can forecast beyond observed values could be implemented. Nonlinear (e.g. splines) or double linear integration (Di Lorenzo and Ohman, 2013) techniques could also be worth pursuing, as well as time-delay embedding methods (Munch et al., 2018, 2017), which not only allow for nonstationary dynamics but also do not require a specified functional form. The best approach to making stochastic projections of productivity responses to environmental conditions that are rapidly changing and at time-series extremes merits further research.

In addition to the empirical (i.e. random effects) approach, it is worth considering the mechanistic approach (i.e. explicitly modeled links to environmental covariates) for a given stock whose recruitment or M is suspected to have shifted due to a clear, plausible hypothesized external influence. Several factors may result in a higher likelihood that the mechanistic approach is useful in an assessment: a history of overfishing (Free et al., 2019; but high F can also swamp the signal of an environmental influence, Haltuch and Punt, 2011), more rapid environmental change, stocks at the edge of species' range, opportunistic (short-lived) species (*sensu* Winemiller and Rose, 1992), lower trophic level species, longer time series, periodic signals in which more than once cycle has been recorded (e.g. Pacific Decadal Oscillation and sardine), and stronger signals (wider ranges of observed stock status and environmental conditions) (Free et al., 2019; Haltuch et al., 2019; Haltuch and Punt, 2011; Marshall et al., 2019). Although developing mature mechanistic hypotheses will take significant effort and collaboration with ecologists and oceanographers, these rules of thumb can guide researchers in prioritizing which stocks to investigate environment-productivity relationships. Our

ability to forecast and understand oceanographic and climate variables continues to improve and increasingly will present opportunities for including well-founded mechanistic environmental effects in stock assessments (Tommasi et al., 2017).

The perception of precision in model output such as projected population biomass or catch advice is affected by whether productivity components are treated as either constant or stochastic processes (Figs. S1–S10), much like whether certain parameters are either fixed or estimated in traditional assessment models. Modeling frameworks such as WHAM allow the performance of these alternative models to be compared and when allowing temporal variation in productivity components is justified, the greater uncertainty in the assessment output is more realistic. Moreover, model uncertainty could be included in our preception of output uncertainty by fitting ensembles of WHAM models that represent alternative states of nature (Anderson et al., 2017; Möllmann et al., 2014). Finally, the simulation capabilities of WHAM can also be useful in situations where variation in productivity is hypothesized, but information to estimate this variation is unavailable. WHAM can be configured as an operating model to simulate plausible environmentally-driven changes in recruitment or M or plausible stochastic variation in order to estimate the sensitivity of status quo models.

4.5 Conclusion

A major present-day challenge in fisheries is to assess and manage stocks in a changing environment. We foresee environment-linked stock assessments becoming more feasible and realistic as fisheries and oceanographic time-series lengthen and our ecological understanding deepens. We have developed WHAM with this in mind. Finally, we note that the development of TMB has been a critical advance for fisheries assessment modeling frameworks such as WHAM, allowing us to rapidly fit models that treat population and environmental processes as time-varying random effects in a state-space framework.

Acknowledgements

We appreciate the efforts of the organizers of the special issue and CAPAM workshop on next-generation stock assessment models held on November 4-8, 2019 in Wellington, New Zealand. We also thank Charles Perretti and two anonymous reviewers for helpful suggestions on the manuscript. This research was performed while BCS held an NRC Research Associateship award at the NEFSC under the Northeast Groundfish and Climate Initiative. We thank Chris Melrose for updating the Cold Pool Index with data through 2018, using code originally written by Jon Hare.

References

- Aeberhard, W.H., Mills Flemming, J., Nielsen, A., 2018. Review of State-Space Models for Fisheries Science. *Annu. Rev. Stat. Appl.* 5, 215–235. <https://doi.org/10.1146/annurev-statistics-031017-100427>
- Agostini, V., Hendrix, A., Hollowed, A., Wilson, C., Pierce, S., Francis, R., 2008. Climate-ocean variability and Pacific hake: A geostatistical modeling approach. *J. Mar. Sys.* 71, 237–248. <https://doi.org/10.1016/j.jmarsys.2007.01.010>
- Aitchison, J., Shen, S.M., 1980. Logistic-Normal Distributions: Some Properties and Uses. *Biometrika* 67, 261–272.
- Aldrin, M., Tvete, I., Aanes, S., Subbey, S., 2020. The specification of the data model part in the SAM model matters. *Fish. Res.* 229, 105585. <https://doi.org/10.1016/j.fishres.2020.105585>
- Anderson, S.C., Cooper, A.B., Jensen, O.P., Minto, C., Thorson, J.T., Walsh, J.C., Afflerbach, J., Dickey-Collas, M., Kleisner, K.M., Longo, C., Osio, G.C., Ovando, D., Mosqueira, I., Rosenberg, A.A., Selig, E.R., 2017. Improving estimates of population status and trend with superensemble models. *Fish Fish.* 18, 732–741. <https://doi.org/10.1111/faf.12200>
- Anon., 2015. Assessment Model for Alaska Description of GUI and Instructions.
- Berg, C.W., Nielsen, A., 2016. Accounting for correlated observations in an age-based state-space stock assessment model. *ICES J. Mar. Sci.* 73, 1788–1797. <https://doi.org/10.1093/icesjms/fsw046>
- Brett, J.R., 1971. Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus Nerka*). *Am. Zool.* 11, 99–113. <https://doi.org/10.1093/icb/11.1.99>
- Brooks, E.N., Legault, C.M., 2016. Retrospective forecasting evaluating performance of stock projections for New England groundfish stocks. *Can. J. Fish. Aquat. Sci.* 73, 935–950. <https://doi.org/10.1139/cjfas-2015-0163>
- Cadigan, N.G., 2016. A state-space stock assessment model for northern cod, including under-reported catches and variable natural mortality rates. *Can. J. Fish. Aquat. Sci.* 73, 296–308. <https://doi.org/10.1139/cjfas-2015-0047>
- Chen, Z., Kwon, Y.-O., Chen, K., Fratantoni, P., Gawarkiewicz, G., Joyce, T.M., 2020. Long-Term SST Variability on the Northwest Atlantic Continental Shelf and Slope. *Geophys. Res. Lett.* 47. <https://doi.org/10.1029/2019GL085455>

De Oliveira, J., Butterworth, D., 2005. Limits to the use of environmental indices to reduce risk and/or increase yield in the South African anchovy fishery. *Afr. J. Mar. Sci.* 27, 191–203. <https://doi.org/10.2989/18142320509504078>

Deroba, J.J., Butterworth, D.S., Methot, R.D., De Oliveira, J.a.A., Fernandez, C., Nielsen, A., Cadrin, S.X., Dickey-Collas, M., Legault, C.M., Ianelli, J., Valero, J.L., Needle, C.L., O'Malley, J.M., Chang, Y.-J., Thompson, G.G., Canales, C., Swain, D.P., Miller, D.C.M., Hintzen, N.T., Bertignac, M., Ibaibarriaga, L., Silva, A., Murta, A., Kell, L.T., de Moor, C.L., Parma, A.M., Dichmont, C.M., Restrepo, V.R., Ye, Y., Jardim, E., Spencer, P.D., Hanselman, D.H., Blaylock, J., Mood, M., Hulson, P.-J.F., 2015. Simulation testing the robustness of stock assessment models to error: Some results from the ICES strategic initiative on stock assessment methods. *ICES J. Mar. Sci.* 72, 19–30. <https://doi.org/10.1093/icesjms/fst237>

Di Lorenzo, E., Ohman, M.D., 2013. A double-integration hypothesis to explain ocean ecosystem response to climate forcing. *Proc. Natl. Acad. Sci. U.S.A.* 110, 2496–2499. <https://doi.org/10.1073/pnas.1218022110>

Francis, R.C., 2014. Replacing the multinomial in stock assessment models: A first step. *Fish. Res.* 151, 70–84. <https://doi.org/10.1016/j.fishres.2013.12.015>

Free, C.M., Thorson, J.T., Pinsky, M.L., Oken, K.L., Wiedenmann, J., Jensen, O.P., 2019. Impacts of historical warming on marine fisheries production. *Science* 363, 979–983. <https://doi.org/10.1126/science.aau1758>

Fry, F., 1971. The effect of environmental factors on the physiology of fish, in: *Fish Physiology*. Elsevier, pp. 1–98. [https://doi.org/10.1016/S1546-5098\(08\)60146-6](https://doi.org/10.1016/S1546-5098(08)60146-6)

Garstang, W., 1900. The Impoverishment of the Sea. A Critical Summary of the Experimental and Statistical Evidence bearing upon the Alleged Depletion of the Trawling Grounds. *J. Mar. Biol. Assoc. U.K.* 6, 1–69. <https://doi.org/10.1017/S0025315400072374>

Gudmundsson, G., 1994. Time series analysis of catch-at-age observations. *Appl. Stat.* 43, 117–126. <https://doi.org/10.2307/2986116>

Gudmundsson, G., Gunnlaugsson, T., 2012. Selection and estimation of sequential catch-at-age models. *Can. J. Fish. Aquat. Sci.* 69, 1760–1772. <https://doi.org/10.1139/f2012-095>

Haltuch, M.A., Brooks, E., Brodziak, J., Devine, J., Johnson, K., Klibansky, N., Nash, R., Payne, M., Shertzer, K., Subbey, S., Wells, B., 2019. Unraveling the recruitment problem: A review of environmentally-informed forecasting and management strategy evaluation. *Fish. Res.* 217, 198–216. <https://doi.org/10.1016/j.fishres.2018.12.016>

Haltuch, M.A., Punt, A.E., 2011. The promises and pitfalls of including decadal-scale climate forcing of

recruitment in groundfish stock assessment. *Can. J. Fish. Aquat. Sci.* 68, 912–926. <https://doi.org/10.1139/f2011-030>

Hare, J.A., Borggaard, D.L., Friedland, K.D., Anderson, J., Burns, P., Chu, K., Clay, P.M., Collins, M.J., Cooper, P., Fratantoni, P.S., Johnson, M.R., Manderson, J.P., Milke, L., Miller, T.J., Orphanides, C.D., Saba, V.S., 2016. Northeast Regional Action Plan - NOAA Fisheries Climate Science Strategy (No. NMFS-NE-239). NOAA Fisheries, Northeast Fisheries Science Center, Woods Hole, MA.

Harville, D.A., 1974. Bayesian inference for variance components using only error contrasts. *Biometrika* 61, 383–385. <https://doi.org/10.1093/biomet/61.2.383>

Hill, K.T., Crone, P.R., Zwolinski, J.P., 2018. Assessment of the Pacific sardine resource in 2018 for U.S. Management in 2018-2019 (No. NOAA Technical Memorandum NMFS-SWFSC-600). US Department of Commerce.

Hjort, J., 1914. Fluctuations in the great fisheries of Northern Europe viewed in the light of biological research. *Rapports et Procès-Verbaux des Réunions du Conseil Permanent International Pour L'Exploration de la Mer* 20, 1–228.

ICES, 2020. Workshop on the review and future of state space stock assessment models in ICES (WKRFSAM). *ICES Sci. Rep.* 2, 23p. <https://doi.org/10.17895/ices.pub.6004>

ICES, 2017a. Report of the North Western Working Group (NWWG). *ICES CM 2017/ACOM:08*, 27 April-4 May 2017, Copenhagen, Denmark.

ICES, 2017b. Report of the working group on assessment of demersal stocks in the North Sea and Skagerrak (2017). *ICES CM 2017/ACOM:21*, 26 April-5 May 2017, ICES HQ.

Iles, T.C., Beverton, R.J.H., 1998. Stock, recruitment and moderating processes in flatfish. *J. Sea Res.*, Proceedings of the Third International Symposium on Flatfish Ecology, Part II 39, 41–55. [https://doi.org/10.1016/S1385-1101\(97\)00022-1](https://doi.org/10.1016/S1385-1101(97)00022-1)

Kristensen, K., Nielsen, A., Berg, C., Skaug, H., Bell, B.M., 2016. TMB: Automatic differentiation and Laplace approximation. *J. Stat. Soft.* 70, 1–21. <https://doi.org/10.18637/jss.v070.i05>

Kumar, R., Cadigan, N.G., Zheng, N., Varkey, D.A., Morgan, M.J., 2020. A state-space spatial survey-based stock assessment (SSURBA) model to inform spatial variation in relative stock trends. *Can. J. Fish. Aquat. Sci.* 77, 21.

Larkin, P., 1996. Concepts and issues in marine ecosystem management. *Rev. Fish Biol. Fisher.* 6, 139–164.

654 <https://doi.org/10.1007/BF00182341>

655 Legault, C.M., Restrepo, V.R., 1998. A Flexible Forward Age-Structured Assessment Program (No. 49).

656 Link, J.S., 2002. What Does Ecosystem-Based Fisheries Management Mean? *Fisheries* 27, 5.

657 Lorenzen, K., 1996. The relationship between body weight and natural mortality in juvenile and adult fish: A
658 comparison of natural ecosystems and aquaculture. *J. Fish Biol.* 49, 627–642. [https://doi.org/10.1111/j.1095-](https://doi.org/10.1111/j.1095-8649.1996.tb00060.x)
659 [8649.1996.tb00060.x](https://doi.org/10.1111/j.1095-8649.1996.tb00060.x)

660 Lynch, P.D., Methot, R.D., Link, J.S. (Eds.), 2018. Implementing a Next Generation Stock Assessment
661 Enterprise. An Update to the NOAA Fisheries Stock Assessment Improvement Plan, in: U.S. Dep. Commer.,
662 NOAA Tech. Memo. NMFS-F/ SPO-183, p. 127. <https://doi.org/10.7755/TMSPO.183>

663 Marshall, K.N., Koehn, L.E., Levin, P.S., Essington, T.E., Jensen, O.P., 2019. Inclusion of ecosystem
664 information in US fish stock assessments suggests progress toward ecosystem-based fisheries management.
665 *ICES J. Mar. Sci.* 76, 1–9. <https://doi.org/10.1093/icesjms/fsy152>

666 Martell, S.J.D., Lima, C.W., 2015. Technical documentation for the integrated statistical catch age model
667 (iSCAM).

668 McClatchie, S., Goericke, R., Auad, G., Hill, K., 2010. Re-assessment of the stockRecruit and tempera-
669 tureRecruit relationships for Pacific sardine (*Sardinops sagax*). *Can. J. Fish. Aquat. Sci.* 67, 1782–1790.
670 <https://doi.org/10.1139/F10-101>

671 Mendelsohn, R., 1988. Some problems in estimating population sizes from catch-at-age data. *Fish. Bull.* 86,
672 617–630.

673 Methot, R.D., Taylor, I.G., 2011. Adjusting for bias due to variability of estimated recruitments in fishery
674 assessment models. *Can. J. Fish. Aquat. Sci.* 68, 1744–1760. <https://doi.org/10.1139/f2011-092>

675 Methot, R.D., Wetzel, C.R., 2013. Stock synthesis: A biological and statistical framework for fish stock
676 assessment and fishery management. *Fish. Res.* 142, 86–99. <https://doi.org/10.1016/j.fishres.2012.10.012>

677 Miller, T.J., Hare, J.A., Alade, L.A., 2016. A state-space approach to incorporating environmental effects on
678 recruitment in an age-structured assessment model with an application to southern New England yellowtail
679 flounder. *Can. J. Fish. Aquat. Sci.* 73, 1261–1270. <https://doi.org/10.1139/cjfas-2015-0339>

680 Miller, T.J., Hyun, S.-Y., 2018. Evaluating evidence for alternative natural mortality and process error
681 assumptions using a state-space, age-structured assessment model. *Can. J. Fish. Aquat. Sci.* 75, 691–703.
682 <https://doi.org/10.1139/cjfas-2017-0035>

683 Miller, T.J., Legault, C.M., 2015. Technical details for ASAP version 4 (No. Ref Doc. 15-17). US Dept
684 Commer, Northeast Fish Sci Cent.

685 Miller, T.J., O'Brien, L., Fratantoni, P.S., 2018. Temporal and environmental variation in growth and
686 maturity and effects on management reference points of Georges Bank Atlantic cod. Can. J. Fish. Aquat.
687 Sci. 75, 2159–2171. <https://doi.org/10.1139/cjfas-2017-0124>

688 Miller, T.J., Stock, B.C., 2020. The Woods Hole Assessment Model (WHAM).

689 Möllmann, C., Lindegren, M., Blenckner, T., Bergström, L., Casini, M., Diekmann, R., Flinkman, J.,
690 Müller-Karulis, B., Neuenfeldt, S., Schmidt, J.O., Tomczak, M., Voss, R., Gårdmark, A., 2014. Implementing
691 ecosystem-based fisheries management: From single-species to integrated ecosystem assessment and advice
692 for Baltic Sea fish stocks. ICES J. Mar. Sci. 71, 1187–1197. <https://doi.org/10.1093/icesjms/fst123>

693 Munch, S.B., Giron-Nava, A., Sugihara, G., 2018. Nonlinear dynamics and noise in fisheries recruitment: A
694 global meta-analysis. Fish Fish. 19, 964–973. <https://doi.org/10.1111/faf.12304>

695 Munch, S.B., Poynor, V., Arriaza, J.L., 2017. Circumventing structural uncertainty: A Bayesian perspective on
696 nonlinear forecasting for ecology. Ecol. Complex. 32, 134–143. <https://doi.org/10.1016/j.ecocom.2016.08.006>

697 Myers, R.A., 1998. When do environment-recruitment correlations work? Rev. Fish Biol. Fisher. 8, 285–305.

698 NEFSC, 2020a. Operational assessment of 14 Northeast groundfish stocks, updated through 2018. U.S. Dept.
699 Commer., NOAA, NMFS, NEFSC, Woods Hole, MA.

700 NEFSC, 2020b. Butterfish 2020 assessment update report. U.S. Dept. Commer., NOAA, NMFS, NEFSC,
701 Woods Hole, MA.

702 Nielsen, A., Berg, C.W., 2014. Estimation of time-varying selectivity in stock assessments using state-space
703 models. Fish. Res. 158, 96–101. <https://doi.org/10.1016/j.fishres.2014.01.014>

704 O'Leary, C.A., Miller, T.J., Thorson, J.T., Nye, J.A., 2019. Understanding historical summer flounder
705 (*Paralichthys Dentatus*) abundance patterns through the incorporation of oceanography-dependent vital rates
706 in Bayesian hierarchical models. Can. J. Fish. Aquat. Sci. 76, 1275–1294. [https://doi.org/10.1139/cjfas-](https://doi.org/10.1139/cjfas-2018-0092)
707 2018-0092

708 Patrick, W.S., Link, J.S., 2015. Myths that Continue to Impede Progress in Ecosystem-Based Fisheries
709 Management. Fisheries 40, 155–160. <https://doi.org/10.1080/03632415.2015.1024308>

710 Perreault, A.M., Zheng, N., Cadigan, N.G., 2020. Estimation of growth parameters based on length-stratified
711 age samples. Can. J. Fish. Aquat. Sci. 77, 439–450. <https://doi.org/10.1139/cjfas-2019-0129>

712 Perretti, C.T., Deroba, J.J., Legault, C.M., 2020. Simulation testing methods for estimating misreported
713 catch in a state-space stock assessment model. *ICES J. Mar. Sci.* 77, 911–920. [https://doi.org/10.1093/](https://doi.org/10.1093/icesjms/fsaa034)
714 [icesjms/fsaa034](https://doi.org/10.1093/icesjms/fsaa034)

715 Perretti, C.T., Fogarty, M.J., Friedland, K.D., Hare, J.A., Lucey, S.M., McBride, R.S., Miller, T.J., Morse,
716 R.E., O’Brien, L., Pereira, J.J., Smith, L.A., Wuenschel, M.J., 2017. Regime shifts in fish recruitment on the
717 Northeast US Continental Shelf. *Mar. Ecol. Prog. Ser.* 574, 1–11. <https://doi.org/10.3354/meps12183>

718 Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Bris, A.L., Mills, K.E., Nye, J.A., Record,
719 N.R., Scannell, H.A., Scott, J.D., Sherwood, G.D., Thomas, A.C., 2015. Slow adaptation in the face of rapid
720 warming leads to collapse of the Gulf of Maine cod fishery. *Science* 350, 809–812. [https://doi.org/10.1126/](https://doi.org/10.1126/science.aac9819)
721 [science.aac9819](https://doi.org/10.1126/science.aac9819)

722 Punt, A.E., A’mar, T., Bond, N.A., Butterworth, D.S., de Moor, C.L., De Oliveira, J.A.A., Haltuch, M.A.,
723 Hollowed, A.B., Szuwalski, C., 2014. Fisheries management under climate and environmental uncertainty:
724 Control rules and performance simulation. *ICES J. Mar. Sci.* 71, 2208–2220. [https://doi.org/10.1093/](https://doi.org/10.1093/icesjms/fst057)
725 [icesjms/fst057](https://doi.org/10.1093/icesjms/fst057)

726 R Core Team, 2020. R: A Language and Environment for Statistical Computing.

727 Rose, G.A., Rowe, S., 2015. Northern cod comeback. *Can. J. Fish. Aquat. Sci.* 72, 1789–1798. <https://doi.org/10.1139/cjfas-2015-0346>
728

729 Schnute, J.T., Haigh, R., 2007. Compositional analysis of catch curve data, with an application to *Sebastes*
730 *maliger*. *ICES J. Mar. Sci.* 64, 218–233. <https://doi.org/10.1093/icesjms/fsl024>

731 Schnute, J.T., Richards, L.J., 1995. The influence of error on population estimates from catch-age models.
732 *Can. J. Fish. Aquat. Sci.* 52, 2063–2077. <https://doi.org/10.1139/f95-800>

733 Shelton, P.A., Sinclair, A.F., Chouinard, G.A., Mohn, R., Duplisea, D.E., 2006. Fishing under low productivity
734 conditions is further delaying recovery of Northwest Atlantic cod (*Gadus Morhua*). *Can. J. Fish. Aquat. Sci.*
735 63, 235–238. <https://doi.org/10.1139/f05-253>

736 Shotwell, S.K., Hanselman, D.H., Belkin, I.M., 2014. Toward biophysical synergy: Investigating advection
737 along the Polar Front to identify factors influencing Alaska sablefish recruitment. *Deep-Sea Res. Pt. II.* 107,
738 40–53. <https://doi.org/10.1016/j.dsr2.2012.08.024>

739 Stock, B.C., Xu, H., Miller, T.J., Thorson, J.T., Nye, J.A., 2021. Implementing two-dimensional autocorrela-
740 tion in either survival or natural mortality improves a state-space assessment model for Southern New England-
741 Mid Atlantic yellowtail flounder. *Fish. Res.* 237, 105873. <https://doi.org/10.1016/j.fishres.2021.105873>

742 Stock, C.A., Alexander, M.A., Bond, N.A., Brander, K.M., Cheung, W.W., Curchitser, E.N., Delworth, T.L.,
743 Dunne, J.P., Griffies, S.M., Haltuch, M.A., Hare, J.A., Hollowed, A.B., Lehodey, P., Levin, S.A., Link, J.S.,
744 Rose, K.A., Rykaczewski, R.R., Sarmiento, J.L., Stouffer, R.J., Schwing, F.B., Vecchi, G.A., Werner, F.E.,
745 2011. On the use of IPCC-class models to assess the impact of climate on Living Marine Resources. *Prog.*
746 *Oceanogr.* 88, 1–27. <https://doi.org/10.1016/j.pocean.2010.09.001>

747 Sullivan, P.J., 1992. A Kalman filter approach to catch-at-length analysis. *Biometrics* 48, 237–257.

748 Tableau, A., Collie, J.S., Bell, R.J., Minto, C., 2018. Decadal changes in the productivity of New England
749 fish populations. *Can. J. Fish. Aquat. Sci.* 76, 1528–1540. <https://doi.org/10.1139/cjfas-2018-0255>

750 Thompson, W.R., 1936. On confidence ranges for the median and other expectation distributions for
751 populations of unknown distribution form. *Ann. Math. Statist.* 7, 122–128. [https://doi.org/10.1214/aoms/](https://doi.org/10.1214/aoms/1177732502)
752 [1177732502](https://doi.org/10.1214/aoms/1177732502)

753 Thorson, J.T., 2019. Perspective: Let’s simplify stock assessment by replacing tuning algorithms with
754 statistics. *Fish. Res.* 217, 133–139. <https://doi.org/10.1016/j.fishres.2018.02.005>

755 Tommasi, D., Stock, C.A., Hobday, A.J., Methot, R., Kaplan, I.C., Eveson, J.P., Holsman, K., Miller, T.J.,
756 Gaichas, S., Gehlen, M., Pershing, A., Vecchi, G.A., Msadek, R., Delworth, T., Eakin, C.M., Haltuch, M.A.,
757 Séférian, R., Spillman, C.M., Hartog, J.R., Siedlecki, S., Samhuri, J.F., Muhling, B., Asch, R.G., Pinsky,
758 M.L., Saba, V.S., Kapnick, S.B., Gaitan, C.F., Rykaczewski, R.R., Alexander, M.A., Xue, Y., Pegion, K.V.,
759 Lynch, P., Payne, M.R., Kristiansen, T., Lehodey, P., Werner, F.E., 2017. Managing living marine resources
760 in a dynamic environment: The role of seasonal to decadal climate forecasts. *Prog. Oceanogr.* 152, 15–49.
761 <https://doi.org/10.1016/j.pocean.2016.12.011>

762 Walters, C.J., Collie, J.S., 1988. Is research on environmental factors useful to fisheries management? *Can. J.*
763 *Fish. Aquat. Sci.* 45, 1848–1854. <https://doi.org/10.1139/f88-217>

764 Williams, E.H., Shertzer, K.W., 2015. Technical documentation of the Beaufort Assessment Model (BAM).
765 U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SEFSC-671.

766 Winemiller, K.O., Rose, K.A., 1992. Patterns of life-history diversification in North American fishes:
767 Implications for population regulation. *Can. J. Fish. Aquat. Sci.* 49, 2196–2218. [https://doi.org/10.1139/f92-](https://doi.org/10.1139/f92-242)
768 [242](https://doi.org/10.1139/f92-242)

769 Xu, H., Miller, T.J., Hameed, S., Alade, L.A., Nye, J.A., 2018. Evaluating the utility of the Gulf Stream
770 Index for predicting recruitment of Southern New England-Mid Atlantic yellowtail flounder. *Fish. Oceanogr.*
771 27, 85–95. <https://doi.org/10.1111/fog.12236>

772 Xu, H., Thorson, J.T., Methot, R.D., 2020. Comparing the performance of three data weighting methods when
 773 allowing for time-varying selectivity. *Can. J. Fish. Aquat. Sci.* 77, 247–263. [https://doi.org/10.1139/cjfas-](https://doi.org/10.1139/cjfas-2019-0107)
 774 2019-0107

775 Xu, H., Thorson, J.T., Methot, R.D., Taylor, I.G., 2019. A new semi-parametric method for autocorrelated
 776 age- and time-varying selectivity in age-structured assessment models. *Can. J. Fish. Aquat. Sci.* 76, 268–285.
 777 <https://doi.org/10.1139/cjfas-2017-0446>

778 Zwolinski, J.P., Demer, D.A., 2012. A cold oceanographic regime with high exploitation rates in the
 779 Northeast Pacific forecasts a collapse of the sardine stock. *Proc. Natl. Acad. Sci. U.S.A.* 109, 4175–4180.
 780 <https://doi.org/10.1073/pnas.1113806109>

Appendix A

Table A1 lists the stock-recruit functions, f , available in WHAM, along with possible effects of a modeled environmental covariate, X_y . Note that the user must specify the lag, i.e. number of time steps, ψ , at which X_y affects recruitment (where $\psi = 1$ is common and indicates that X_y affects recruitment the following year). Then

$$\log R_y \sim \mathcal{N} \left(\log [f(SSB_{y-1}, X_{y-\psi})] - \frac{\sigma_R^2}{2}, \sigma_R^2 \right)$$

where SSB_{y-1} is the spawning stock biomass in year $y - 1$. In Table A1, we have abbreviated SSB_{y-1} to S and $X_{y-\psi}$ to X . Each of the environmental effect terms, $e^{\gamma X}$, can be extended to include polynomials as in Eqn. 6, i.e. $e^{\gamma_1 X + \gamma_2 X^2 + \dots}$.

Table A1: Stock-recruit functions available in WHAM with possible environmental effects.

| Recruit model | Environmental link | | | |
|-------------------|--------------------------------|---|---|---|
| | None | Controlling | Limiting | Masking |
| Random walk | R_{y-1} | | | |
| Random about mean | R_0 | $e^{\gamma X} R_0$ | | |
| Beverton-Holt | $\frac{\alpha S}{1 + \beta S}$ | $\frac{\alpha S e^{\gamma X}}{1 + \beta S}$ | $\frac{\alpha S}{1 + \beta S e^{\gamma X}}$ | $\frac{\alpha S}{e^{\gamma X} + \beta S}$ |
| Ricker | $\alpha S e^{-\beta S}$ | $\alpha S e^{-\beta S + \gamma X}$ | | $\alpha S e^{-\beta S(1 + \gamma X)}$ |

The environment-recruitment link terminology follows Fry (1971) and Iles and Beverton (1998):

- “controlling”: density-independent mortality,
- “limiting”: carrying capacity effect, e.g. X_y determines the amount of suitable habitat,
- “lethal”: threshold effect, i.e. R_y goes to 0 at some X_y value,
- “masking”: X_y decreases $dR/dSSB$, as expected if X_y affects metabolism or growth, and
- “directive”: e.g. behavioral.

Appendix B

Below we list the available age composition likelihood equations currently implemented in WHAM. For each dataset d and year y , the predicted proportion at age vector is $\hat{\mathbf{p}}_{d,y}$ and the observed proportion at age vector is $\mathbf{p}_{d,y}^*$. The effective sample sizes, $\text{Neff}_{d,y}$, are treated as fixed. The number of non-zero age classes, $n_{d,y}$, youngest non-zero age class, $a_{d,y}^*$, and oldest non-zero age class, $A_{d,y}^*$, also vary by d and y . In the equations below, all d and y have been dropped for simplicity, i.e. they represent the likelihood of a single observed $\mathbf{p}_{d,y}^*$ for a given dataset and year.

The predicted proportion vector is

$$\hat{p}_{d,y,a} = \frac{\hat{C}_{a,y,d}}{\sum_{a=1}^A \hat{C}_{a,y,d}}$$

if the dataset is catch, and

$$\hat{p}_{d,y,a} = \frac{\hat{I}_{a,y,d}}{\sum_{a=1}^A \hat{I}_{a,y,d}}$$

if the dataset is a survey, where $\hat{C}_{a,y,d}$ and $\hat{I}_{a,y,d}$ are defined in Eqns. 7 and 9.

Multinomial

$$-\log \mathcal{L}(\hat{\mathbf{p}} | \mathbf{p}^*, \text{Neff}) = \Gamma(\text{Neff} + 1) - \sum_a \Gamma(p_a^* \text{Neff} + 1) + \sum_a p_a^* \text{Neff} \cdot \log \hat{p}_a$$

with no estimated parameters (`age_comp = "multinomial"`).

Dirichlet-Multinomial

$$-\log \mathcal{L}(\hat{\mathbf{p}}, \phi | \mathbf{p}^*, \text{Neff}) = \Gamma(\text{Neff} + 1) + \Gamma(\phi) - \Gamma(\text{Neff} + \phi) - \sum_a [\Gamma(p_a^* \text{Neff} + 1) - \Gamma(p_a^* \text{Neff} + \phi \hat{p}_a) + \Gamma(\phi \hat{p}_a)]$$

where $\phi > 0$ is an estimated parameter (`age_comp = "dir-mult"`).

Dirichlet

$$-\log \mathcal{L}(\hat{\mathbf{p}}, \phi | \mathbf{p}^*) = \Gamma(\phi) - \sum_a [\Gamma(\phi \hat{p}_a) + (\phi \hat{p}_a - 1) \log p_a^*]$$

810 where $\phi > 0$ is an estimated parameter (`age_comp = "dirichlet"`).

811 **Logistic normal**

812 WHAM implements the logistic normal as in Miller et al. (2016). In a given year y and dataset (fishery or
813 survey) d , the negative log-likelihood of the predicted proportion at age vector, $\hat{\mathbf{p}}_{d,y}$, and observation error
814 variance, τ_d^2 , given the observed proportion at age vector, $\mathbf{p}_{d,y}^*$, is

$$v_{d,y} = \frac{\tau_d^2}{\text{Neff}_{d,y}}$$

$$-\log \mathcal{L}(\hat{\mathbf{p}}, v | \mathbf{p}^*) = \frac{n-1}{2} \log(2\pi v) + \frac{1}{2v} \sum_{a=a^*}^{n-1} \left(\log \frac{p_a^*}{p_{A^*}^*} - \log \frac{\hat{p}_a}{\hat{p}_{A^*}} \right)^2$$

815 where n , a^* , and A^* are the number of non-zero age classes, and the youngest and oldest non-zero age
816 classes in dataset d and year y (subscripts d and y dropped for simplicity). $\text{Neff}_{d,y}$ is an effective sample
817 size that can scale the estimated observation error variance, τ_d^2 , for dataset d based on changes in sampling
818 effort in time. Zero observations can either be pooled with adjacent non-zero age classes (`age_comp =`
819 `"logistic-normal-pool0"`) or treated as missing (`age_comp = "logistic-normal-miss0"`).

Table 1: Model descriptions and estimated parameters. Parameter descriptions and equations are given in text. Note that the base model in the M module is NAA-1, and the base model in the Selectivity and Ecov-Recruitment modules is NAA-3. Ecov-1 fits the Cold Pool Index data and estimates σ_x in order to allow comparison to Ecov-2 through Ecov-5 using AIC (same data needed in likelihood).

| Model | Description | Parameters estimated | No. |
|--------------------------------|--|---|-----------------|
| Numbers-at-age (NAA) | | | |
| Base | Recruitment deviations are fixed effects | R_y for $y > 1$ | $n_{years} - 1$ |
| NAA-1: Indep. R_y | Recruitment deviations are independent random effects | σ_R | 1 |
| NAA-2: AR(1) R_y | Recruitment deviations are autocorrelated, AR(1), random effects | σ_R, ρ_{year} | 2 |
| NAA-3: Indep. $N_{a,y}$ | All NAA deviations are independent random effects | σ_R, σ_a | 2 |
| NAA-4: 2D AR(1) $N_{a,y}$ | All NAA deviations are random effects with correlation by year and age, 2D AR(1) | $\sigma_R, \sigma_a, \rho_{year}, \rho_{age}$ | 4 |
| Natural mortality (M) | | | |
| M-1: None | No random effects on M | σ_R | 1 |
| M-2: Indep. | M deviations are independent random effects | σ_R, σ_M | 2 |
| M-3: 2D AR(1) | M deviations are random effects with correlation by year and age, 2D AR(1) | $\sigma_R, \sigma_M, \varphi_{year}, \varphi_{age}$ | 4 |
| Selectivity (Sel) | | | |
| Sel-1: None | No random effects on selectivity | σ_R, σ_a | 2 |
| Sel-2: Indep. | Selectivity deviations are independent random effects | $\sigma_R, \sigma_a, \sigma_{Sel}$ | 3 |
| Sel-3: 2D AR(1) | Selectivity deviations are random effects with correlation by year and parameter | $\sigma_R, \sigma_a, \sigma_{Sel}, \phi_{year}, \phi_{par}$ | 5 |
| Ecov-Recruitment (Ecov) | | | |
| Ecov-1: RW-none | Ecov: random walk (RW), effect on β : none | $\sigma_R, \sigma_a, \sigma_x$ | 3 |
| Ecov-2: RW-linear | Ecov: random walk (RW), effect on β : linear | $\sigma_R, \sigma_a, \sigma_x, \beta_1$ | 4 |
| Ecov-3: RW-poly | Ecov: random walk (RW), effect on β : 2nd order polynomial (poly) | $\sigma_R, \sigma_a, \sigma_x, \beta_1, \beta_2$ | 5 |
| Ecov-4: AR(1)-linear | Ecov: autocorrelated, AR(1), effect on β : linear | $\sigma_R, \sigma_a, \sigma_x, \phi_x, \beta_1$ | 5 |
| Ecov-5: AR(1)-poly | Ecov: autocorrelated, AR(1), effect on β : 2nd order polynomial (poly) | $\sigma_R, \sigma_a, \sigma_x, \phi_x, \beta_1, \beta_2$ | 6 |

Table 2: Stocks used in simulation tests.

| Stock | Processes tested | | | | Model dim | | Biol. par. | | Stock status | | Source | | |
|---------------------------|------------------|---|-----|------|-----------|------|------------|-------|--------------|------------|--------|--------------------|--------------------|
| | NAA | M | Sel | Ecov | # | Ages | # | Years | M | σ_R | | $\frac{B}{B_{40}}$ | $\frac{F}{F_{40}}$ |
| SNEMA yellowtail flounder | x | x | | x | | 6 | | 49 | 0.2-0.4 | 1.67 | 0.01 | 0.44 | NEFSC (2020a) |
| Butterfish | x | x | | | | 5 | | 31 | 1.3 | 0.23 | 2.57 | 0.03 | NEFSC (2020b) |
| North Sea cod | x | x | | | | 6 | | 54 | 0.2-1.2 | 0.87 | 0.14 | 2.00 | ICES (2017a) |
| Icelandic herring | x | | | | | 11 | | 30 | 0.1 | 0.55 | 0.40 | 1.81 | ICES (2017b) |
| Georges Bank haddock | x | | x | | | 9 | | 86 | 0.2 | 1.65 | 5.16 | 0.12 | NEFSC (2020a) |




Figure 1: AIC differences by model and stock when fit to original datasets. Random effects were only used for one process at a time (numbers at age, natural mortality, selectivity, and recruitment linked to the Cold Pool Index, CPI). Stock abbreviations: SNEMA yellowtail flounder (SNEMAYT), North Sea cod (NScod), Icelandic herring (ICEherring), and Georges Bank haddock (GBhaddock).




Figure 2: Change in Mohn's ρ relative to the statistical catch at age model (Base) for full state-space models (numbers at all ages are random effects, NAA-3 and NAA-4). Changes in Mohn's ρ are shown for all five stocks (whisker and box ends and median line in boxplots). Reduction in Mohn's ρ_R for SNEMA yellowtail flounder NAA-4 is off chart. Mohn's ρ values for all stocks and models are given in Table S1.




Figure 3: Mean change in coefficient of variation (CV) for key quantities in model and projection years for full state-space models (numbers at all ages are random effects, NAA-3 and NAA-4) relative to NAA-1. Boxplots show mean change in CV across years for all five stocks (whisker and box ends, median line). F was fixed at 0 in projection years.




Figure 4: Abundance-at-age deviations estimated for Georges Bank haddock using alternative models of numbers-at-age (NAA) random effects. Base = a traditional statistical catch-at-age (SCAA) model. NAA-1 = only recruitment deviations are independent random effects (most similar to Base). NAA-2 = as NAA-1, but with autocorrelated recruitment deviations, AR(1). NAA-3 = all NAA deviations are independent random effects. NAA-4 = as NAA-3, but deviations are correlated by age ($\rho_{age} = -0.12$, 95% CI: -0.32-0.08) and year ($\rho_{year} = 0.73$, 95% CI: 0.58-0.87). The vertical dashed line indicates the terminal year in the assessment.




Figure 5: Natural mortality, $\log(M)$, estimated for Southern New England-Mid Atlantic yellowtail flounder using three random effects models. M-1 = no random effects on M . M-2 = independent M deviations. M-3 = M deviations are correlated by age ($\varphi_{age} = 0.40$, 95% CI: 0.09-0.70) and year ($\varphi_{year} = 0.63$, 95% CI: 0.31-0.94). The vertical dashed line indicates the terminal year in the assessment.




Figure 6: Selectivity estimated for Georges Bank haddock using three random effects models. Sel-1 = no random effects (constant logistic selectivity). Sel-2 = independent selectivity deviations. Sel-3 = selectivity deviations are correlated by parameter ($\phi_{par} = 0.60$, 95% CI: 0.30-0.80) and year ($\phi_{year} = 0.87$, 95% CI: 0.74-0.94).




Figure 7: Beverton-Holt stock-recruit relationships fit for Southern New England-Mid Atlantic yellowtail flounder, with and without effects of the Cold Pool Index (CPI). A) CPI estimated from the model with lowest AIC (Ecov-4, AR(1)-linear). Points are observations with 95% CI, and the line with shading is the model-estimated CPI with 95% CI. Note the increased uncertainty surrounding the CPI estimate in 2017 (no observation). B) Estimates of spawning stock biomass (SSB), recruitment, and the stock-recruit function from the model without a CPI effect, Ecov-1. C) Estimates of SSB and recruitment from Ecov-4, with an effect of the CPI on β . Lines depict the expected stock-recruit relationship in each year y , given the CPI in year $y - 1$ (color).




Figure 8: Relative error of key quantities estimated for Georges Bank haddock using four models of numbers-at-age (NAA) random effects. NAA-1 = only recruitment deviations are random effects (most similar to Base, a traditional statistical catch-at-age model), and deviations are independent. NAA-2 = as NAA-1, but with autocorrelated recruitment deviations. NAA-3 = all NAA deviations are independent random effects. NAA-4 = as NAA-3, but deviations are correlated by age and year. Points without lines indicate that 95% CI are smaller than the points.




Figure 9: Relative error of key quantities estimated for Southern New England-Mid Atlantic yellowtail flounder using three models of natural mortality (M) random effects. M-1 = no random effects on M . M-2 = independent M deviations. M-3 = M deviations are correlated by age and year. Points without lines indicate that 95% CI are smaller than the points.




Figure 10: Relative error of key quantities estimated for Georges Bank haddock using three models of selectivity random effects. Sel-1 = no random effects (constant logistic selectivity). Sel-2 = independent selectivity deviations. Sel-3 = selectivity deviations are correlated by parameter and year. Points without lines indicate that 95% CI are smaller than the points.




Figure 11: Proportion of simulations in which each model had the lowest AIC. A) Numbers-at-age (NAA), aggregated across all five stocks. B) Natural mortality (M), aggregated over two stocks (SNEMAYT and butterfish). C) Selectivity (GBhaddock). Not all estimation models converged for each simulation, even when the operating model matched (Fig. S25).