- ¹ The Woods Hole Assessment Model (WHAM): a general state-space
- assessment framework that incorporates time- and age-varying
- processes via random effects and links to environmental covariates
- Brian C. Stock¹, Timothy J. Miller¹
- ⁵ brian.stock@noaa.gov, timothy.j.miller@noaa.gov, Northeast Fisheries Science Center, National Marine
- ⁶ Fisheries Service, 166 Water Street, Woods Hole, MA 02543, USA

7

1

8 Abstract

The rapid changes observed in many marine ecosystems that support fisheries pose a challenge to stock assessment and management predicated on time-invariant productivity and considering species in isolation. In 10 single-species assessments, two main approaches have been used to account for productivity changes: allowing 11 biological parameters to vary stochastically over time (empirical), or explicitly linking population processes 12 such as recruitment (R) or natural mortality (M) to environmental covariates (mechanistic). Here, we describe the Woods Hole Assessment Model (WHAM) framework and software package, which combines these two 14 approaches. WHAM can estimate time- and age-varying random effects on annual transitions in numbers at age (NAA), M, and selectivity, as well as fit environmental time-series with process and observation errors, missing data, and nonlinear links to R and M. WHAM can also be configured as a traditional statistical catch-at-age (SCAA) model in order to easily bridge from status quo models and test them against models 18 with state-space and environmental effects, all within a single framework. We fit models with and without (independent or autocorrelated) random effects on NAA, M, and selectivity to data from five stocks with a broad range of life history, fishing pressure, number of ages, and time-series length. Models that included random effects performed well across stocks and processes, especially random effects models with a two dimensional (2D) first-order autoregressive, AR(1), covariance structure over age and year. We conducted simulation tests and found negligible or no bias in estimation of important assessment outputs (SSB, F, stock status, and catch) when the operating and estimation models matched. However, bias in SSB and F was often non-trivial when the estimation model was less complex than the operating model, especially when models without random effects were fit to data simulated from models with random effects. Bias of the variance and correlation parameters controlling random effects was also negligible or slightly negative as expected. Our results suggest that WHAM can be a useful tool for stock assessment when environmental effects on R or M, or stochastic variation in NAA transitions, M, or selectivity are of interest. In the U.S. Northeast, where the productivity of several groundfish stocks has declined, conducting assessments in WHAM with time-varying processes via random effects or environment-productivity links may

34 Keywords

- state-space; stock assessment; random effects; time-varying; environmental effects; recruitment; natural
- 36 mortality; Template Model Builder (TMB)

account for these trends and potentially reduce retrospective bias.

₃₇ 1 Introduction

The last two decades have increasingly seen a push for more holistic, ecosystem-based fisheries management (Larkin 1996; Link 2002). In part, this is due to the view that considering single species in isolation produces riskier and less robust outcomes long-term (Patrick and Link 2015). In several high-profile cases, fisheries management has failed to prevent collapses because they did not reduce fishing pressure in responses to changes in natural mortality (M), recruitment, or migration patterns caused by dynamics external to the 42 stock in question (Northern cod: Shelton et al. 2006; Rose and Rowe 2015; Gulf of Maine cod: Pershing et al. 2015; Pacific sardine: Zwolinski and Demer 2012). This is particularly concerning in the context of climate change and the wide range of biological processes—often assumed to be constant—in stock assessments that are likely to be affected (Stock et al. 2011; Tommasi et al. 2017). One approach to account for changing productivity is to explicitly link population processes to environmental covariates in single-species stock assessments, i.e. the mechanistic approach sensu Punt et al. (2014). Traditional single-species assessments assume that population dynamics are based on time-constant biological processes and variable fishing mortality (F), even though fisheries scientists have long known about important drivers of time-varying population processes, e.g. recruitment, mortality, growth, and movement (Garstang 1900; Hjort 1914). Effects of the environment or interactions with other species can be considered contextually, rather than explicitly as estimated parameters (although temporal variation in empirical weight and maturity at age can affect reference points). Despite how counterintuitive this may seem to ecologists and oceanographers who study such relationships, the evidence for direct linkages to specific environmental covariates is often weak and can break down over time (Myers 1998; McClatchie et al. 2010). Additionally, the primary goal of most assessments is to provide management advice on near-term sustainable harvest levels—not to explain ecological relationships. Even if an environmental covariate directly affects fish productivity, including the effect in an assessment may not improve management advice if the effect is weak (De Oliveira and Butterworth 2005). Moreover, including environmental effects in an assessment or management system has been shown to actually provide worse management in some cases (Walters and Collie 1988; De Oliveira and Butterworth 2005; Punt et al. 2014). This can be true even in cases of relatively well-understood mechanistic links between oceanic conditions and fish populations, as in the case of sea surface temperature and Pacific sardine (Zwolinski and Demer 2012; Hill et al. 2018). Still, incorporating mechanistic environment-productivity links in assessments does have the potential to reduce residual variance, particularly in periods when few demographic data exist (Shotwell et al. 2014; Miller et al. 2016).

57 An alternative approach is to allow biological parameters to vary stochastically over time, i.e. the empirical

approach sensu Punt et al. (2014). In this case, the variation is caused by a range of sources that are not explicitly modeled. Statistical catch-at-age (SCAA) models typically only estimate year-specific recruitment (R_y) and F_y , often as deviations from a mean, R_0 , that may or may not be a function of spawning biomass, e.g. $\log R_y = \log R_0 + \epsilon_y$. The main reason that other parameters are assumed constant is simply that there are not enough degrees of freedom to estimate many time-varying parameters. A common solution is to penalize the deviations, e.g. $\epsilon_y \sim \mathcal{N}(0, \sigma_\epsilon^2)$, although the penalty terms, σ_ϵ^2 , must be fixed or iteratively tuned and are therefore somewhat subjective (Methot and Taylor 2011; Methot and Wetzel 2013; Aeberhard et al. 2018; Xu et al. 2020). State-space models that treat parameters as unobserved states can, in principle, avoid such subjectivity by estimating the penalty terms as variance parameters constraining random effects and maximizing the marginal likelihood (Thorson 2019). In this way, state-space models can allow processes to vary in time while simultaneously estimating fewer parameters.

Although state-space stock assessments have existed for some time (Mendelssohn 1988; Sullivan 1992; Gudmundsson 1994), the recent development of Template Model Builder (TMB, Kristensen et al. 2016) software to perform efficient Laplace approximation has greatly expanded their use (Nielsen and Berg 2014; Cadigan 2016; Miller et al. 2016). In addition to the key advantage of objectively estimating variance, or "data weighting", parameters, state-space models naturally predict unobserved states, and therefore handle missing data and short-term projections in a straightforward way (ICES 2020). In comparisons with SCAA models, state-space models have been shown to have larger, more realistic, uncertainty and lower retrospective bias (Miller and Hyun 2018; Stock et al. 2021).

Retrospective bias can occur when changing environmental conditions lead to changes in productivity that
are unaccounted for in stock assessments, and this is a concern common to several groundfish stocks on the
Northeast U.S. Shelf (Brooks and Legault 2016; Tableau et al. 2018). The Northeast U.S. Shelf ecosystem is
rapidly changing, and this has motivated managers to make the "continue[d] development of stock assessment
models that include environmental terms" a top priority (Hare et al. 2016). Applications of state-space
models with environmental effects on recruitment, growth, M, and maturity have proven promising (Miller et
al. 2018; Xu et al. 2018; Miller and Hyun 2018; O'Leary et al. 2019). In addition to providing short-term
(1-3 years) catch advice with reduced retrospective bias, it is hoped that environment-linked assessments will
help create realistic evaluation of sustainable stock and harvest levels in the medium-term (3-10 years) for
stocks that have not rebounded in response to dramatic decreases in F.

To address the needs of fisheries management in a changing climate, we seek an assessment framework that combines both the empirical and mechanistic approaches. Namely, it should be able to 1) estimate time-varying parameters as random effects (i.e. a state-space model), and 2) include environmental effects directly on biological parameters. The framework should also allow for easy testing against status quo SCAA models to ease gradual adoption through the "research track" or "benchmark" assessment process (Lynch et al. 2018). The objectives of this manuscript are to introduce the Woods Hole Assessment Model (WHAM) framework and demonstrate its ability to:

- 1. estimate time- and age-varying random effects on annual changes in abundance at age, M, and selectivity;
- 2. fit environmental time-series with process and observation error, missing data, and a link to a population process;
- 3. project processes treated as random effects in short-term population forecasts; and
- 4. simulate new data and random effects to conduct self- and cross-tests (sensu Deroba et al. 2015) to estimate bias in parameters and derived quantities.

Throughout, we describe how the above are implemented using the open-source WHAM software package (Miller and Stock 2020).

$_{\scriptscriptstyle{12}}$ 2 Methods

104

3 2.1 Model description

WHAM is a generalization and extension of Miller et al. (2016) in TMB. It is in many respects similar to the
Age-Structured Assessment Program (ASAP, Legault and Restrepo 1998; Miller and Legault 2015) and can
be configured to fit statistical catch-at-age models nearly identically. There is functionality built into WHAM
to migrate ASAP input files to R inputs needed for WHAM, and WHAM uses many of the same types of
data inputs, such as empirical weight-at-age, so that existing assessments in the U.S. Northeast can be easily
replicated and tested against models with state-space and environmental effects in a single framework.

2.1.1 Processes with random effects

WHAM primarily diverges from ASAP through the implementation of random effects on three processes: interannual transitions in numbers at age (NAA), natural mortality (M), and selectivity (s), as well as allowing
effects of environmental covariates (Ecov) on recruitment and natural mortality (but see ASAP4; Miller and
Legault 2015). The environmental covariates and their observations are part of the state-space framework
with true, unobserved values treated as random effects and observation on them having error. Other than
environmental covariates, the processes are assumed to have a two dimensional (2D) first-order autoregressive,

AR(1), covariance structure over age and year, although correlation in either or both dimensions can be turned 127 off. The 2D AR(1) structure has been widely used to model deviations by age and year in the parameters $F_{a,y}$ (Nielsen and Berg 2014; Kumar et al. 2020; Perreault et al. 2020), $M_{a,y}$ (Cadigan 2016; Stock et al. 2021), 129 $s_{a,y}$ (Xu et al. 2019), and $N_{a,y}$ (Stock et al. 2021), as well as in the catch $(C_{a,y})$ and survey index $(I_{a,y})$ observations (Berg and Nielsen 2016). Although other covariance structures are plausible (e.g. compound 131 symmetry, unconstrained, or order-k autoregressive; Nielsen and Berg 2014; Berg and Nielsen 2016), the 2D AR(1) structure is a simple option which requires few parameters and allows for smooth deviations in 133 time and by age; values for nearby years and ages are more correlated than distant ones. This is often a 134 reasonable null hypothesis for biological or fishery processes, but there are also clear examples where change 135 occurs abruptly, e.g. a disease outbreak increases M or changes in gear or area regulations affect selectivity 136 (ICES 2017a).

$_{\scriptscriptstyle{138}}$ 2.1.1.1 Numbers at age (NAA)

The stock equations in WHAM that describe the transitions between numbers at age are identical to Miller et al. (2016) and Nielsen and Berg (2014):

$$\log N_{a,y} = \begin{cases} \log (f(SSB_{y-1})) + \varepsilon_{1,y}, & \text{if } a = 1\\ \log (N_{a-1,y-1}) - Z_{a-1,y-1} + \varepsilon_{a,y}, & \text{if } 1 < a < A\\ \log (N_{A-1,y-1}e^{-Z_{A-1,y-1}} + N_{A,y-1}e^{-Z_{A,y-1}}) + \varepsilon_{A,y}, & \text{if } a = A \end{cases}$$
(1)

where $N_{a,y}$ are the numbers at age a in year y, Z is the total mortality rate (F+M), f is the stock-recruit function (for options see Appendix A), and A represents the plus-group. The $\varepsilon_{a,y}$ deviations are akin to survival, which is clearly related to $M_{a,y}$, but they can also be caused by mis-specification of selectivity of the fishery or indices, movement into or out of the population, or misreported catch (Schnute and Richards 1995; Gudmundsson and Gunnlaugsson 2012; Perretti et al. 2020; Stock et al. 2021). In this analysis we consider the SCAA model, "Base", and then examine four possible models that treat the $\varepsilon_{a,y}$ deviations as random effects.

Base is a SCAA model, which estimates recruitment deviations, $\varepsilon_{1,y}$, in each year y > 1 as unconstrained fixed effect parameters. The NAA deviations for ages a > 1 are fixed at zero so that transitions are deterministic.

NAA-1 is most similar to Base, where only recruitment deviations, $\varepsilon_{1,y}$, are estimated. However, in NAA-1 the recruitment deviations are assumed to be independent and identically distributed (IID) random effects,

$$\varepsilon_{1,y} \sim \mathcal{N}\left(-\frac{\sigma_R^2}{2}, \sigma_R^2\right),$$

with variance σ_R^2 as an estimated parameter. The $-\frac{\sigma_R^2}{2}$ bias correction term is included by default so that the expected recruitment, $E(N_{1,y}=R_y)$ equals the expected R_y from the stock-recruit function (Methot and Taylor 2011; Thorson 2019). This bias correction adjustment can also be turned off.

NAA-2 is the same as NAA-1, except that the recruitment deviations are stationary AR(1) with autocorrelation parameter $-1 < \rho_{year} < 1$:

$$\varepsilon_{1,y+1} \sim \mathcal{N}\left(\rho_{year}\varepsilon_{1,y} - \frac{\sigma_R^2}{2(1-\rho_{year}^2)}, \sigma_R^2\right)$$

NAA-3 is the "full state-space" model from Nielsen and Berg (2014) and Miller et al. (2016), where all numbers at age are independent random effects and:

$$\varepsilon_{a,y} \sim \begin{cases} \mathcal{N}\left(-\frac{\sigma_R^2}{2}, \sigma_R^2\right), & \text{if } a = 1\\ \mathcal{N}\left(-\frac{\sigma_a^2}{2}, \sigma_a^2\right), & \text{if } a > 1 \end{cases}$$
(2)

where σ_a^2 for all ages a>1 are assumed to be the same but different from age a=1, i.e. recruitment.

This assumption is sensible because variability of deviations between expected and realized recruitment are typically larger than deviations from expected abundance at older ages.

NAA-4 treats the numbers at all ages as random effects, as in NAA-3, but the NAA deviations, $\varepsilon_{a,y}$, have a

2D stationary AR(1) structure as in Stock et al. (2021):

$$\mathbf{E} \sim \mathcal{MVN} (0, \Sigma)$$

where $\mathbf{E} = (\varepsilon_{1,1}, \dots, \varepsilon_{1,Y-1}, \varepsilon_{2,1}, \dots, \varepsilon_{2,Y-1}, \dots, \varepsilon_{A,1}, \dots, \varepsilon_{A,Y-1})'$ is a vector of all NAA deviations, Y is the total number of observation and prediction years, and Σ is the covariance matrix of \mathbf{E} defined by:

$$Cov\left(\varepsilon_{a,y}, \varepsilon_{\tilde{a},\tilde{y}}\right) = \frac{\sigma_{a}\sigma_{\tilde{a}}\rho_{age}^{|a-\tilde{a}|}\rho_{year}^{|y-\tilde{y}|}}{\left(1 - \rho_{age}^{2}\right)\left(1 - \rho_{uear}^{2}\right)}$$

and $-1 < \rho_{age} < 1$ and $-1 < \rho_{year} < 1$ are the AR(1) coefficients in age and year, respectively. As in NAA-3, σ_a^2 for all ages a > 1 are assumed to be the same but different from age a = 1, σ_R^2 . The bias correction term for age a > 1 in NAA-4 is $-\frac{\sigma_a^2}{2(1-\rho_{year}^2)(1-\rho_{age}^2)}$.

169 2.1.1.2 Natural mortality (M)

For natural mortality, there are mean parameters for each age, μ_{M_a} , each of which may be estimated freely or fixed at the initial values. The μ_{M_a} may also be estimated in sets of ages, e.g. estimate one mean M shared across ages 3-5, $\mu_{M_3} = \mu_{M_4} = \mu_{M_5}$. There is also an option for M to be specified as a function of weight-at-age, $M_{a,y} = \mu_M W_{a,y}^b$, as in Lorenzen (1996) and Miller and Hyun (2018). Regardless of whether μ_{M_a} are fixed or estimated, WHAM can also be configured to estimate deviations in M, $\delta_{a,y}$, as random effects analogous to the NAA deviations (Cadigan 2016; Stock et al. 2021):

$$\log(M_{a,y}) = \mu_{M_a} + \delta_{a,y}$$

$$\operatorname{Cov}(\delta_{a,y}, \delta_{\tilde{a},\tilde{y}}) = \frac{\sigma_M^2 \varphi_{age}^{|a-\tilde{a}|} \varphi_{year}^{|y-\tilde{y}|}}{(1 - \varphi_{age}^2)(1 - \varphi_{year}^2)}$$
(3)

where σ_M^2 , φ_{age} , and φ_{year} are the AR(1) variance and correlation coefficients in age and year, respectively.

In this analysis, we demonstrate three alternative M random effects models. For simplicity, all models treat μ_{M_a} as known, as in most of the original assessments. M-1 is identical to the NAA-1 model, with no random effects on M ($\delta_{a,y}=0$ and not estimated). M-2 allows IID M deviations, estimating σ_M^2 but fixing $\varphi_{age}=\varphi_{year}=0$. M-3 estimates the full 2D AR(1) structure for M deviations.

As in ASAP and many other SCAA assessment frameworks, WHAM assumes separability in the fishing

2.1.1.3 Selectivity (Sel)

mortality rate by age and year, e.g. $F_{a,y} = F_y s_a$, where F_y is the "fully selected" fishing mortality rate in 183 year y and s_a is the selectivity at age a. We note that this differs from the approach in SAM (Nielsen and Berg 2014), where the $F_{a,y}$ are estimated directly as multivariate random effects without the separability 185 assumption. Three parametric forms are available (logistic, double-logistic, and decreasing-logistic), as well as a non-parametric option to estimate each s_a individually ("age-specific"). WHAM estimates selectivity 187 parameters on the logit scale to avoid boundary problems during estimation. To allow for abrupt temporal 188 changes in selectivity as in ASAP, WHAM can estimate selectivity in user-specified time blocks. WHAM can also estimate smooth changes in selectivity using autoregressive random effects as described below. 190 WHAM estimates annual full F_y and mean selectivity parameters as fixed effects. Deviations in selectivity 191 parameters can be estimated as random effects, $\zeta_{p,y}$, with autocorrelation by parameter (p), year (y), both, 192 or neither. This is done similarly to Xu et al. (2019), except that the deviations are placed on the parameters 193 instead of the mean $s_{a,y}$ in order to guarantee that $0 < s_{a,y} < 1$. For example, logistic selectivity with two 194 parameters a_{50} and k is estimated as: 195

$$s_{a,y} = \frac{1}{1+e^{-(a-a_{50}y)/k_y}}$$

$$a_{50y} = l_{a_{50}} + \frac{u_{a_{50}} - l_{a_{50}}}{1+e^{-(\nu_1 + \zeta_{1,y})}}$$

$$k_y = l_k + \frac{u_k - l_k}{1+e^{-(\nu_2 + \zeta_{2,y})}}$$

$$\operatorname{Cov}(\zeta_{1,y}, \zeta_{2,\tilde{y}}) = \frac{\sigma_s^2 \phi_{par} \phi_{year}^{|y-\tilde{y}|}}{(1-\phi_{par}^2)(1-\phi_{year}^2)}$$

$$(4)$$

where ν_1 is the logit-scale mean a_{50} parameter with lower and upper bounds $l_{a_{50}}$ and $u_{a_{50}}$, ν_2 is the logit-scale mean k parameter with lower and upper bounds l_k and u_k , σ_s^2 is the AR(1) variance, and ϕ_{par} , and ϕ_{year} are the AR(1) correlation coefficients by parameter and year.

Below, we demonstrate three models with random effect deviations on logistic selectivity, akin to those for M.

Sel-1 treats all numbers at age as independent random effects (i.e. NAA-3) but with no random effects on s($\zeta_{p,y} = 0$ and not estimated). Sel-2 allows IID s random effects, estimating σ_s^2 but fixing $\phi_{par} = \phi_{year} = 0$.

Sel-3 estimates the full 2D AR(1) structure for s random effects.

203 2.1.1.4 Environmental covariates (*Ecov*)

WHAM models environmental covariate data using state-space models with process and observation components. The true, unobserved values (or "latent states", X_y) are then linked to the population dynamics
equations with user-specified lag. For example, recruitment in year y may be influenced by X_{y-1} (lag 1),
while natural mortality in year y may be influenced by X_y (lag 0). Multiple environmental covariates may be
included, but currently only as independent processes. The environmental and population model years do not
need to match, and missing years are allowed. In particular, including environmental data in the projection
period can be useful.

2.1.1.4.1 *Ecov* process model

There are currently two options in WHAM for the Ecov process model: a normal random walk and AR(1).

We model the random walk as in Miller et al. (2016):

$$X_{y+1}|X_y \sim \mathcal{N}\left(X_y, \sigma_X^2\right)$$

where σ_X^2 is the process variance and X_1 is estimated as a fixed effect parameter. One disadvantage of the random walk is that it is nonstationary. In short-term projections, \hat{X}_y will be equal to the last estimate with an observation and the uncertainty of \hat{X}_y will increase over time. If \hat{X}_y influences the dynamices of the

population and reference points, this leads to increasing uncertainty in projections of the stock and status over time as well (Miller et al. 2016). For these reason, we generally prefer to model X_y as a stationary AR(1) process as in Miller et al. (2018):

$$X_1 \sim \mathcal{N}\left(\mu_X, \frac{\sigma_X^2}{1 - \phi_X^2}\right)$$

$$X_y \sim \mathcal{N}\left(\mu_X(1 - \phi_X) + \phi_X X_{y-1}, \sigma_X^2\right)$$
(5)

where μ_X , σ_X^2 , and $|\phi_X| < 1$ are the marginal mean, conditional variance, and autocorrelation parameters. In addition to stationarity, another important difference between the random walk and AR(1) in short-term population projections is that the AR(1) will gradually revert to the mean over time, unless environmental covariate observations are included in the projection period.

224 2.1.1.4.2 Ecov observation model

The environmental covariate observations, x_y , are assumed to be normally distributed with mean X_y and variance $\sigma_{x_y}^2$:

$$x_y|X_y \sim \mathcal{N}\left(X_y, \sigma_{x_y}^2\right)$$

The observation variance in each year, $\sigma_{x_y}^2$, can be treated as known with year-specific values (as in Miller et al. 2016) or one overall value shared among years. They can also be estimated as parameters, likewise either as yearly values or one overall value. If yearly $\sigma_{x_y}^2$ estimates are desired, WHAM estimates the hyperparameters μ_{σ_x} and σ_{σ_x} as fixed effects and treats the $\sigma_{x_y}^2$ as random effects:

$$\sigma_{x_y}^2 \sim \mathcal{N}\left(\mu_{\sigma_x}, \sigma_{\sigma_x}^2\right)$$

231 **2.1.1.4.3** Link to population

WHAM currently provides options to link the modeled environmental covariate, X_y , to the population dynamics via recruitment or natural mortality. It is also sometimes useful to fit the Ecov model without a link to the population dynamics so that models with and without environmental effects have the same data in the likelihood and can be compared via AIC.

In the case of recruitment, the options follow the framework laid out by Fry (1971) and Iles and Beverton

(1998): "controlling" (density-independent mortality), "limiting" (carrying capacity effect, e.g. X_y determines

the amount of suitable habitat), "lethal" (threshold effect, i.e. R_y goes to 0 at some X_y value), "masking" (X_y decreases dR/dSSB, as expected if X_y affects metabolism or growth), and "directive" (e.g. behavioral). Of these, WHAM currently allows controlling, limiting, or masking effects in the Beverton-Holt stock-recruit function, and controlling or masking effects in the Ricker function (see Appendix A for all equations). For natural mortality, environmental effects are placed on μ_M , shared across ages by default.

Regardless of where the environment-population link is, the effect can be either linear or polynomial. Nonlinear effects of environmental covariates are common in ecology, and quadratic effects are to be expected in cases where intermediate values are optimal (Brett 1971; Agostini et al. 2008). WHAM includes a function to calculate orthogonal polynomials in TMB, akin to the poly() function in R. Orthogonal polynomials remove correlation between covariates, e.g. X and X^2 , which improves estimability and allows for straightforward

249 In this analysis, we compare five models with limiting effects on Beverton-Holt recruitment:

evaluation of the significance of adding X^2 to a model that includes X.

$$\hat{R}_{y+1} = \frac{\alpha SSB_y}{1 + e^{\beta_0 + \beta_1 X_y + \beta_2 X_y^2} SSB_y}$$

$$\tag{6}$$

where SSB_y is spawning stock biomass in year y, α and β_0 are the standard parameters of the Beverton-Holt function, and β_1 and β_2 are polynomial effect terms that modify β_0 based on the value of the estimated environmental covariate, X_y . Eqn. 6 is one of several stock-recruit functions with environmental links implemented in WHAM (see Appendix A for complete list). Ecov-1 treats X_y as a random walk ($\phi_X = 1$) but does not include an effect on recruitment ($\beta_1 = \beta_2 = 0$). We include Ecov-1 in order to compare AIC of the original model to those with environmental effects on recruitment. Ecov-2 and Ecov-3 also treat X_y as a random walk, but Ecov-2 estimates β_1 and Ecov-3 estimates both β_1 and β_2 . Ecov-4 and Ecov-5 estimate X_y as an AR(1) process instead of a random walk (estimate ϕ_X), and Ecov-4 estimates β_1 and Ecov-5 estimates both β_1 and β_2 .

59 2.1.2 Population observation model

Like ASAP, there are likelihood components for annual aggregate catch and relative abundance observations for each fleet and index, and annual age compositions for each fleet and index.

262 2.1.2.1 Aggregate catch and indices

The predicted catch at age for fleet i, $\hat{C}_{a,y,i}$, is a function of $N_{a,y}$, $M_{a,y}$, $F_{y,i}$, and $s_{a,y,i}$:

$$\hat{C}_{a,y,i} = N_{a,y} \left(1 - e^{-Z_{a,y}} \right) \frac{F_{a,y,i}}{Z_{a,y}}. \tag{7}$$

The log-aggregate catch $\hat{C}_{y,i} = \sum_a \hat{C}_{a,y,i} W_{a,y,i}$ observation, where $W_{a,y,i}$ is the empirical weight at age, is assumed to have a normal distribution

$$\log(C_{y,i}) \sim \mathcal{N}\left(\log(\hat{C}_{y,i}) - \frac{\sigma_{C_{y,i}}^2}{2}, \sigma_{C_{y,i}}^2\right)$$
(8)

where the standard deviation

$$\sigma_{C_{y,i}} = e^{\eta_i} \sigma_{\tilde{C}_{y,i}}$$

is a function of an input standard deviation $\sigma_{\tilde{C}_{y,i}}$ and a fleet-specific parameter η_i that is fixed at 0 by default, but may be estimated. The bias correction term, $-\frac{\sigma_{C_{y,i}}^2}{2}$, is included by default based on Aldrin et al. (2020) but can be turned off.

Observations of aggregate indices of abundance are handled identically to the aggregate catch as in Eqn. 8 except that for index i the predicted index at age is

$$\hat{I}_{a,y,i} = q_i s_{a,y,i} N_{a,y} W_{a,y,i} e^{-Z_{a,y} f_{y,i}}.$$
(9)

where q_i is the catchability and $f_{y,i}$ is the fraction of the annual time step elapsed when the index is observed.

There are options for indices to be in terms of abundance (numbers) or biomass and $W_{a,y,i} = 1$ for the former.

2.1.2.2 Catch and index age composition

WHAM includes several options for the catch and index age compositions, including the multinomial (the
default to match ASAP), Dirichlet, Dirichlet-multinomial (Thorson 2019), and logistic normal (Aitchison
and Shen 1980; Schnute and Haigh 2007) distributions (Appendix B). Although the multinomial has been
commonly used in stock assessment, it is generally inferior to more flexible options that are self-weighting
and allow for correlations, such as the Dirichlet and logistic normal (Francis 2014). We therefore assumed a
logistic normal distribution for age composition observations in all of the applications and simulation studies
here.

2.1.3**Projections**

The default settings for short-term projections follow common practice for stock assessments in the U.S. 283 Northeast: the population is projected three years using the average selectivity, maturity, weight, and natural 284 mortality at age from the last five model years to calculate reference points (NEFSC 2020a). WHAM implements similar options as ASAP for specifying F_y in the projection years: terminal year F_y , average F_y 286 over specified years, $F_{X\%}$ (F at X% SPR, where X is specified and 40 by default), user-specified F_y , or F 287 derived from user-specified catch. For all options except user-specified F_y , the uncertainty in projected F288 is propagated into the uncertainty of projected population attributes. For models with random effects on NAA, M, or Ecov, the default is to continue the stochastic process into the projection years. WHAM does 290 not currently do this for selectivity because, like F, it is a function of management. Instead, selectivity is taken as the average of recent model years. 292 The decision whether or not to continue autocorrelated processes into the projection years is important 293 because it can substantially affect short-term SSB forecasts (Stock et al. 2021). This occurs because any non-zero NAA or M deviations near the end of the assessment are propagated into the projection period. 295 If the Ecov data extend beyond the population model years, WHAM will fit the Ecov model to all available 296 data and use the estimated X_y in projections. This may often be the case because lags can exist in both the physical-biological mechanism and the assessment process. As an example, a model with an Ecov effect on 298 recruitment may link physical oceanographic conditions, e.g. surface or bottom temperature, in year t to recruitment in year t+1, and the assessment conducted in year t may only use population data through year 300 t-1. In this case, 3-year population projections only need the Ecov model to be projected one year. While the default handling of Ecov projections is to continue the stochastic process, WHAM includes options to use 302 terminal year x_y , x_y averaged over specified years, or specified x_y . The option to specify x_y allows users to 303 investigate how alternative climate projections may affect the stock.

2.2Fits to original datasets

304

We fit the models described above to data from five stocks with a broad range of life history, status, and 306 model dimension (number of ages and years): Southern New England-Mid Atlantic (SNEMA) yellowtail flounder, butterfish, North Sea cod, Icelandic herring, and Georges Bank (GB) haddock (Tables 1 and 2). We 308 assumed the same form of logistic normal for SNEMA yellowtail flounder age composition observations as Miller et al. (2016) where unobserved ages are pooled with adjascent ages, but for other stocks unobserved 310 ages are treated as missing. The variance parameters associated with the logistic normal distributions were estimated for all stocks.

Although WHAM can treat multiple processes as random effects simultaneously (e.g. NAA and M; Stock et 313 al. 2021), here we only fit alternate models for one process at a time for simplicity. We fit the NAA random 314 effects models to all five stocks since these represent core WHAM functionality. We chose to highlight one 315 stock each for the M, Sel, and Ecov processes because all models did not converge for all stocks and processes: SNEMA yellowtail flounder (M and Ecov) and GB haddock (Sel). We fixed μ_{M_a} at the values used in the 317 original assessments, except for butterfish where it estimated in the assessment. Aside from the GB haddock Sel models, we used the same selectivity parameterization and time blocks as in the original assessments. We 319 used the NAA-3 model (all NAA deviations are random effects) in the Sel and Ecov demonstrations, but the NAA-1 model (only recruitment deviations are random effects) for the M demonstrations, because deviations 321 in the NAA transitions can be caused by unmodeled deviations in M. For the SNEMA yellowtail flounder 322 Ecov models, we used the Cold Pool Index (CPI) as calculated in Miller et al. (2016). We updated the CPI 323 through 2018, except that the 2017 observation was missing because the NEFSC fall bottom trawl survey 324 was not completed in the SNEMA region. Within each process, we compared the performance of the different random effects models using AIC. 326 We fit all models using the open-source statistical software R (R Core Team 2020) and TMB (Kristensen et al. 327 2016), as implemented in the R package WHAM (Miller and Stock 2020). Documentation and tutorials for how to specify additional random effect structures in WHAM are available at https://timjmiller.github.io/wham/. 329

332 2.3 Simulation tests

NOAA/wham-sim.

331

After fitting each model to the original datasets, we used the simulation feature of TMB to conduct self- and cross-tests (sensu Deroba et al. 2015). For each model, we generated 100 sets of new data and random effects, keeping the fixed effect parameters constant at values estimated in original fits. We then re-fit all models to datasets simulated under each as an operating model. We calculated the relative error in parameters constraining random effects (Table 1) and quantities of interest, such as spawning stock biomass (SSB), F, $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R. We calculated relative error as $\frac{\hat{\theta}_i}{\theta_i} - 1$, where θ_i is the true value for simulated dataset i and $\hat{\theta}_i$ is the value estimated from fitting the model to the simulated data. To estimate bias of each estimation model for a given operating model, we calculated 95% confidence intervals of the median relative error using the binomial distribution (Thompson 1936). To summarize the bias across simulations and years for each

Code and data files to run the analysis presented here are available at https://github.com/brianstock-

- model, we calculated the median quantity across years and took the mean of the medians across simulations.
- Finally, for each operating model we calculated the proportion of simulations in which each estimation model
- converged and had the lowest AIC.

345 Results

$_{ ext{\tiny 346}}$ 3.1 Original datasets

The 2D AR(1) covariance structure for random effects on NAA, M, and selectivity performed well across stocks and processes, evidenced by lower AIC than the IID random effects models (Fig. 1). This AIC difference was larger for selectivity than NAA or M, but the differences for NAA and M were also non-trivial, ranging from 11.1–53.2. Incorporating random effects on NAA, M or selectivity did not have consistently positive or negative effects on key assessment model output (SSB, F, or recruitment); these effects differed by stock and process (Figs. S1–S10). One pair of 2D AR(1) covariance parameters, σ and ρ_{year} , was consistently estimated with negative correlation across all stocks and processes, -0.43 on average.

3.1.1 Numbers-at-age (NAA)

Treating numbers at all ages as random effects was strongly supported by AIC, compared to treating only recruitment deviations as random effects (Fig. 1). Including autoregressive NAA random effects was also generally supported by AIC, either the AR(1) when only recruitment deviations were random effects or the 2D AR(1) when all ages were random effects. Across the five stocks, models that treated all NAA as random effects (i.e., full state-space models) also had reduced retrospective pattern and increased uncertainty in SSB, F, and recruitment (Figs. 2-3). Including 2D AR(1) autocorrelation on NAA deviations further reduced retrospective patterns and increased uncertainty on average.

The estimated AR(1) and IID random effects were similar, with the AR(1) random effects slightly smoothed compared to the IID random effects when the estimated correlations were positive (e.g. $\rho_{year} = 0.73$, 95% CI: 0.58-0.87, for Georges Bank haddock in Fig. 4).

$_{365}$ $\;\;3.1.2$ Natural mortality (M)

As for the NAA models, including random effect deviations on M was supported by AIC (Fig. 1). Although
the 2D AR(1) M model did not converge for North Sea cod, it had the lowest AIC for butterfish and SNEMA

yellowtail flounder. The M-3 model with 2D AR(1) autocorrelation estimated similar M deviations as the M-2 model, with smoothing when estimated correlations were positive (e.g. $\varphi_{year} = 0.63$, 95% CI: 0.31-0.94, and $\varphi_{age} = 0.40$, 95% CI: 0.09-0.70, for SNEMA yellowtail flounder in Fig. 5). Both M-2 and M-3 estimated elevated M for fish aged 5-6+ and lower M for ages 2-3 for several years in the middle of the time-series, and then higher M for ages 1-3 at the end of the time-series.

373 3.1.3 Selectivity (Sel)

For Georges Bank haddock, including 2D AR(1) random effect deviations on selectivity was strongly supported by AIC (Fig. 1). Sel-3 estimated similar patterns in selectivity compared to Sel-2, except with smoother variations by age and year (Fig. 6). Compared to the model with constant selectivity, Sel-2 and Sel-3 estimated lower s for age 3 in recent years and higher s for age 3 before 1990. They also estimated higher s for age 2 before 1990, especially from 1973 to 1976.

3.1.4 Environmental covariate effect on recruitment (Ecov)

As in previous analyses (Miller et al. 2016; Xu et al. 2018), including an effect of the CPI on recruitment for 380 SNEMA yellowtail flounder was clearly supported by AIC (\triangle AIC of 20.3-33.0 between Ecov-1 and Ecov-2 through Ecov-5, Fig. 1). The AR(1)-linear model (Ecov-4) had the lowest AIC—fitting the CPI using an 382 AR(1) model was preferred over the random walk (\triangle AIC of 12.7 between Ecov-2 and Ecov-4), and the quadratic term, β_2 , was deemed unnecessary (Δ AIC of 1.3 between Ecov-5 and Ecov-4). As expected, the 384 AR(1) process model estimated higher uncertainty in years with higher observation error (e.g. 1982–1992) and missing observations (2017, Fig. 7A). The CPI negatively influenced recruitment, i.e. \hat{R} was higher following years with lower CPI (lower fall bottom temperature, Fig. 7C). Including the CPI-recruitment link 387 changed the estimates of R_y by up to 30%, but in most years the relative difference in R_y was less than 10% (Figs. 7B–C and S3). The relative differences in SSB and F were small, less than 4% in most years, compared 389 to the influence from adding random effects on all NAA or M (Figs. S1–S3).

3.2 Simulation tests

Several findings were consistent across processes and stocks. When the estimation and operating model were consistent, bias of SSB, F, $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R was generally less than 2% and not significant based on confidence intervals (Figs. 8–10 and S11–S16). Bias was also generally less than 2% when more complex models were fitted to less complicated operating model simulations. In contrast, the bias was often greater

than 5% when the estimation model was less complex than the operating model, especially when models without random effects were fit to data simulated from models with random effects. Bias in SSB and F were always opposite, i.e. when SSB was biased high, F was biased low, and vice versa. Predicted catch was never biased. Bias of the variance and correlation parameters controlling random effects was generally not significant or negative, as expected using maximum likelihood estimation (Figs. S17-S20). Restricted maximum likelihood (REML) should be used if more accurate estimation of these parameters is a priority.

In cross-tests, the percentage of simulations in which AIC selected the correct model varied between 62–99% by process, stock, and operating model (Fig. 11). Estimation models more complex than the operating model were more likely to be chosen for NAA models than for M or selectivity.

405 3.2.1 Numbers-at-age

Across the five stocks, all NAA models estimated SSB, F, $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R with very minimal bias in 406 self-tests. An exception was the estimation of F for Icelandic herring, particularly for NAA-1 and NAA-2 (median relative error in F with 95% CI for NAA-1: -0.060 (-0.073, -0.047), NAA-2: -0.060 (-0.073, -0.047). 408 NAA-3: 0.015 (-0.004, 0.034), and NAA-4: 0.049 (0.017, 0.080); Figs. S14 and S21-S23). The convergence rate for most models and stocks was above 95%, with the same exception of NAA-1 and NAA-2 for Icelandic 410 herring (Fig. S20). In cross-tests, NAA-1 and NAA-2 exhibited non-trivial bias when estimating data 411 simulated from models that treated numbers at all ages as random effects. The degree of bias varied by 412 quantity and stock between -25% and 25% (Figs. 8 and S11–S14). The more complex NAA models estimated 413 all quantities without bias regardless of operating model. 414 For four of the five stocks, σ_R^2 was estimated without bias in self-tests (Fig. S17). The exception was 415 butterfish, for which σ_R^2 was negatively biased in NAA-1, NAA-2, and NAA-4 (but not NAA-3). In contrast, 416 both NAA-3 and NAA-4 estimated σ_a^2 with negative bias for all five stocks. There was no consistent pattern 417

419 3.2.2 Natural mortality

418

in bias for the correlation parameters ρ_{age} and ρ_{year} .

SSB, F, $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R were estimated without significant bias in self-tests, although less complex models exhibited bias when fit to data simulated from more complex models (Figs. 9 and S15–S16).

As for the NAA models, σ_R^2 was estimated without bias in the M model self-tests (Fig. S18). σ_M^2 was biased low, φ_{year} had no bias, and the direction of bias in φ_{age} was inconsistent.

3.2.3 Selectivity

Models with IID or 2D AR(1) s random effects estimated SSB, F, $\frac{B}{B_{40\%}}$, $\frac{F}{F_{40\%}}$, and R with little to no bias across operating models (Fig. 10). The model without s random effects, Sel-1, showed substantial bias when fit to data simulated from Sel-2 or Sel-3. For Georges Bank haddock, the variance and correlation parameters σ_a^2 , σ_{Sel}^2 , ϕ_{year} , and ϕ_{par} were all estimated with slight negative bias in self-tests of models with s random effects (Fig. S19).

430 3.2.4 Ecov-Recruitment

The random walk CPI models estimated σ_X^2 with less bias than the AR(1) CPI models (Fig. S20). β_0 was biased high in all models, although this was not significant for the model with lowest AIC (Ecov-4, AR(1)-linear). All models estimated the parameters ϕ_X , α , β_1 , and β_2 without significant bias.

434 4 Discussion

4.1 Overview

Our results suggest that the WHAM package can be a useful tool for stock assessment when environmental 436 effects on recruitment, or stochastic changes in the numbers at age transitions, selectivity, or natural mortality 437 are of interest. The simulation tests showed negligible or no bias in estimation of important assessment 438 outputs (SSB, F, stock and harvest status) when the operating and estimation models matched. The less 439 complex models, without random effects or autoregressive structure, exhibited some bias in cross-tests, while the more complex models did not. In these cases, bias in SSB and F were opposite such that predicted catch 441 was unbiased. WHAM models with independent or 2D AR(1) random effect deviations performed well across stocks and processes, which suggests that they warrant consideration in future stock-specific studies. The 2D 443 AR(1) covariance structure performed particularly well, generally having the lowest AIC and retrospective patterns (Figs. 1-2) and outperforming the model with independent random effects in cross-tests (Figs. 8-11 445 and S11-S15).

4.7 4.2 Limitations

- For simplicity, we only demonstrated alternative models for a single random effect process in each of our comparisons. WHAM can, however, treat multiple processes as random effects simultaneously. For SNEMA yellowtail flounder, Stock et al. (2021) found that models with process errors in both M and NAA converged and performed best by AIC and Mohn's ρ . Models with 2D autocorrelation in either process were identifiable and estimated similar F and SSB trends. Preliminary work on other stocks in the region have also shown this is possible. In general, which combinations of process errors are expected to work jointly is difficult to say without testing different combinations. The answer is very likely to be stock-specific as well, depending on the information content in the data, number of ages and years, life history, and exploitation history.
- Due to the number of models and stocks we tested, we did not thoroughly examine diagnostics, such as residual patterns, for each model fit that would be checked in a full assessment. It is possible that doing so might lead to the rejection of some model configurations for some stocks.
- Further improvements could be made to the handling of age compositions in WHAM. The current Dirichlet,
 Dirichlet-multinomial, and logistic normal options are more flexible than the multinomial distribution used in
 ASAP (Francis 2014). Yet, extensions to the logistic normal are possible; most notably, additional correlation
 structures as in Francis (2014) and Martell and Lima (2015).
- Finally, some of the 2D AR(1) covariance parameters were estimated with negative bias, particularly σ_a (Figs. S17-S20). This is the expected direction of bias using maximum likelihood, and using REML instead would likely reduce it (Miller et al. 2018). This is relatively easy to do in WHAM, or more generally in TMB, by treating all parameters except the 2D AR(1) covariance parameters as random effects. This directs TMB to use the Laplace approximation to integrate over the other fixed effect parameters (treated as random effects) with flat priors (Harville 1974). We did not do this because it requires constructing and fitting the model in TMB a second time for each simulation.

4.3 Relationships to other existing assessment model frameworks

WHAM assumes separability in $F_{a,y}$, i.e. $F_{a,y} = F_y s_a$, and estimates annual full F_y as fixed effects and s_a as constant or time-varying random effects with various autoregressive assumptions possible. Most other assessment frameworks in the U.S, e.g. ASAP, Stock Synthesis (Methot and Wetzel 2013), an Assessment Model for Alaska (AMAK; Anon. 2015), and the Beaufort Assessment Model (BAM; Williams and Shertzer 2015), make this same separability assumption which is useful for specifying a fully-selected F in projections

to calculate reference points or generating catch advice. Estimating selectivity in time blocks is common practice in these frameworks. In contrast, SAM estimates $F_{a,y}$ directly as a multivariate random walk process (Nielsen and Berg 2014). The autoregressive models for selectivity parameters under certain configurations of WHAM should allow for similar F at age patterns as in SAM. Stock Synthesis allows 2D AR(1) random effects on $s_{a,y}$ instead of the parameters, and then the variance parameter is estimated by an iterative tuning algorithm (Xu et al. 2019). It is unclear whether estimating time-varying selectivity as random effects produces better results than assuming time blocks, or if there are advantages to using any of the three approaches for estimating time-varying selectivity used by WHAM, SAM, or Stock Synthesis.

Likewise, WHAM and the other U.S.-based assessment models treat catch, index, and composition observations differently than SAM. The U.S.-based assessment models treat aggregate and composition observations separately for fisheries and indices whereas SAM treats observations of catch and indices at age, $C_{a,y}$ and $I_{a,y}$, as multivariate log-normal. Separate observation models for aggregate and composition observations is natural when sampling for total catch differs from that for length and age composition.

Like SAM, WHAM can estimate interannual transitions in NAA as a random walk processes, but WHAM
can also be configured to treat these deviations as stationary autoregressive processes. Alternatively (or
simultaneously, Stock et al. 2021) WHAM can estimate deviations in natural mortality as autoregressive
processes like NCAM (Cadigan 2016). Uniquely, WHAM can model multiple environmental covariate time
series as state-space processes and include their effects, possibly nonlinearly, in various ways on recruitment or
natural mortality. Although most applications thusfar investigate effects of physical processes on demographic
parameters, indices of predation might also be considered (Marshall et al. 2019).

96 4.4 Future use of WHAM

The productivity of several groundfish stocks in the U.S. Northeast has declined in recent decades, and conducting assessments in WHAM with time-varying processes via random effects or environment-productivity links could account for these trends and potentially reduce retrospective bias (Perretti et al. 2017; Tableau et al. 2018; Stock et al. 2021). WHAM can be configured to fit SCAA models very similar to ASAP and therefore bridge between the two frameworks. Adding random effects or environmental covariates to an assessment model is a large structural change, and simulation self- and cross-tests such as we have demonstrated here should be conducted through the research track process (Lynch et al. 2018). If comparisons against status quo SCAA models prove favorable, WHAM could transition to being used in operational assessments. However, because the details of how to include time-varying processes, as well as the effect on the assessment output,

will vary by stock, this evaluation may need to be conducted on a stock-by-stock basis.

Random effects allow for changing productivity and, if they are considered as an autoregressive process, can 507 propagate the effect of these changes on assessment output in short-term projections (Stock et al. 2021). However, the AR(1) process demonstrated here trends to the mean in projections and will not predict 509 values beyond extremes in observed time-series. This is an issue worth examining because many marine ecosystems are changing to such extent that recent conditions are time-series extremes, and conditions in the 511 near future may continue to expand the range of observations (e.g. sea surface temperatures over the U.S. Northeast Shelf in the last decade; Chen et al. 2020). More complex nonstationary time-series models such 513 as autoregressive integrated moving average (ARIMA) that can forecast beyond observed values could be implemented. Nonlinear (e.g. splines) or double linear integration (Di Lorenzo and Ohman 2013) techniques 515 could also be worth pursuing, as well as time-delay embedding methods (Munch et al. 2017, 2018), which 516 not only allow for nonstationary dynamics but also do not require a specified functional form. The best 517 approach to making stochastic projections of productivity responses to environmental conditions that are 518 rapidly changing and at time-series extremes merits further research. 519

In addition to the empirical (i.e. random effects) approach, it is worth considering the mechanistic approach 520 (i.e. explicitly modeled links to environmental covariates) for a given stock whose recruitment or M is 521 suspected to have shifted due to a clear, plausible hypothesized external influence. Several factors may result 522 in a higher likelihood that the mechanistic approach is useful in an assessment: a history of overfishing (Free 523 et al. 2019; but high F can also swamp the signal of an environmental influence, Haltuch and Punt 2011), 524 more rapid environmental change, stocks at the edge of species' range, opportunistic (short-lived) species 525 (sensu Winemiller and Rose 1992), lower trophic level species, longer time series, periodic signals in which more than once cycle has been recorded (e.g. Pacific Decadal Oscillation and sardine), and stronger signals 527 (wider ranges of observed stock status and environmental conditions) (Haltuch and Punt 2011; Haltuch et al. 2019; Marshall et al. 2019; Free et al. 2019). Although developing mature mechanistic hypotheses 529 will take significant effort and collaboration with ecologists and oceanographers, these rules of thumb can guide researchers in prioritizing which stocks to investigate environment-productivity relationships. Our 531 ability to forecast and understand oceanographic and climate variables continues to improve and increasingly 532 will present opportunities for including well-founded mechanistic environmental effects in stock assessments 533 (Tommasi et al. 2017). 534

The perception of precision in model output such as projected population biomass or catch advice is affected by whether productivity componenents are treated as either constant or stochastic processes (Figs. S1– S10), much like whether certain parameters are either fixed or estimated in traditional assessment models. Modeling frameworks such as WHAM allow the performance of these alternative models to be compared and when allowing temporal variation in productivity components is justified, the greater uncertainty in the assessment output is more realistic. Moreover, model uncertainty could be included in our preception of output uncertainty by fitting ensembles of WHAM models that represent alternative states of nature (Möllmann et al. 2014; Anderson et al. 2017). Finally, the simulation capabilities of WHAM can also be useful in situations where variation in productivity is hypothesized, but information to estimate this variation is unavialable. WHAM can be configured as an operating model to simulate plausible environmentally-driven changes in recruitment or M or plausible stochastic variation in order to estimate the sensitivity of status quo models.

547 4.5 Conclusion

A major present-day challenge in fisheries is to assess and manage stocks in a changing environment. We foresee environment-linked stock assessments becoming more feasible and realistic as fisheries and oceanographic time-series lengthen and our ecological understanding deepens. We have developed WHAM with this in mind. Finally, we note that the development of TMB has been a critical advance for fisheries assessment modeling frameworks such as WHAM, allowing us to rapidly fit models that treat population and environmental processes as time-varying random effects in a state-space framework.

554 Acknowledgements

We appreciate the efforts of the organizers of the special issue and CAPAM workshop on next-generation stock assessment models held on November 4-8, 2019 in Wellington, New Zealand. We also thank Charles Perretti and two anonymous reviewers for helpful suggestions on the manuscript. This research was performed while BCS held an NRC Research Associateship award at the NEFSC under the Northeast Groundfish and Climate Initiative. We thank Chris Melrose for updating the Cold Pool Index with data through 2018, using code originally written by Jon Hare.

561 References

- Aeberhard, W.H., Mills Flemming, J., and Nielsen, A. 2018. Review of State-Space Models for Fisheries
- ⁵⁶³ Science. Annu. Rev. Stat. Appl. **5**(1): 215–235. doi:10.1146/annurev-statistics-031017-100427.
- ⁵⁶⁴ Agostini, V., Hendrix, A., Hollowed, A., Wilson, C., Pierce, S., and Francis, R. 2008. Climate-ocean
- variability and Pacific hake: A geostatistical modeling approach. Journal of Marine Systems 71: 237–248.
- doi:10.1016/j.jmarsys.2007.01.010.
- 567 Aitchison, J., and Shen, S.M. 1980. Logistic-Normal Distributions: Some Properties and Uses. Biometrika
- **67**(2): 261–272.
- Aldrin, M., Tvete, I., Aanes, S., and Subbey, S. 2020. The specification of the data model part in the SAM
- model matters. Fisheries Research **229**: 105585. doi:10.1016/j.fishres.2020.105585.
- Anderson, S.C., Cooper, A.B., Jensen, O.P., Minto, C., Thorson, J.T., Walsh, J.C., Afflerbach, J., Dickey-
- ⁵⁷² Collas, M., Kleisner, K.M., Longo, C., Osio, G.C., Ovando, D., Mosqueira, I., Rosenberg, A.A., and Selig,
- 573 E.R. 2017. Improving estimates of population status and trend with superensemble models. Fish Fish 18(4):
- 732-741. doi:10.1111/faf.12200.
- Anon. 2015. Assessment Model for Alaska Description of GUI and Instructions. Available from https:
- //github.com/afsc-assessments/AMAK/blob/master/docs/AMAK%20Documentation.pdf.
- Berg, C.W., and Nielsen, A. 2016. Accounting for correlated observations in an age-based state-space stock
- ssessment model. ICES J Mar Sci **73**(7): 1788–1797. doi:10.1093/icesjms/fsw046.
- 579 Brett, J.R. 1971. Energetic responses of salmon to temperature. A study of some thermal relations in
- the physiology and freshwater ecology of sockeye salmon (Oncorhynchus Nerka). Am Zool 11(1): 99–113.
- ⁵⁸¹ doi:10.1093/icb/11.1.99.
- Brooks, E.N., and Legault, C.M. 2016. Retrospective forecasting evaluating performance of stock projections
- ₅₈₃ for New England groundfish stocks. Can. J. Fish. Aquat. Sci. **73**(6): 935–950. doi:10.1139/cjfas-2015-0163.
- Cadigan, N.G. 2016. A state-space stock assessment model for northern cod, including under-reported catches
- and variable natural mortality rates. Canadian Journal of Fisheries and Aquatic Sciences 73(2): 296–308.
- doi:10.1139/cjfas-2015-0047.
- ⁵⁸⁷ Chen, Z., Kwon, Y.-O., Chen, K., Fratantoni, P., Gawarkiewicz, G., and Joyce, T.M. 2020. Long-Term
- 588 SST Variability on the Northwest Atlantic Continental Shelf and Slope. Geophys. Res. Lett. 47(1).
- doi:10.1029/2019GL085455.

- 590 De Oliveira, J., and Butterworth, D. 2005. Limits to the use of environmental indices to reduce risk and/or
- increase yield in the South African anchovy fishery. African Journal of Marine Science 27(1): 191–203.
- ⁵⁹² doi:10.2989/18142320509504078.
- Deroba, J.J., Butterworth, D.S., Methot, R.D., De Oliveira, J.a.A., Fernandez, C., Nielsen, A., Cadrin,
- 594 S.X., Dickey-Collas, M., Legault, C.M., Ianelli, J., Valero, J.L., Needle, C.L., O'Malley, J.M., Chang, Y.-J.,
- Thompson, G.G., Canales, C., Swain, D.P., Miller, D.C.M., Hintzen, N.T., Bertignac, M., Ibaibarriaga, L.,
- 596 Silva, A., Murta, A., Kell, L.T., de Moor, C.L., Parma, A.M., Dichmont, C.M., Restrepo, V.R., Ye, Y.,
- Jardim, E., Spencer, P.D., Hanselman, D.H., Blaylock, J., Mood, M., and Hulson, P.-J.F. 2015. Simulation
- testing the robustness of stock assessment models to error: Some results from the ICES strategic initiative on
- stock assessment methods. ICES J Mar Sci 72(1): 19–30. doi:10.1093/icesjms/fst237.
- 600 Di Lorenzo, E., and Ohman, M.D. 2013. A double-integration hypothesis to explain ocean ecosystem
- response to climate forcing. Proceedings of the National Academy of Sciences 110(7): 2496–2499.
- doi:10.1073/pnas.1218022110.
- ⁶⁰³ Francis, R.C. 2014. Replacing the multinomial in stock assessment models: A first step. Fisheries Research
- 604 **151**: 70–84. doi:10.1016/j.fishres.2013.12.015.
- Free, C.M., Thorson, J.T., Pinsky, M.L., Oken, K.L., Wiedenmann, J., and Jensen, O.P. 2019. Impacts of
- historical warming on marine fisheries production. Science **363**(6430): 979–983. doi:10.1126/science.aau1758.
- ₆₀₇ Fry, F. 1971. The effect of environmental factors on the physiology of fish. *In Fish Physiology*. Elsevier. pp.
- 608 1-98. doi:10.1016/S1546-5098(08)60146-6.
- 609 Garstang, W. 1900. The Impoverishment of the Sea. A Critical Summary of the Experimental and Statistical
- 610 Evidence bearing upon the Alleged Depletion of the Trawling Grounds. Journal of the Marine Biological
- 451 Association of the United Kingdom 6(01): 1–69. doi:10.1017/S0025315400072374.
- 612 Gudmundsson, G. 1994. Time series analysis of catch-at-age observations. Applied Statistics 43(1): 117–126.
- 613 Gudmundsson, G., and Gunnlaugsson, T. 2012. Selection and estimation of sequential catch-at-age models.
- 614 Canadian Journal of Fisheries and Aquatic Sciences **69**(11): 1760–1772. doi:10.1139/f2012-095.
- Haltuch, M.A., Brooks, E., Brodziak, J., Devine, J., Johnson, K., Klibansky, N., Nash, R., Payne,
- 616 M., Shertzer, K., Subbey, S., and Wells, B. 2019. Unraveling the recruitment problem: A review of
- environmentally-informed forecasting and management strategy evaluation. Fisheries Research 217: 198–216.
- doi:10.1016/j.fishres.2018.12.016.
- Haltuch, M.A., and Punt, A.E. 2011. The promises and pitfalls of including decadal-scale climate forcing

- of recruitment in groundfish stock assessment. Canadian Journal of Fisheries and Aquatic Sciences 68(5):
- 621 912-926. doi:10.1139/f2011-030.
- Hare, J.A., Borggaard, D.L., Friedland, K.D., Anderson, J., Burns, P., Chu, K., Clay, P.M., Collins, M.J.,
- 623 Cooper, P., Fratantoni, P.S., Johnson, M.R., Manderson, J.P., Milke, L., Miller, T.J., Orphanides, C.D.,
- and Saba, V.S. 2016. Northeast Regional Action Plan NOAA Fisheries Climate Science Strategy. NOAA
- Fisheries, Northeast Fisheries Science Center, Woods Hole, MA. Available from https://repository.library.
- noaa.gov/view/noaa/13138 [accessed 17 September 2019].
- Harville, D.A. 1974. Bayesian inference for variance components using only error contrasts. Biometrika 61(2):
- 628 383-385. doi:10.1093/biomet/61.2.383.
- 629 Hill, K.T., Crone, P.R., and Zwolinski, J.P. 2018. Assessment of the Pacific sardine resource in 2018 for U.S.
- Management in 2018-2019. US Department of Commerce. Available from https://doi.org/10.7289/V5/TM-
- 631 SWFSC-600.
- Hjort, J. 1914. Fluctuations in the great fisheries of Northern Europe viewed in the light of biological research.
- 653 Rapports et Procès-Verbaux des Réunions du Conseil Permanent International Pour L'Exploration de la Mer
- 634 **20**: 1–228.
- 655 ICES. 2017a. Report of the North Western Working Group (NWWG). ICES CM 2017/ACOM:08, 27 April-4
- 656 May 2017, Copenhagen, Denmark. Available from http://www.ices.dk/sites/pub/Publication%20Reports/
- Expert%20Group%20Report/acom/2017/NWWG/NWWG%202017%20Report.pdf.
- 6538 ICES. 2017b. Report of the working group on assessment of demersal stocks in the North Sea
- 659 and Skagerrak (2017). ICES CM 2017/ACOM:21, 26 April-5 May 2017, ICES HQ. Available
- from https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2017/
- 641 WGNSSK/01%20WGNSSK%20Report%202017.pdf.
- 642 ICES. 2020. Workshop on the review and future of state space stock assessment models in ICES (WKRFSAM).
- ICES Scientific Reports 2(32): 23p. ICES. doi:10.17895/ices.pub.6004.
- Iles, T.C., and Beverton, R.J.H. 1998. Stock, recruitment and moderating processes in flatfish. Journal of
- Sea Research **39**(1): 41–55. doi:10.1016/S1385-1101(97)00022-1.
- 646 Kristensen, K., Nielsen, A., Berg, C., Skaug, H., and Bell, B.M. 2016. TMB: Automatic differentiation and
- Laplace approximation. Journal of Statistical Software 70: 1–21. doi:10.18637/jss.v070.i05.
- 648 Kumar, R., Cadigan, N.G., Zheng, N., Varkey, D.A., and Morgan, M.J. 2020. A state-space spatial
- survey-based stock assessment (SSURBA) model to inform spatial variation in relative stock trends. 77: 21.

- Larkin, P. 1996. Concepts and issues in marine ecosystem management. Reviews in Fish Biology and Fisheries
- 6: 139-164. doi:10.1007/BF00182341.
- 652 Legault, C.M., and Restrepo, V.R. 1998. A Flexible Forward Age-Structured Assessment Program.
- 653 Link, J.S. 2002. What Does Ecosystem-Based Fisheries Management Mean? Fisheries 27(4): 5.
- 654 Lorenzen, K. 1996. The relationship between body weight and natural mortality in juvenile and adult
- 655 fish: A comparison of natural ecosystems and aquaculture. Journal of Fish Biology 49(4): 627–642.
- 656 doi:10.1111/j.1095-8649.1996.tb00060.x.
- Lynch, P.D., Methot, R.D., and Link, J.S. (Editors). 2018. Implementing a Next Generation Stock Assessment
- Enterprise. An Update to the NOAA Fisheries Stock Assessment Improvement Plan. U.S. Dep. Commer.,
- NOAA Tech. Memo. NMFS-F/ SPO-183. p. 127. doi:10.7755/TMSPO.183.
- 660 Marshall, K.N., Koehn, L.E., Levin, P.S., Essington, T.E., and Jensen, O.P. 2019. Inclusion of ecosystem
- 661 information in US fish stock assessments suggests progress toward ecosystem-based fisheries management.
- 662 ICES J Mar Sci **76**(1): 1–9. Oxford Academic. doi:10.1093/icesjms/fsy152.
- 663 Martell, S.J.D., and Lima, C.W. 2015. Technical documentation for the integrated statistical catch age model
- 664 (iSCAM). Available from https://github.com/smartell/iSCAM [accessed 23 January 2021].
- McClatchie, S., Goericke, R., Auad, G., and Hill, K. 2010. Re-assessment of the stockRecruit and tempera-
- tureRecruit relationships for Pacific sardine (Sardinops sagax). Can. J. Fish. Aquat. Sci. 67(11): 1782–1790.
- 667 NRC Research Press. doi:10.1139/F10-101.
- 668 Mendelssohn, R. 1988. Some problems in estimating population sizes from catch-at-age data. Fishery Bulletin
- 669 **86**(4): 617–630.
- 670 Methot, R.D., and Taylor, I.G. 2011. Adjusting for bias due to variability of estimated recruitments in fishery
- assessment models. Can. J. Fish. Aquat. Sci. 68(10): 1744–1760. NRC Research Press. doi:10.1139/f2011-
- 672 092.
- 673 Methot, R.D., and Wetzel, C.R. 2013. Stock synthesis: A biological and statistical framework for fish stock
- assessment and fishery management. Fisheries Research 142: 86–99. doi:10.1016/j.fishres.2012.10.012.
- Miller, T.J., Hare, J.A., and Alade, L.A. 2016. A state-space approach to incorporating environmental effects
- on recruitment in an age-structured assessment model with an application to southern New England yellowtail
- flounder. Canadian Journal of Fisheries and Aquatic Sciences 73(8): 1261–1270. doi:10.1139/cjfas-2015-0339.
- 678 Miller, T.J., and Hyun, S.-Y. 2018. Evaluating evidence for alternative natural mortality and process error

- 679 assumptions using a state-space, age-structured assessment model. Canadian Journal of Fisheries and Aquatic
- 680 Sciences **75**(5): 691–703. doi:10.1139/cjfas-2017-0035.
- 681 Miller, T.J., and Legault, C.M. 2015. Technical details for ASAP version 4. US Dept Commer, Northeast
- Fish Sci Cent. Available from doi:10.7289/V57W695G [accessed 17 October 2019].
- 683 Miller, T.J., O'Brien, L., and Fratantoni, P.S. 2018. Temporal and environmental variation in growth and
- maturity and effects on management reference points of Georges Bank Atlantic cod. Can. J. Fish. Aquat.
- 685 Sci. **75**(12): 2159–2171. doi:10.1139/cjfas-2017-0124.
- 686 Miller, T.J., and Stock, B.C. 2020. The Woods Hole Assessment Model (WHAM). Available from https:
- //timjmiller.github.io/wham/.
- Möllmann, C., Lindegren, M., Blenckner, T., Bergström, L., Casini, M., Diekmann, R., Flinkman, J., Müller-
- 689 Karulis, B., Neuenfeldt, S., Schmidt, J.O., Tomczak, M., Voss, R., and Gårdmark, A. 2014. Implementing
- ecosystem-based fisheries management: From single-species to integrated ecosystem assessment and advice
- 691 for Baltic Sea fish stocks. ICES J Mar Sci 71(5): 1187–1197. Oxford Academic. doi:10.1093/icesjms/fst123.
- ⁶⁹² Munch, S.B., Giron-Nava, A., and Sugihara, G. 2018. Nonlinear dynamics and noise in fisheries recruitment:
- 693 A global meta-analysis. Fish and Fisheries 19: 964–973. doi:10.1111/faf.12304.
- 694 Munch, S.B., Poynor, V., and Arriaza, J.L. 2017. Circumventing structural uncertainty: A
- ₆₉₅ Bayesian perspective on nonlinear forecasting for ecology. Ecological Complexity **32**: 134–143
- doi:10.1016/j.ecocom.2016.08.006.
- Myers, R.A. 1998. When do environment-recruitment correlations work? Reviews in Fish Biology and
- 698 Fisheries 8: 285–305.
- 699 NEFSC. 2020a. Operational assessment of 14 Northeast groundfish stocks, updated through 2018. U.S. Dept.
- 700 Commer., NOAA, NMFS, NEFSC, Woods Hole, MA. Available from https://s3.amazonaws.com/nefmc.org/
- ₇₀₁ 8_Prepublication-NE-GrndfshOp-Assessment-1-7-2020-revision.pdf [accessed 5 August 2020].
- 702 NEFSC. 2020b. Butterfish 2020 assessment update report. U.S. Dept. Commer., NOAA, NMFS, NEFSC,
- 703 Woods Hole, MA.
- Nielsen, A., and Berg, C.W. 2014. Estimation of time-varying selectivity in stock assessments using state-space
- models. Fisheries Research **158**: 96–101. doi:10.1016/j.fishres.2014.01.014.
- 706 O'Leary, C.A., Miller, T.J., Thorson, J.T., and Nye, J.A. 2019. Understanding historical summer flounder (
- 707 Paralichthys Dentatus) abundance patterns through the incorporation of oceanography-dependent vital rates

- in Bayesian hierarchical models. Can. J. Fish. Aquat. Sci. **76**(8): 1275–1294. doi:10.1139/cjfas-2018-0092.
- Patrick, W.S., and Link, J.S. 2015. Myths that Continue to Impede Progress in Ecosystem-Based Fisheries
- 710 Management. Fisheries **40**(4): 155–160. doi:10.1080/03632415.2015.1024308.
- Perreault, A.M., Zheng, N., and Cadigan, N.G. 2020. Estimation of growth parameters based on length-
- 712 stratified age samples. Can. J. Fish. Aquat. Sci. **77**(3): 439–450. doi:10.1139/cjfas-2019-0129.
- Perretti, C.T., Deroba, J.J., and Legault, C.M. 2020. Simulation testing methods for estimating misreported
- catch in a state-space stock assessment model. ICES J Mar Sci 77(3): 911–920. Oxford Academic.
- ⁷¹⁵ doi:10.1093/icesjms/fsaa034.
- Perretti, C.T., Fogarty, M.J., Friedland, K.D., Hare, J.A., Lucey, S.M., McBride, R.S., Miller, T.J., Morse,
- R.E., O'Brien, L., Pereira, J.J., Smith, L.A., and Wuenschel, M.J. 2017. Regime shifts in fish recruitment on
- the Northeast US Continental Shelf. Marine Ecology Progress Series 574: 1–11. doi:10.3354/meps12183.
- Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Bris, A.L., Mills, K.E., Nye, J.A., Record,
- 720 N.R., Scannell, H.A., Scott, J.D., Sherwood, G.D., and Thomas, A.C. 2015. Slow adaptation in the
- face of rapid warming leads to collapse of the Gulf of Maine cod fishery. Science 350(6262): 809-812.
- ⁷²² doi:10.1126/science.aac9819.
- Punt, A.E., A'mar, T., Bond, N.A., Butterworth, D.S., de Moor, C.L., De Oliveira, J.A.A., Haltuch, M.A.,
- Hollowed, A.B., and Szuwalski, C. 2014. Fisheries management under climate and environmental uncertainty:
- Control rules and performance simulation. ICES J Mar Sci 71(8): 2208–2220. doi:10.1093/icesjms/fst057.
- R Core Team. 2020. R: A Language and Environment for Statistical Computing. R Foundation for Statistical
- Computing, Vienna, Austria. Available from https://www.R-project.org.
- ⁷²⁸ Rose, G.A., and Rowe, S. 2015. Northern cod comeback. Can. J. Fish. Aquat. Sci. **72**(12): 1789–1798.
- 729 doi:10.1139/cjfas-2015-0346.
- 750 Schnute, J.T., and Haigh, R. 2007. Compositional analysis of catch curve data, with an application to
- 731 Sebastes maliger. ICES Journal of Marine Science 64(2): 218–233. doi:10.1093/icesjms/fsl024.
- Schnute, J.T., and Richards, L.J. 1995. The influence of error on population estimates from catch-age models.
- ⁷³³ Can. J. Fish. Aquat. Sci. **52**(10): 2063–2077. doi:10.1139/f95-800.
- Shelton, P.A., Sinclair, A.F., Chouinard, G.A., Mohn, R., and Duplisea, D.E. 2006. Fishing under low
- 735 productivity conditions is further delaying recovery of Northwest Atlantic cod (Gadus morhua). Can. J. Fish.
- 736 Aquat. Sci. **63**(2): 235–238. doi:10.1139/f05-253.

- 737 Shotwell, S.K., Hanselman, D.H., and Belkin, I.M. 2014. Toward biophysical synergy: Investigating advection
- along the Polar Front to identify factors influencing Alaska sablefish recruitment. Deep Sea Research Part II:
- ⁷³⁹ Topical Studies in Oceanography **107**: 40–53. doi:10.1016/j.dsr2.2012.08.024.
- 5tock, B.C., Xu, H., Miller, T.J., Thorson, J.T., and Nye, J.A. 2021. Implementing two-dimensional autocor-
- relation in either survival or natural mortality improves a state-space assessment model for Southern New
- Fisheries Research 237: 105873. doi:10.1016/j.fishres.2021.105873.
- Stock, C.A., Alexander, M.A., Bond, N.A., Brander, K.M., Cheung, W.W., Curchitser, E.N., Delworth, T.L.,
- Dunne, J.P., Griffies, S.M., Haltuch, M.A., Hare, J.A., Hollowed, A.B., Lehodey, P., Levin, S.A., Link, J.S.,
- Rose, K.A., Rykaczewski, R.R., Sarmiento, J.L., Stouffer, R.J., Schwing, F.B., Vecchi, G.A., and Werner,
- 746 F.E. 2011. On the use of IPCC-class models to assess the impact of climate on Living Marine Resources.
- Progress in Oceanography 88(1-4): 1-27. doi:10.1016/j.pocean.2010.09.001.
- Sullivan, P.J. 1992. A Kalman filter approach to catch-at-length analysis. Biometrics 48(1): 237–257.
- Tableau, A., Collie, J.S., Bell, R.J., and Minto, C. 2018. Decadal changes in the productivity of New England
- ₇₅₀ fish populations. Can. J. Fish. Aquat. Sci. **76**(9): 1528–1540. doi:10.1139/cjfas-2018-0255.
- Thompson, W.R. 1936. On Confidence Ranges for the Median and Other Expectation Distributions for Pop-
- ₇₅₂ ulations of Unknown Distribution Form. Ann. Math. Statist. **7**(3): 122–128. doi:10.1214/aoms/1177732502.
- 753 Thorson, J.T. 2019. Perspective: Let's simplify stock assessment by replacing tuning algorithms with statistics.
- Fisheries Research **217**: 133–139. doi:10.1016/j.fishres.2018.02.005.
- Tommasi, D., Stock, C.A., Hobday, A.J., Methot, R., Kaplan, I.C., Eveson, J.P., Holsman, K., Miller,
- T.J., Gaichas, S., Gehlen, M., Pershing, A., Vecchi, G.A., Msadek, R., Delworth, T., Eakin, C.M., Haltuch,
- ⁷⁵⁷ M.A., Séférian, R., Spillman, C.M., Hartog, J.R., Siedlecki, S., Samhouri, J.F., Muhling, B., Asch, R.G.,
- 758 Pinsky, M.L., Saba, V.S., Kapnick, S.B., Gaitan, C.F., Rykaczewski, R.R., Alexander, M.A., Xue, Y.,
- Pegion, K.V., Lynch, P., Payne, M.R., Kristiansen, T., Lehodey, P., and Werner, F.E. 2017. Managing living
- marine resources in a dynamic environment: The role of seasonal to decadal climate forecasts. Progress in
- Oceanography **152**: 15–49. doi:10.1016/j.pocean.2016.12.011.
- ₇₆₂ Walters, C.J., and Collie, J.S. 1988. Is Research on Environmental Factors Useful to Fisheries Management?
- ⁷⁶³ Can. J. Fish. Aquat. Sci. **45**(10): 1848–1854. doi:10.1139/f88-217.
- Williams, E.H., and Shertzer, K.W. 2015. Technical documentation of the Beaufort Assessment Model
- ₇₆₅ (BAM). U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SEFSC-671. Available from
- http://dx.doi.org/10.7289/V57M05W6 [accessed 21 January 2021].

- Winemiller, K.O., and Rose, K.A. 1992. Patterns of Life-History Diversification in North American Fishes:
- ₇₆₈ Implications for Population Regulation. Canadian Journal of Fisheries and Aquatic Sciences 49(10): 2196–
- 769 2218. doi:10.1139/f92-242.
- 770 Xu, H., Miller, T.J., Hameed, S., Alade, L.A., and Nye, J.A. 2018. Evaluating the utility of the Gulf
- 771 Stream Index for predicting recruitment of Southern New England-Mid Atlantic yellowtail flounder. Fisheries
- Oceanography **27**(1): 85–95. doi:10.1111/fog.12236.
- 773 Xu, H., Thorson, J.T., and Methot, R.D. 2020. Comparing the performance of three data weighting methods
- when allowing for time-varying selectivity. Can. J. Fish. Aquat. Sci. 77(2): 247–263. doi:10.1139/cjfas-2019-
- 775 0107.
- 776 Xu, H., Thorson, J.T., Methot, R.D., and Taylor, I.G. 2019. A new semi-parametric method for autocorrelated
- age- and time-varying selectivity in age-structured assessment models. Can. J. Fish. Aquat. Sci. 76(2):
- ⁷⁷⁸ 268–285. doi:10.1139/cjfas-2017-0446.
- Zwolinski, J.P., and Demer, D.A. 2012. A cold oceanographic regime with high exploitation rates in the
- 780 Northeast Pacific forecasts a collapse of the sardine stock. Proceedings of the National Academy of Sciences
- 781 **109**(11): 4175–4180. doi:10.1073/pnas.1113806109.

782 Appendix A

Table A1 lists the stock-recruit functions, f, available in WHAM, along with possible effects of a modeled environmental covariate, X_y . Note that the user must specify the lag, i.e. number of time steps, ψ , at which X_y affects recruitment (where $\psi = 1$ is common and indicates that X_y affects recruitment the following year). Then

$$\log R_y \sim \mathcal{N}\left(\log\left[f(SSB_{y-1}, X_{y-\psi})\right] - \frac{\sigma_R^2}{2}, \sigma_R^2\right)$$

where SSB_{y-1} is the spawning stock biomass in year y-1. In Table A1, we have abbreviated SSB_{y-1} to S and $X_{y-\psi}$ to X. Each of the environmental effect terms, $e^{\gamma X}$, can be extended to include polynomials as in Eqn. 6, i.e. $e^{\gamma_1 X + \gamma_2 X^2 + \cdots}$.

Table A1: Stock-recruit functions available in WHAM with possible environmental effects.

	Environmental link							
Recruit model	None	Controlling	Limiting	Masking				
Random walk	R_{y-1}							
Random about mean	R_0	$e^{\gamma X}R_0$						
Beverton-Holt	$\frac{\alpha S}{1+\beta S}$	$\frac{\alpha S e^{\gamma X}}{1+\beta S}$	$\frac{\alpha S}{1+\beta Se^{\gamma X}}$	$\frac{\alpha S}{e^{\gamma X} + \beta S}$				
Ricker	$\alpha S e^{-\beta S}$	$\alpha S e^{-\beta S + \gamma X}$		$\alpha S e^{-\beta S(1+\gamma X)}$				

The environment-recruitment link terminology follows Fry (1971) and Iles and Beverton (1998):

- "controlling": density-independent mortality,
- "limiting": carrying capacity effect, e.g. X_y determines the amount of suitable habitat,
- "lethal": threshold effect, i.e. R_y goes to 0 at some X_y value,
- "masking": X_y decreases dR/dSSB, as expected if X_y affects metabolism or growth, and
- "directive": e.g. behavioral.

796 Appendix B

Below we list the available age composition likelihood equations currently implemented in WHAM. For each dataset d and year y, the predicted proportion at age vector is $\hat{\mathbf{p}}_{d,y}$ and the observed proportion at age vector is $\mathbf{p}_{d,y}^*$. The effective sample sizes, Neff_{d,y}, are treated as fixed. The number of non-zero age classes, $n_{d,y}$, youngest non-zero age class, $a_{d,y}^*$, and oldest non-zero age class, $A_{d,y}^*$, also vary by d and y. In the equations below, all d and y have been dropped for simplicity, i.e. they represent the likelihood of a single observed $\mathbf{p}_{d,y}^*$ for a given dataset and year.

803 The predicted proportion vector is

$$\hat{p}_{d,y,a} = \frac{\hat{C}_{a,y,d}}{\sum_{a=1}^{A} \hat{C}_{a,y,d}}$$

if the dataset is catch, and

$$\hat{p}_{d,y,a} = \frac{\hat{I}_{a,y,d}}{\sum_{a=1}^{A} \hat{I}_{a,y,d}}$$

if the dataset is a survey, where $\hat{C}_{a,y,d}$ and $\hat{I}_{a,y,d}$ are defined in Eqns. 7 and 9.

806 Multinomial

$$-\log \mathcal{L}(\hat{\mathbf{p}}|\mathbf{p}^*, \text{Neff}) = \Gamma(\text{Neff} + 1) - \sum_{a} \Gamma(p_a^* \text{Neff} + 1) + \sum_{a} p_a^* \text{Neff} \cdot \log \hat{p}_a$$

with no estimated parameters (age_comp = "multinomial").

808 Dirichlet-Multinomial

$$-\log\mathcal{L}(\hat{\mathbf{p}},\phi|\mathbf{p}^*,\mathrm{Neff}) = \Gamma(\mathrm{Neff}+1) + \Gamma(\phi) - \Gamma(\mathrm{Neff}+\phi) - \sum_a \left[\Gamma(p_a^*\mathrm{Neff}+1) - \Gamma(p_a^*\mathrm{Neff}+\phi\hat{p}_a) + \Gamma(\phi\hat{p}_a)\right]$$

where $\phi > 0$ is an estimated parameter (age_comp = "dir-mult").

10 Dirichlet

$$-\log \mathcal{L}(\hat{\mathbf{p}}, \phi | \mathbf{p}^*) = \Gamma(\phi) - \sum_{a} \left[\Gamma(\phi \hat{p}_a) + (\phi \hat{p}_a - 1) \log p_a^* \right]$$

where $\phi > 0$ is an estimated parameter (age_comp = "dirichlet").

812 Logistic normal

WHAM implements the logistic normal as in Miller et al. (2016). In a given year y and dataset (fishery or survey) d, the negative log-likelihood of the predicted proportion at age vector, $\hat{\mathbf{p}}_{d,y}$, and observation error variance, τ_d^2 , given the observed proportion at age vector, $\mathbf{p}_{d,y}^*$, is

$$v_{d,y} = \frac{\tau_d^2}{\text{Neff}_{d,y}} - \log \mathcal{L}(\hat{\mathbf{p}}, v | \mathbf{p}^*) = \frac{n-1}{2} \log (2\pi v) + \frac{1}{2v} \sum_{a=a^*}^{n-1} \left(\log \frac{p_a^*}{p_{A^*}^*} - \log \frac{\hat{p}_a}{\hat{p}_{A^*}} \right)^2$$

where n, a^* , and A^* are the number of non-zero age classes, and the youngest and oldest non-zero age classes in dataset d and year y (subscripts d and y dropped for simplicity). Neff_{d,y} is an effective sample size that can scale the estimated observation error variance, τ_d^2 , for dataset d based on changes in sampling effort in time. Zero observations can either be pooled with adjacent non-zero age classes (age_comp = "logistic-normal-pool0") or treated as missing (age_comp = "logistic-normal-miss0").

Table 1: Model descriptions and estimated parameters. Parameter descriptions and equations are given in text. Note that the base model in the M module is NAA-1, and the base model in the Selectivity and Ecov-Recruitment modules is NAA-3. Ecov-1 fits the Cold Pool Index data and estimates σ_x in order to allow comparison to Ecov-2 through Ecov-5 using AIC (same data needed in likelihood).

Model	Description	Parameters estimated	No.					
Numbers-at-age (NAA)								
Base	Recruitment deviations are fixed effects	R_y for $y > 1$	$n_{years} - 1$					
NAA-1: Indep. R_y	Recruitment deviations are independent random effects	σ_R	1					
NAA-2: AR(1) R_y	Recruitment deviations are autocorrelated, AR(1), random effects	σ_R, ho_{year}	2					
NAA-3: Indep. $N_{a,y}$	All NAA deviations are independent random effects	σ_R,σ_a	2					
NAA-4: 2D AR(1) $N_{a,y}$	All NAA deviations are random effects with correlation by year and age, 2D AR(1)	$\sigma_R,\sigma_a, ho_{year}, ho_{age}$	4					
Natural mortality (M)								
M-1: None	No random effects on M	σ_R	1					
M-2: Indep.	M deviations are independent random effects	σ_R,σ_M	2					
M-3: $2D AR(1)$	M deviations are random effects with correlation by year and age, 2D AR(1)	$\sigma_R,\sigma_M,arphi_{year},arphi_{age}$	4					
Selectivity (Sel)								
Sel-1: None	No random effects on selectivity	σ_R,σ_a	2					
Sel-2: Indep.	Selectivity deviations are independent random effects	$\sigma_R,\sigma_a,\sigma_{Sel}$	3					
Sel-3: 2D AR(1)	Selectivity deviations are random effects with correlation by year and parameter	$\sigma_R, \sigma_a, \sigma_{Sel}, \phi_{year}, \phi_{par}$	5					
Ecov-Recruitment (Ecov)								
Ecov-1: RW-none	Ecov: random walk (RW), effect on β : none	$\sigma_R,\sigma_a,\sigma_x$	3					
Ecov-2: RW-linear	Ecov: random walk (RW), effect on β : linear	$\sigma_R,\sigma_a,\sigma_x,\beta_1$	4					
Ecov-3: RW-poly	Ecov: random walk (RW), effect on β : 2nd order polynomial (poly)	$\sigma_R,\sigma_a,\sigma_x,eta_1,eta_2$	5					
Ecov-4: $AR(1)$ -linear	Ecov: autocorrelated, AR(1), effect on β : linear	$\sigma_R,\sigma_a,\sigma_x,\phi_x,\beta_1$	5					
Ecov-5: AR(1)-poly	Ecov: autocorrelated, AR(1), effect on β : 2nd order polynomial (poly)	$\sigma_R, \sigma_a, \sigma_x, \phi_x, \beta_1, \beta_2$	6					

35

Table 2: Stocks used in simulation tests.

	Processes tested		Model dim		Biol. par.		Stock status				
Stock	NAA	M	Sel	Ecov	# Ages	# Years	\overline{M}	σ_R	$\frac{B}{B_{40}}$	$\frac{F}{F_{40}}$	Source
SNEMA yellowtail flounder	X	X		x	6	49	0.2-0.4	1.67	0.01	0.44	NEFSC (2020a)
Butterfish	X	\mathbf{X}			5	31	1.3	0.23	2.57	0.03	NEFSC (2020b)
North Sea cod	X	X			6	54	0.2 - 1.2	0.87	0.14	2.00	ICES (2017a)
Icelandic herring	X				11	30	0.1	0.55	0.40	1.81	ICES $(2017b)$
Georges Bank haddock	X		X		9	86	0.2	1.65	5.16	0.12	NEFSC $(2020a)$




Figure 1: AIC differences by model and stock when fit to original datasets. Random effects were only used for one process at a time (numbers at age, natural mortality, selectivity, and recruitment linked to the Cold Pool Index, CPI). Stock abbreviations: SNEMA yellowtail flounder (SNEMAYT), North Sea cod (NScod), Icelandic herring (ICEherring), and Georges Bank haddock (GBhaddock).




Figure 2: Change in Mohn's ρ relative to the statistical catch at age model (Base) for full state-space models (numbers at all ages are random effects, NAA-3 and NAA-4). Changes in Mohn's ρ are shown for all five stocks (whisker and box ends and median line in boxplots). Reduction in Mohn's ρ_R for SNEMA yellowtail flounder NAA-4 is off chart.




Figure 3: Mean change in coefficient of variation (CV) for key quantities in model and projection years for full state-space models (numbers at all ages are random effects, NAA-3 and NAA-4) relative to NAA-1. Boxplots show mean change in CV across years for all five stocks (whisker and box ends, median line). F was fixed at 0 in projection years.




Figure 4: Abundance-at-age deviations estimated for Georges Bank haddock using alternative models of numbers-at-age (NAA) random effects. Base = a traditional statistical catch-at-age (SCAA) model. NAA-1 = only recruitment deviations are independent random effects (most similar to Base). NAA-2 = as NAA-1, but with autocorrelated recruitment deviations, AR(1). NAA-3 = all NAA deviations are independent random effects. NAA-4 = as NAA-3, but deviations are correlated by age ($\rho_{age} = -0.12, 95\%$ CI: -0.32-0.08) and year ($\rho_{year} = 0.73, 95\%$ CI: 0.58-0.87). The vertical dashed line indicates the terminal year in the assessment.




Figure 5: Natural mortality, $\log(M)$, estimated for Southern New England-Mid Atlantic yellowtail flounder using three random effects models. M-1 = no random effects on M. M-2 = independent M deviations. M-3 = M deviations are correlated by age ($\varphi_{age} = 0.40, 95\%$ CI: 0.09-0.70) and year ($\varphi_{year} = 0.63, 95\%$ CI: 0.31-0.94). The vertical dashed line indicates the terminal year in the assessment.




Figure 6: Selectivity estimated for Georges Bank haddock using three random effects models. Sel-1 = no random effects (constant logistic selectivity). Sel-2 = independent selectivity deviations. Sel-3 = selectivity deviations are correlated by parameter ($\phi_{par}=0.60,\,95\%$ CI: 0.30-0.80) and year ($\phi_{year}=0.87,\,95\%$ CI: 0.74-0.94).




Figure 7: Beverton-Holt stock-recruit relationships fit for Southern New England-Mid Atlantic yellowtail flounder, with and without effects of the Cold Pool Index (CPI). A) CPI estimated from the model with lowest AIC (Ecov-4, AR(1)-linear). Points are observations with 95% CI, and the line with shading is the model-estimated CPI with 95% CI. Note the increased uncertainty surrounding the CPI estimate in 2017 (no observation). B) Estimates of spawning stock biomass (SSB), recruitment, and the stock-recruit function from the model without a CPI effect, Ecov-1. C) Estimates of SSB and recruitment from Ecov-4, with an effect of the CPI on β . Lines depict the expected stock-recruit relationship in each year y, given the CPI in year y-1 (color).

Operating model NAA-1 NAA-2 0.50 0.25 SSB 0.00 0 0 -0.25 -0.50 0.50 0.25 0 0 -0.25 -0.50 0.50 Relative error 0.25 B/B_{40%} 0.00 -0.25 -0.50 0.50 0.25 0 F/F_{40%} 0 0 0.00 -0.25 -0.50 0.50 Recruitment 0.25 Ó Ó 0 -0.25 -0.50 **Estimation model**

Figure 8: Relative error of key quantities estimated for Georges Bank haddock using four models of numbers-at-age (NAA) random effects. NAA-1 = only recruitment deviations are random effects (most similar to Base, a traditional statistical catch-at-age model), and deviations are independent. NAA-2 = as NAA-1, but with autocorrelated recruitment deviations. NAA-3 = all NAA deviations are independent random effects. NAA-4 = as NAA-3, but deviations are correlated by age and year. Points without lines indicate that 95% CI are smaller than the points.




Figure 9: Relative error of key quantities estimated for Southern New England-Mid Atlantic yellowtail flounder using three models of natural mortality (M) random effects. M-1 = no random effects on M. M-2 = independent M deviations. M-3 = M deviations are correlated by age and year. Points without lines indicate that 95% CI are smaller than the points.




Figure 10: Relative error of key quantities estimated for Georges Bank haddock using three models of selectivity random effects. Sel-1 = no random effects (constant logistic selectivity). Sel-2 = independent selectivity deviations. Sel-3 = selectivity deviations are correlated by parameter and year. Points without lines indicate that 95% CI are smaller than the points.




Figure 11: Proportion of simulations in which each model had the lowest AIC. A) Numbers-at-age (NAA), aggregated across all five stocks. B) Natural mortality (M), aggregated over two stocks (SNEMAYT and butterfish). C) Selectivity (GBhaddock). Not all estimation models converged for each simulation, even when the operating model matched.