

1 The Woods Hole Assessment Model (WHAM): a general state-space
2 assessment framework that incorporates time- and age-varying
3 processes via random effects and environmental covariates

4 Brian C. Stock¹, Timothy J. Miller ¹

5 ¹brian.stock@noaa.gov, timothy.j.miller@noaa.gov, Northeast Fisheries Science Center, National Marine
6 Fisheries Service, 166 Water Street, Woods Hole, MA 02543, USA

8 **Abstract**

9 WHAM is great.

10 **Keywords**

11 state-space; stock assessment; random effects; time-varying; environmental effects; recruitment; survival;
12 natural mortality; Template Model Builder (TMB)

1 Introduction

The last two decades have increasingly seen a push for more holistic, ecosystem-based fisheries management (Larkin, 1996; Link, 2002). In part, this is a recognition that considering single species in isolation produces riskier and less robust outcomes long-term (Patrick and Link, 2015). In several high-profile cases, fisheries management has failed to prevent collapses because they did not reduce fishing pressure in responses to changes in natural mortality (M), recruitment, or migration patterns caused by dynamics external to the stock in question (Northern cod: Shelton et al., 2006, @rose2015Northern; Gulf of Maine cod: Pershing et al., 2015; Pacific sardine: Zwolinski and Demer, 2012). This is particularly concerning in the context of climate change and the wide range of biological processes—often assumed to be constant—in stock assessments that are likely to be affected (Stock et al., 2011).

One approach to account for changing productivity is to explicitly link population processes to environmental covariates in single-species stock assessments, i.e. the mechanistic approach *sensu* Punt et al. (2014). Traditional single-species assessments are based on internal population dynamics and the effect of fishing mortality (F), and typically ignore effects of the environment or interactions with other species, even though fisheries scientists have long known that these are important drivers of time-varying population processes, e.g. recruitment, mortality, growth, and movement (Garstang, 1900; Hjort, 1914). Despite how counterintuitive this may seem to ecologists and oceanographers who study such relationships, the evidence for direct linkages to specific environmental covariates is often weak and can break down over time (McClatchie et al., 2010; Myers, 1998). Additionally, the primary goal of most assessments is to provide management advice on near-term sustainable harvest levels—not to explain ecological relationships. Even if an environmental covariate directly affects fish productivity, including the effect in an assessment may not improve management advice if the effect is weak (De Oliveira and Butterworth, 2005). Worse, including environmental effects in an assessment or management system has been shown to actually provide worse management in some cases (De Oliveira and Butterworth, 2005; Punt et al., 2014; Walters and Collie, 1988). This can be true even in cases of relatively well-understood mechanistic links between oceanic conditions and fish populations, as in the case of sea surface temperature and Pacific sardine (Hill et al., 2018; Zwolinski and Demer, 2012).

An alternative approach is to allow biological parameters to vary stochastically over time, without explanation, i.e. the empirical approach *sensu* Punt et al. (2014). Statistical catch-at-age (SCAA) models typically only estimate year-specific recruitment (R_t) and F_t , often as deviations from a mean, e.g. $\log R_t = \log R_0 + \epsilon_t$. The main reason that other parameters are assumed constant is simply that there are not enough degrees of freedom to estimate many time-varying parameters. One common solution is to penalize the deviations,

e.g. $\epsilon_t \sim \mathcal{N}(0, \sigma_\epsilon^2)$, although the penalty terms, σ_ϵ^2 , must be fixed and are therefore subjective (Methot and Wetzel, 2013; Xu et al., 2019). State-space models that treat parameters as unobserved states can avoid such subjectivity by estimating the penalty terms as variance parameters constraining random effects (Aeberhard et al., 2018). In this way, state-space models can allow processes to vary in time while simultaneously estimating fewer parameters.

Although state-space stock assessments have existed for some time (Gudmundsson, 1994; Mendelsohn, 1988; Sullivan, 1992), the recent development of Template Model Builder (TMB, Kristensen et al., 2016) software to perform efficient Laplace approximation has greatly expanded their use (Cadigan, 2016; Miller et al., 2016; Nielsen and Berg, 2014). In addition to the key advantage of objectively estimating variance, or “data weighting”, parameters, state-space models naturally predict unobserved states, and therefore handle missing data and short-term projections in a straightforward way (ICES, 2020). In comparisons with SCAA models, they generally have larger (more realistic) uncertainty and lower retrospective bias (Miller et al. in prep).

Retrospective bias can occur when changing environmental conditions lead to changes in productivity that are unaccounted for in stock assessments, and this is a concern common to several groundfish stocks on the Northeast U.S. Shelf (Brooks and Legault, 2016). The Northeast U.S. Shelf ecosystem is rapidly changing, and this has motivated managers to make the “continue[d] development of stock assessment models that include environmental terms” a top priority (Hare et al., 2016). In addition to providing short-term (1-3 years) catch advice with reduced retrospective bias, it is hoped that environment-linked assessments will help create realistic rebuilding plans in the medium-term (3-10 years) for stocks that have not responded to dramatic decreases in F . Miller et al. (2016) developed a state-space model for yellowtail flounder with an environmental effect on recruitment, which reduced retrospective patterns and residual variance. Similar applications that include environmental effects on growth, M , and maturity in a state-space framework have also proven promising (Miller and Hyun, 2018; Miller et al., 2018; O’Leary et al., 2019; Xu et al., 2018).

In summary, to address the needs of fisheries management in a changing climate we seek an assessment framework that 1) estimates time-varying parameters as random effects (i.e. a state-space model), and 2) includes environmental effects. The framework should also allow for easy testing against status quo SCAA models to ease gradual adoption through the “research track” or “benchmark” assessment process (Lynch et al., 2018). The objectives of this manuscript are to introduce the Woods Hole Assessment Model (WHAM) framework and demonstrate its ability to:

1. estimate time- and age-varying random effects on survival, M , and selectivity;
2. fit environmental time-series with process and observation error, missing data, and a link to a population

process; and

3. simulate new data and random effects to conduct self- and cross-tests (*sensu* Deroba et al., 2015).

Finally, we describe how the above are implemented using an open-source software package available to the stock assessment community (Miller and Stock, 2020).

2 Methods

2.1 Model description

Similar to ASAP. Empirical weight-at-age.

2.1.1 Modules with random effects

See Table 1.

2.1.1.1 Numbers-at-age (survival)

2.1.1.2 Natural mortality (M)

2.1.1.3 Selectivity

2.1.1.4 Environmental covariate(s)

2.1.1.4.1 Time-series model

2.1.1.4.2 Observation model

2.1.1.4.3 Link to population

2.2 How is WHAM different from SAM?

Not sure where to put this... some could be in Intro or Discussion. Definitely will be a question in readers' minds so may be good to introduce early?

Most assessments in the U.S. assume separability in $F_{a,t}$, estimate F_t and Sel_a . WHAM does this. SAM estimates $F_{a,t}$ directly. WHAM and SAM also make different separability assumptions for the catch/index data (aggregate total + age comps vs. $C_{a,t}$ directly). Should be similar (?) but could test.

Goal is to replicate ASAP assessments in the U.S. Northeast. Can easily turn on/off random effects.

Observation model is natural for landings data that are measured as total weight plus age composition sampling. Age composition sampling often done separately with survey data.

Treating F and Sel separately can be useful for projections. Oftentimes we want to specify F in projections to calculate a reference point, as opposed to continuing a F time-series process.

2.2.1 Data/observation model

2.2.1.1 Catch (agg, age comp)

2.2.1.2 Index (agg, age comp)

2.2.2 Bias correction

- Analytical obs error. (Aldrin et al., 2020).
- Analytical process error.
- TMB epsilon. (Thorson, 2019; Thorson and Kristensen, 2016)

Should these all be used?

2.3 Simulation tests

Fit each model to original dataset. Use each model to simulate new data and random effects, keeping fixed effect parameters constant at values estimated in original fits. Re-fit each model to datasets simulated under each operating model. We used the stocks in Table 2.

We used R (R Core Team, 2020). WHAM is available as an R package (Miller and Stock, 2020).

3 Results

3.1 Original datasets

2D AR1 structure performed well across modules and stocks (Fig. 1).

3.1.1 Numbers-at-age

Fig. 2.

3.1.2 Natural mortality

Fig. 3.

3.1.3 Selectivity

Fig. 4.

3.1.4 Ecov-Recruitment

Fig. 5.

3.2 Simulation tests

Not all models converged. Fig. 6.

3.2.1 Numbers-at-age

Fig. 7.

Fig. 8.

3.2.2 Natural mortality

Fig. 9.

Fig. 10.

3.2.3 Selectivity

Fig. 11.

Fig. 12.

3.2.4 Ecov-Recruitment

Fig. 13.

4 Discussion

4.1 Overview

We described WHAM. Sim tests showed no bias in self-tests (when estimation model matched operating model). Some bias in cross-tests.

4.2 Future work

WHAM will be used in upcoming research track assessments. Could transition to operational. Potential to improve several NEFSC assessments.

- 2D AR(1) selectivity. Most assessments in the U.S. assume separability in $F_{a,t}$, i.e. estimate F_t and Sel_a . WHAM does this. SAM estimates $F_{a,t}$ directly. WHAM and SAM make different separability assumptions for the catch/index data as well (aggregate total + age comps vs. $C_{a,t}$ directly). Should be similar (?) but could test.
- How many time/age-varying random effects can be estimated simultaneously? Stock et al. (n.d.) estimated random effect deviations in survival and M , as well as an environmental covariate effect on recruitment.
- Ecov-Recruitment simulation study. How much information does Ecov need to have to be useful?

4.3 Extensions

4.3.1 Multivariate spatiotemporal environmental data

4.3.2 Length/growth estimation

4.3.3 Ecov models

- AR(k)
- splines
- Gaussian process/EDM/Munch/Sugihara

4.4 Conclusion

Development of TMB has facilitated significant advancement in fisheries assessment, allowing us to treat population processes as random effects. A grand challenge in fisheries is to assess and manage stocks in a changing environment. Increasingly have the environmental data. Population time-series are lengthening. WHAM is a step in this direction.

Acknowledgements

This research was performed while BCS held an NRC Research Associateship award at the NEFSC. NOAA Fish & Climate grant number ??.

¹⁶⁹ **Supplementary material**

¹⁷⁰ More figures.

References

- Aeberhard, W.H., Mills Flemming, J., Nielsen, A., 2018. Review of State-Space Models for Fisheries Science. *Annu. Rev. Stat. Appl.* 5, 215–235. <https://doi.org/10.1146/annurev-statistics-031017-100427>
- Aldrin, M., Tvete, I., Aanes, S., Subbey, S., 2020. The specification of the data model part in the SAM model matters. *Fisheries Research* 229, 105585. <https://doi.org/10.1016/j.fishres.2020.105585>
- Brooks, E.N., Legault, C.M., 2016. Retrospective forecasting evaluating performance of stock projections for New England groundfish stocks. *Can. J. Fish. Aquat. Sci.* 73, 935–950. <https://doi.org/10.1139/cjfas-2015-0163>
- Cadigan, N.G., 2016. A state-space stock assessment model for northern cod, including under-reported catches and variable natural mortality rates. *Canadian Journal of Fisheries and Aquatic Sciences* 73, 296–308. <https://doi.org/10.1139/cjfas-2015-0047>
- De Oliveira, J., Butterworth, D., 2005. Limits to the use of environmental indices to reduce risk and/or increase yield in the South African anchovy fishery. *African Journal of Marine Science* 27, 191–203. <https://doi.org/10.2989/18142320509504078>
- Deroba, J.J., Butterworth, D.S., Methot, R.D., De Oliveira, J.a.A., Fernandez, C., Nielsen, A., Cadrin, S.X., Dickey-Collas, M., Legault, C.M., Ianelli, J., Valero, J.L., Needle, C.L., O'Malley, J.M., Chang, Y.-J., Thompson, G.G., Canales, C., Swain, D.P., Miller, D.C.M., Hintzen, N.T., Bertignac, M., Ibaibarriaga, L., Silva, A., Murta, A., Kell, L.T., de Moor, C.L., Parma, A.M., Dichmont, C.M., Restrepo, V.R., Ye, Y., Jardim, E., Spencer, P.D., Hanselman, D.H., Blaylock, J., Mood, M., Hulson, P.-J.F., 2015. Simulation testing the robustness of stock assessment models to error: Some results from the ICES strategic initiative on stock assessment methods. *ICES J Mar Sci* 72, 19–30. <https://doi.org/10.1093/icesjms/fst237>
- Garstang, W., 1900. The Impoverishment of the Sea. A Critical Summary of the Experimental and Statistical Evidence bearing upon the Alleged Depletion of the Trawling Grounds. *Journal of the Marine Biological Association of the United Kingdom* 6, 1–69. <https://doi.org/10.1017/S0025315400072374>
- Gudmundsson, G., 1994. Time series analysis of catch-at-age observations. *Applied Statistics* 43, 117–126.
- Hare, J.A., Borggaard, D.L., Friedland, K.D., Anderson, J., Burns, P., Chu, K., Clay, P.M., Collins, M.J., Cooper, P., Fratantoni, P.S., Johnson, M.R., Manderson, J.P., Milke, L., Miller, T.J., Orphanides, C.D., Saba, V.S., 2016. Northeast Regional Action Plan - NOAA Fisheries Climate Science Strategy (No. NMFS-NE-239). NOAA Fisheries, Northeast Fisheries Science Center, Woods Hole, MA.

- Hill, K.T., Crone, P.R., Zwolinski, J.P., 2018. Assessment of the Pacific sardine resource in 2018 for U.S. Management in 2018-2019 (No. NOAA Technical Memorandum NMFS-SWFSC-600). US Department of Commerce.
- Hjort, J., 1914. Fluctuations in the great fisheries of Northern Europe viewed in the light of biological research. *Rapports et Procès-Verbaux des Réunions du Conseil Permanent International Pour L'Exploration de la Mer* 20, 1–228.
- ICES, 2020. Workshop on the Review and Future of State Space Stock Assessment Models in ICES (WKRFSA). ICES Scientific Reports 2, 23p. <https://doi.org/10.17895/ices.pub.6004>
- Kristensen, K., Nielsen, A., Berg, C., Skaug, H., Bell, B.M., 2016. TMB: Automatic differentiation and Laplace approximation. *Journal of Statistical Software* 70, 1–21. <https://doi.org/10.18637/jss.v070.i05>
- Larkin, P., 1996. Concepts and issues in marine ecosystem management. *Reviews in Fish Biology and Fisheries* 6, 139–164. <https://doi.org/10.1007/BF00182341>
- Link, J.S., 2002. What Does Ecosystem-Based Fisheries Management Mean? *Fisheries* 27, 5.
- Lynch, P.D., Methot, R.D., Link, J.S. (Eds.), 2018. Implementing a Next Generation Stock Assessment Enterprise. An Update to the NOAA Fisheries Stock Assessment Improvement Plan, in: U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/ SPO-183, p. 127. <https://doi.org/10.7755/TMSPO.183>
- McClatchie, S., Goericke, R., Auad, G., Hill, K., 2010. Re-assessment of the stockRecruit and temperatureRecruit relationships for Pacific sardine (*Sardinops sagax*). *Can. J. Fish. Aquat. Sci.* 67, 1782–1790. <https://doi.org/10.1139/F10-101>
- Mendelssohn, R., 1988. Some problems in estimating population sizes from catch-at-age data. *Fishery Bulletin* 86, 617–630.
- Methot, R.D., Wetzel, C.R., 2013. Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research* 142, 86–99. <https://doi.org/10.1016/j.fishres.2012.10.012>
- Miller, T.J., Hare, J.A., Alade, L.A., 2016. A state-space approach to incorporating environmental effects on recruitment in an age-structured assessment model with an application to southern New England yellowtail flounder. *Canadian Journal of Fisheries and Aquatic Sciences* 73, 1261–1270. <https://doi.org/10.1139/cjfas-2015-0339>
- Miller, T.J., Hyun, S.-Y., 2018. Evaluating evidence for alternative natural mortality and process error

assumptions using a state-space, age-structured assessment model. Canadian Journal of Fisheries and Aquatic Sciences 75, 691–703. <https://doi.org/10.1139/cjfas-2017-0035>

Miller, T.J., O'Brien, L., Fratantoni, P.S., 2018. Temporal and environmental variation in growth and maturity and effects on management reference points of Georges Bank Atlantic cod. Can. J. Fish. Aquat. Sci. 1–13. <https://doi.org/10.1139/cjfas-2017-0124>

Miller, T.J., Stock, B.C., 2020. The Woods Hole Assessment Model (WHAM).

Myers, R.A., 1998. When do environment-recruitment correlations work? Reviews in Fish Biology and Fisheries 8, 285–305.

Nielsen, A., Berg, C.W., 2014. Estimation of time-varying selectivity in stock assessments using state-space models. Fisheries Research 158, 96–101. <https://doi.org/10.1016/j.fishres.2014.01.014>

O'Leary, C.A., Miller, T.J., Thorson, J.T., Nye, J.A., 2019. Understanding historical summer flounder (*Paralichthys Dentatus*) abundance patterns through the incorporation of oceanography-dependent vital rates in Bayesian hierarchical models. Can. J. Fish. Aquat. Sci. 76, 1275–1294. <https://doi.org/10.1139/cjfas-2018-0092>

Patrick, W.S., Link, J.S., 2015. Myths that Continue to Impede Progress in Ecosystem-Based Fisheries Management. Fisheries 40, 155–160. <https://doi.org/10.1080/03632415.2015.1024308>

Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Bris, A.L., Mills, K.E., Nye, J.A., Record, N.R., Scannell, H.A., Scott, J.D., Sherwood, G.D., Thomas, A.C., 2015. Slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery. Science 350, 809–812. <https://doi.org/10.1126/science.aac9819>

Punt, A.E., A'mar, T., Bond, N.A., Butterworth, D.S., de Moor, C.L., De Oliveira, J.A.A., Haltuch, M.A., Hollowed, A.B., Szuwalski, C., 2014. Fisheries management under climate and environmental uncertainty: Control rules and performance simulation. ICES J Mar Sci 71, 2208–2220. <https://doi.org/10.1093/icesjms/fst057>

R Core Team, 2020. R: A Language and Environment for Statistical Computing.

Rose, G.A., Rowe, S., 2015. Northern cod comeback. Can. J. Fish. Aquat. Sci. 72, 1789–1798. <https://doi.org/10.1139/cjfas-2015-0346>

Shelton, P.A., Sinclair, A.F., Chouinard, G.A., Mohn, R., Duplisea, D.E., 2006. Fishing under low productivity conditions is further delaying recovery of Northwest Atlantic cod (*Gadus morhua*). Can. J. Fish. Aquat. Sci.

258 63, 235–238. <https://doi.org/10.1139/f05-253>

259 Stock, B.C., Xu, H., Miller, T.J., Thorson, J.T., Nye, J.A., n.d. Implementing a 2-dimensional smoother on
 260 either survival or natural mortality improves a state-space assessment model for Southern New England-Mid
 261 Atlantic yellowtail flounder.

262 Stock, C.A., Alexander, M.A., Bond, N.A., Brander, K.M., Cheung, W.W., Curchitser, E.N., Delworth, T.L.,
 263 Dunne, J.P., Griffies, S.M., Haltuch, M.A., Hare, J.A., Hollowed, A.B., Lehodey, P., Levin, S.A., Link, J.S.,
 264 Rose, K.A., Rykaczewski, R.R., Sarmiento, J.L., Stouffer, R.J., Schwing, F.B., Vecchi, G.A., Werner, F.E.,
 265 2011. On the use of IPCC-class models to assess the impact of climate on Living Marine Resources. *Progress*
 266 *in Oceanography* 88, 1–27. <https://doi.org/10.1016/j.pocean.2010.09.001>

267 Sullivan, P.J., 1992. A Kalman filter approach to catch-at-length analysis. *Biometrics* 48, 237–257.

268 Thorson, J.T., 2019. Perspective: Let’s simplify stock assessment by replacing tuning algorithms with
 269 statistics. *Fisheries Research* 217, 133–139. <https://doi.org/10.1016/j.fishres.2018.02.005>

270 Thorson, J.T., Kristensen, K., 2016. Implementing a generic method for bias correction in statistical
 271 models using random effects, with spatial and population dynamics examples. *Fisheries Research* 175, 66–74.
 272 <https://doi.org/10.1016/j.fishres.2015.11.016>

273 Walters, C.J., Collie, J.S., 1988. Is Research on Environmental Factors Useful to Fisheries Management?
 274 *Can. J. Fish. Aquat. Sci.* 45, 1848–1854. <https://doi.org/10.1139/f88-217>

275 Xu, H., Miller, T.J., Hameed, S., Alade, L.A., Nye, J.A., 2018. Evaluating the utility of the Gulf Stream Index
 276 for predicting recruitment of Southern New England-Mid Atlantic yellowtail flounder. *Fisheries Oceanography*
 277 27, 85–95. <https://doi.org/10.1111/fog.12236>

278 Xu, H., Thorson, J.T., Methot, R.D., 2019. Comparing the performance of three data-weighting methods when
 279 allowing for time-varying selectivity. *Can. J. Fish. Aquat. Sci.* 1–17. <https://doi.org/10.1139/cjfas-2019-0107>

280 Zwolinski, J.P., Demer, D.A., 2012. A cold oceanographic regime with high exploitation rates in the Northeast
 281 Pacific forecasts a collapse of the sardine stock. *Proceedings of the National Academy of Sciences* 109,
 282 4175–4180. <https://doi.org/10.1073/pnas.1113806109>

Table 1: Model descriptions and estimated parameters. Parameter descriptions and equations are given in text. Note that the base model in the M module is NAA m1, and the base model in the Selectivity and Ecov-Recruitment modules is NAA m3. Ecov m1 fits the Cold Pool Index data and estimates σ_x in order to allow comparison to m2-m5 using AIC (same data needed in likelihood).

Model	Description	Estimated parameters
Numbers-at-age (NAA)		
m1: SCAA (IID)	Recruitment deviations are IID random effects	σ_R
m2: SCAA (AR1)	Recruitment deviations are autocorrelated (AR1) random effects	σ_R, ρ_y
m3: NAA (IID)	All NAA deviations are IID random effects	σ_R, σ_a
m4: NAA (2D AR1)	All NAA deviations are random effects with correlation by year and age (2D AR1)	$\sigma_R, \sigma_a, \rho_y, \rho_a$
Natural mortality (M)		
m1: none	No random effects on M	σ_R
m2: IID	M deviations are IID random effects	σ_R, σ_M
m3: 2D AR1	M deviations are random effects with correlation by year and age (2D AR1)	$\sigma_R, \sigma_M, \varphi_y, \varphi_a$
Selectivity (Sel)		
m1: none	No random effects on selectivity	σ_R, σ_a
m2: IID	Selectivity deviations are IID random effects	$\sigma_R, \sigma_a, \sigma_{Sel}$
m3: 2D AR1	Selectivity deviations are random effects with correlation by year and age (2D AR1)	$\sigma_R, \sigma_a, \sigma_{Sel}, \phi_y, \phi_a$
Ecov-Recruitment (Ecov)		
m1: RW-none	Ecov: random walk (RW), effect on β : none	$\sigma_R, \sigma_a, \sigma_x$
m2: RW-linear	Ecov: random walk (RW), effect on β : linear	$\sigma_R, \sigma_a, \sigma_x, \beta_1$
m3: RW-poly	Ecov: random walk (RW), effect on β : 2nd order polynomial (poly)	$\sigma_R, \sigma_a, \sigma_x, \beta_1, \beta_2$
m4: AR1-linear	Ecov: autocorrelated (AR1), effect on β : linear	$\sigma_R, \sigma_a, \sigma_x, \phi_x, \beta_1$
m5: AR1-poly	Ecov: autocorrelated (AR1), effect on β : 2nd order polynomial (poly)	$\sigma_R, \sigma_a, \sigma_x, \phi_x, \beta_1, \beta_2$

Table 2: Stocks used in simulation tests.

Stock	Modules tested				Model dim		Biol. par.		Stock status	
	NAA	M	Sel	Ecov	# Ages	# Years	M	σ_R	$\frac{B}{B_{40}}$	$\frac{F}{F_{40}}$
SNEMA yellowtail flounder	x	x		x	6	49	0.2-0.4	1.67	0.01	0.44
Butterfish	x	x			5	31	1.3	0.23	2.57	0.03
North Sea cod	x	x			6	54	0.2-1.2	0.87	0.14	2.00
Icelandic herring	x				11	30	0.1	0.55	0.40	1.81
Georges Bank haddock	x		x		9	86	0.2	1.65	5.16	0.12




Figure 1: AIC differences by model and stock when fit to original datasets. Stock abbreviations: SNEMA yellowtail flounder (SNEMAYT), North Sea cod (NScod), Icelandic herring (ICEherring), and Georges Bank haddock (GBhaddock).




Figure 2: Survival deviations estimated for Icelandic herring using four models of numbers-at-age (NAA) random effects. m1 = only recruitment deviations are random effects (most similar to traditional statistical catch-at-age, SCAA), and deviations are independent and identically distributed (IID). m2 = as m1, but with autocorrelated recruitment deviations (AR1_y). m3 = all NAA deviations are IID random effects. m4 = as m3, but deviations are correlated by age and year (2D AR1).




Figure 3: Natural mortality (M) estimated for butterfish using three random effects models. m1 = no random effects on M . m2 = M deviations are independent and identically distributed (IID). m3 = M deviations are correlated by age and year (2D AR1).




Figure 4: Selectivity estimated for Georges Bank haddock using three random effects models. m1 = no random effects (constant logistic selectivity). m2 = selectivity deviations are independent and identically distributed (IID). m3 = selectivity deviations are correlated by parameter and year (2D AR1).




Figure 5: Beverton-Holt stock-recruit relationships fit for Southern New England-Mid Atlantic yellowtail flounder, with and without effects of the Cold Pool Index (CPI). A) CPI estimated from the model with lowest AIC (m4, AR1-linear). Points are observations with 95% CI, and the line with shading is the model-estimated CPI with 95% CI. Note the increased uncertainty surrounding the CPI estimate in 2017 (no observation). B) Estimates of spawning stock biomass (SSB), recruitment, and the stock-recruit function from the model without a CPI effect, m1. C) Estimates of SSB and recruitment from m4, with an effect of the CPI on β . Lines depict the expected stock-recruit relationship in each year t , given the CPI in year $t - 1$ (color).




Figure 6: Proportion of simulations in which each model had the lowest AIC. A) Numbers-at-age (NAA), aggregated across all five stocks. B) Natural mortality (M), aggregated over two stocks (SNEMAYT and NScod). C) Selectivity (GBhaddock). Not all estimation models converged for each simulation, even when the operating model matched.




Figure 7: Relative error of parameters constraining numbers-at-age (NAA) random effects. Four models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. m1 = only recruitment deviations are random effects (most similar to traditional statistical catch-at-age, SCAA), and deviations are independent and identically distributed (IID). m2 = as m1, but with autocorrelated recruitment deviations (AR1_y). m3 = all NAA deviations are IID random effects. m4 = as m3, but deviations are correlated by age and year (2D AR1). Relative error was calculated as $\frac{\hat{\theta}_i}{\theta} - 1$, where $\hat{\theta}_i$ was the estimate in simulation i for parameter θ , and θ was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI.




Figure 8: Relative error of key quantities estimated for Icelandic herring using four models of numbers-at-age (NAA) random effects. m1 = only recruitment deviations are random effects (most similar to traditional statistical catch-at-age, SCAA), and deviations are independent and identically distributed (IID). m2 = as m1, but with autocorrelated recruitment deviations (AR1_y). m3 = all NAA deviations are IID random effects. m4 = as m3, but deviations are correlated by age and year (2D AR1).




Figure 9: Relative error of parameters constraining natural mortality (M) random effects. Three models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. m1 = no random effects on M . m2 = M deviations were independent and identically distributed (IID). m3 = M deviations were correlated by age and year (2D AR1). Relative error was calculated as $\frac{\hat{\theta}_i}{\theta} - 1$, where $\hat{\theta}_i$ was the estimate in simulation i for parameter θ , and θ was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI. Stock abbreviations: SNEMAYT yellowtail flounder (SNEMAYT) and North Sea cod (NScod, m3 did not converge).




Figure 10: Relative error of key quantities estimated for butterfish using three models of natural mortality (M) random effects. m1 = no random effects on M . m2 = M deviations are independent and identically distributed (IID). m3 = M deviations are correlated by age and year (2D AR1).




Figure 11: Relative error of parameters constraining selectivity random effects for Georges Bank haddock (GBhaddock). Three models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. m1 = no random effects (constant selectivity). m2 = selectivity deviations were independent and identically distributed (IID). m3 = selectivity deviations were correlated by parameter and year (2D AR1). Relative error was calculated as $\frac{\hat{\theta}_i}{\theta} - 1$, where $\hat{\theta}_i$ was the estimate in simulation i for parameter θ , and θ was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI.




Figure 12: Relative error of key quantities estimated for Georges Bank haddock using three models of selectivity random effects. m1 = no random effects (constant logistic selectivity). m2 = selectivity deviations are independent and identically distributed (IID). m3 = selectivity deviations are correlated by parameter and year (2D AR1).




Figure 13: Relative error of parameters constraining variation in recruitment for Southern New England-Mid Atlantic yellowtail flounder (SNEMAYT). Five models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. All models estimated recruitment using the Beverton-Holt function and included CPI effects on β : $\hat{R}_{t+1} = \frac{\alpha S_t}{1 + e^{\beta_0 + \beta_1 x_t + \beta_2 x_t^2 S_t}}$. m1 = Cold Pool Index (CPI) modeled as a random walk (RW) with no effect on recruitment ($\beta_1 = \beta_2 = 0$). m2 = CPI as RW, linear effect on β . m3 = CPI as RW, 2nd order polynomial effect on β . m4 = CPI as AR1, linear effect. m5 = CPI as AR1, polynomial effect. Relative error was calculated as $\frac{\hat{\theta}_i}{\theta} - 1$, where $\hat{\theta}_i$ was the estimate in simulation i for parameter θ , and θ was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI.