

1 The Woods Hole Assessment Model (WHAM): a general state-space  
2 assessment framework that incorporates time- and age-varying  
3 processes via random effects and links to environmental covariates

4 Brian C. Stock<sup>1</sup>, Timothy J. Miller <sup>1</sup>

5 <sup>1</sup>brian.stock@noaa.gov, timothy.j.miller@noaa.gov, Northeast Fisheries Science Center, National Marine  
6 Fisheries Service, 166 Water Street, Woods Hole, MA 02543, USA

## 8 **Abstract**

9 WHAM is great.

## 10 **Keywords**

11 state-space; stock assessment; random effects; time-varying; environmental effects; recruitment; natural  
12 mortality; Template Model Builder (TMB)

# 1 Introduction

The last two decades have increasingly seen a push for more holistic, ecosystem-based fisheries management (Larkin, 1996; Link, 2002). In part, this is a recognition that considering single species in isolation produces riskier and less robust outcomes long-term (Patrick and Link, 2015). In several high-profile cases, fisheries management has failed to prevent collapses because they did not reduce fishing pressure in responses to changes in natural mortality ( $M$ ), recruitment, or migration patterns caused by dynamics external to the stock in question (Northern cod: Shelton et al., 2006; Rose and Rowe, 2015; Gulf of Maine cod: Pershing et al., 2015; Pacific sardine: Zwolinski and Demer, 2012). This is particularly concerning in the context of climate change and the wide range of biological processes—often assumed to be constant—in stock assessments that are likely to be affected (Stock et al., 2011).

One approach to account for changing productivity is to explicitly link population processes to environmental covariates in single-species stock assessments, i.e. the mechanistic approach *sensu* Punt et al. (2014). Traditional single-species assessments are based on internal population dynamics and the effect of fishing mortality ( $F$ ), even though fisheries scientists have long known that these are important drivers of time-varying population processes, e.g. recruitment, mortality, growth, and movement (Garstang, 1900; Hjort, 1914). Effects of the environment or interactions with other species can be considered contextually, rather than explicitly as estimated parameters (although temporal variation in empirical weight and maturity at age can affect reference points). Despite how counterintuitive this may seem to ecologists and oceanographers who study such relationships, the evidence for direct linkages to specific environmental covariates is often weak and can break down over time (McClatchie et al., 2010; Myers, 1998). Additionally, the primary goal of most assessments is to provide management advice on near-term sustainable harvest levels—not to explain ecological relationships. Even if an environmental covariate directly affects fish productivity, including the effect in an assessment may not improve management advice if the effect is weak (De Oliveira and Butterworth, 2005). Worse, including environmental effects in an assessment or management system has been shown to actually provide worse management in some cases (De Oliveira and Butterworth, 2005; Punt et al., 2014; Walters and Collie, 1988). This can be true even in cases of relatively well-understood mechanistic links between oceanic conditions and fish populations, as in the case of sea surface temperature and Pacific sardine (Hill et al., 2018; Zwolinski and Demer, 2012). Still, incorporating mechanistic environment-productivity links in assessments does have the potential to reduce residual variance, particularly in periods when few demographic data exist (Miller et al., 2016; Shotwell et al., 2014).

An alternative approach is to allow biological parameters to vary stochastically over time, i.e. the empirical

approach *sensu* Punt et al. (2014). In this case, the variation is caused by a range of sources that are not explicitly modeled. Statistical catch-at-age (SCAA) models typically only estimate year-specific recruitment ( $R_t$ ) and  $F_t$ , often as deviations from a mean that may be a function of spawning biomass, e.g.  $\log R_t = \log R_0 + \epsilon_t$ . The main reason that other parameters are assumed constant is simply that there are not enough degrees of freedom to estimate many time-varying parameters. One common solution is to penalize the deviations, e.g.  $\epsilon_t \sim \mathcal{N}(0, \sigma_\epsilon^2)$ , although the penalty terms,  $\sigma_\epsilon^2$ , must be fixed or iteratively tuned and are therefore somewhat subjective (Aeberhard et al., 2018; Methot and Taylor, 2011; Methot and Wetzel, 2013; Xu et al., 2020). State-space models that treat parameters as unobserved states can, in principle, avoid such subjectivity by estimating the penalty terms as variance parameters constraining random effects and maximizing the marginal likelihood (Thorson, 2019). In this way, state-space models can allow processes to vary in time while simultaneously estimating fewer parameters.

Although state-space stock assessments have existed for some time (Gudmundsson, 1994; Mendelsohn, 1988; Sullivan, 1992), the recent development of Template Model Builder (TMB, Kristensen et al., 2016) software to perform efficient Laplace approximation has greatly expanded their use (Cadigan, 2016; Miller et al., 2016; Nielsen and Berg, 2014). In addition to the key advantage of objectively estimating variance, or “data weighting”, parameters, state-space models naturally predict unobserved states, and therefore handle missing data and short-term projections in a straightforward way (ICES, 2020). In comparisons with SCAA models, state-space models have been shown to have larger, more realistic, uncertainty and lower retrospective bias (Miller and Hyun, 2018; Stock et al., n.d.).

Retrospective bias can occur when changing environmental conditions lead to changes in productivity that are unaccounted for in stock assessments, and this is a concern common to several groundfish stocks on the Northeast U.S. Shelf (Brooks and Legault, 2016; Tableau et al., 2018). The Northeast U.S. Shelf ecosystem is rapidly changing, and this has motivated managers to make the “continue[d] development of stock assessment models that include environmental terms” a top priority (Hare et al., 2016). In addition to providing short-term (1-3 years) catch advice with reduced retrospective bias, it is hoped that environment-linked assessments will help create realistic rebuilding plans in the medium-term (3-10 years) for stocks that have not rebounded in response to dramatic decreases in  $F$ . For example, Miller et al. (2016) developed a state-space model for yellowtail flounder with an environmental effect on recruitment, which reduced retrospective patterns and residual variance. Additional applications using a similar state-space framework that include environmental effects on growth,  $M$ , and maturity have also proven promising (Miller and Hyun, 2018; Miller et al., 2018; O’Leary et al., 2019; Xu et al., 2018).

To address the needs of fisheries management in a changing climate, we seek an assessment framework

that combines both the empirical and mechanistic approaches. Namely, it should be able to 1) estimate time-varying parameters as random effects (i.e. a state-space model), and 2) include environmental effects directly on biological parameters. The framework should also allow for easy testing against status quo SCAA models to ease gradual adoption through the “research track” or “benchmark” assessment process (Lynch et al., 2018). The objectives of this manuscript are to introduce the Woods Hole Assessment Model (WHAM) framework and demonstrate its ability to:

1. estimate time- and age-varying random effects on annual changes in abundance at age,  $M$ , and selectivity;
2. fit environmental time-series with process and observation error, missing data, and a link to a population process; and
3. simulate new data and random effects to conduct self- and cross-tests (*sensu* Deroba et al., 2015) to estimate bias in parameters and derived quantities.

Throughout, we describe how the above are implemented using the open-source WHAM software package (Miller and Stock, 2020).

## 2 Methods

### 2.1 Model description

WHAM is a generalization and extension of Miller et al. (2016) in TMB. It is in many respects similar to the Age-Structured Assessment Program (ASAP, Legault and Restrepo, 1998; Miller and Legault, 2015) and can be configured to fit statistical catch-at-age models nearly identically. There is functionality built into WHAM to migrate ASAP input files to R inputs needed for WHAM, and WHAM uses many of the same types of data inputs, such as empirical weight-at-age, so that existing assessments in the U.S. Northeast can be easily replicated and tested against models with state-space and environmental effects in a single framework.

*Either put core model equations here or in supplement*

#### 2.1.1 Processes with random effects

WHAM primarily diverges from ASAP through the implementation of random effects on three processes: inter-annual transitions in numbers at age ( $NAA$ ), natural mortality ( $M$ ), and selectivity ( $s$ ), as well as allowing effects of environmental covariates ( $Ecov$ ) on recruitment and natural mortality (but see ASAP4;

Miller and Legault, 2015). The environmental covariates and their observations are treated using state-space models with true, unobserved values treated as random effects and observation on them having error. Other than environmental covariates, the processes are assumed to have a two dimensional (2D) first-order autoregressive (AR1) covariance structure over age and year, although correlation in either or both dimensions can be turned off. The 2D AR1 structure has been widely used to model deviations by age and year in the parameters  $F_{a,y}$  (Nielsen and Berg, 2014),  $M_{a,y}$  (Cadigan, 2016; Stock et al., n.d.),  $s_{a,y}$  (Xu et al., 2019), and  $N_{a,y}$  (Stock et al., n.d.), as well as in the catch ( $C_{a,y}$ ) and survey index ( $I_{a,y}$ ) observations (Berg and Nielsen, 2016).

#### 2.1.1.1 Numbers at age (NAA)

The stock equations in WHAM that describe the transitions between numbers at age are identical to Miller et al. (2016) and Nielsen and Berg (2014):

$$\log N_{a,y} = \begin{cases} \log(f(SSB_{y-1})) + \varepsilon_{1,y}, & \text{if } a = 1 \\ \log(N_{a-1,y-1}) - Z_{a-1,y-1} + \varepsilon_{a,y}, & \text{if } 1 < a < A \\ \log(N_{A-1,y-1}e^{-Z_{A-1,y-1}} + N_{A,y-1}e^{-Z_{A,y-1}}) + \varepsilon_{A,y}, & \text{if } a = A \end{cases} \quad (1)$$

where  $N_{a,y}$  are the numbers at age  $a$  in year  $y$ ,  $Z$  is the total mortality rate ( $F + M$ ),  $f$  is the stock-recruit function,  $Y$  is the total number of observation and prediction years, and  $A$  represents the plus-group. In this analysis we demonstrate four possible models for the NAA deviations,  $\varepsilon_{a,y}$ .

m1 is most similar to a SCAA model, where only recruitment deviations,  $\varepsilon_{1,y}$ , are estimated (i.e.  $\varepsilon_{a,y} = 0$  for  $a > 1$  in Eqn 1). In m1, the recruitment deviations are assumed to be independent and identically distributed (IID):

$$\varepsilon_{1,y} \sim \mathcal{N}\left(-\frac{\sigma_R^2}{2}, \sigma_R^2\right)$$

The  $-\frac{\sigma_R^2}{2}$  bias correction term is included by default so that the expected recruitment,  $E(N_{1,y} = R_y)$  equals the expected  $R_y$  from the stock-recruit function (Methot and Taylor, 2011; Thorson, 2019). This bias correction adjustment can also be turned off. The only difference between m1 and a SCAA is that annual recruitments are random effects and  $\sigma_R^2$  is an estimated parameter within the model.

m2 is the same as m1, except that the recruitment deviations are stationary AR1 with autocorrelation parameter  $-1 < \rho_y < 1$ :

$$\varepsilon_{1,y+1} \sim \mathcal{N}\left(\rho_y \varepsilon_{1,y} - \frac{\sigma_R^2}{2(1-\rho_y^2)}, \sigma_R^2\right)$$

m3 is the “full state-space” model from Nielsen and Berg (2014) and Miller et al. (2016), where all numbers at age are independent random effects and:

$$\varepsilon_{a,y} \sim \begin{cases} \mathcal{N}\left(-\frac{\sigma_R^2}{2}, \sigma_R^2\right), & \text{if } a = 1 \\ \mathcal{N}\left(-\frac{\sigma_a^2}{2}, \sigma_a^2\right), & \text{if } a > 1 \end{cases} \quad (2)$$

where  $\sigma_a^2$  for all ages  $a > 1$  are assumed to be the same but different from age  $a = 1$ , i.e. recruitment. This assumption is sensible because variability of deviations between expected and realized recruitment are typically larger than deviations from expected abundance at older ages.

m4 treats the numbers at all ages as random effects, as in m3, but the *NAA* deviations have a 2D stationary AR1 structure as in Stock et al. (n.d.):

$$\mathbf{E} \sim \mathcal{MVN}(0, \Sigma)$$

where  $\mathbf{E} = (\varepsilon_{1,1}, \dots, \varepsilon_{1,Y-1}, \varepsilon_{2,1}, \dots, \varepsilon_{2,Y-1}, \dots, \varepsilon_{A,1}, \dots, \varepsilon_{A,Y-1})'$  is a vector of all *NAA* deviations,  $\Sigma$  is the covariance matrix of  $\mathbf{E}$  defined by:

$$\text{Cov}(\varepsilon_{a,y}, \varepsilon_{\tilde{a},\tilde{y}}) = \frac{\sigma_a \sigma_{\tilde{a}} \rho_a^{|a-\tilde{a}|} \rho_y^{|y-\tilde{y}|}}{(1-\rho_a^2)(1-\rho_y^2)}$$

and  $-1 < \rho_a < 1$  and  $-1 < \rho_y < 1$  are the AR1 coefficients in age and year, respectively. As in m3,  $\sigma_a^2$  for all ages  $a > 1$  are assumed to be the same but different from age  $a = 1$ ,  $\sigma_R^2$ . The bias correction term for age  $a > 1$  in m4 is  $-\frac{\sigma_a^2}{2(1-\rho_y^2)(1-\rho_a^2)}$ .

#### 2.1.1.2 Natural mortality ( $M$ )

For natural mortality, there are mean parameters for each age,  $\mu_{M_a}$ , each of which may be estimated freely or fixed at the initial values. The  $\mu_{M_a}$  may also be estimated in sets of ages, e.g. estimate one mean  $M$  shared across ages 3-5,  $\mu_{M_3} = \mu_{M_4} = \mu_{M_5}$ . There is also an option for  $M$  to be specified as a function of weight-at-age,  $M_{a,y} = \mu_M W_{a,y}^b$ , as in Lorenzen (1996) and Miller and Hyun (2018). Regardless of whether  $\mu_{M_a}$  are fixed or estimated, WHAM can also be configured to estimate deviations in  $M$ ,  $\delta_{a,y}$ , as random effects analogous to the *NAA* deviations (Cadigan, 2016; Stock et al., n.d.):

$$\begin{aligned}\log(M_{a,y}) &= \mu_{M_a} + \delta_{a,y} \\ \text{Cov}(\delta_{a,y}, \delta_{\tilde{a},\tilde{y}}) &= \frac{\sigma_M^2 \varphi_a^{|a-\tilde{a}|} \varphi_y^{|y-\tilde{y}|}}{(1-\varphi_a^2)(1-\varphi_y^2)}\end{aligned}\tag{3}$$

where  $\sigma_M^2$ ,  $\varphi_a$ , and  $\varphi_y$  are the AR1 variance and correlation coefficients in age and year, respectively.

In this analysis, we demonstrate three alternative  $M$  random effects models. For simplicity, all models treat  $\mu_{M_a}$  as known, as in each of the original assessments. m1 is identical to the base *NAA* model, with no random effects on  $M$  ( $\sigma_M^2 = \varphi_a = \varphi_y = 0$  and not estimated). m2 allows IID  $M$  deviations, estimating  $\sigma_M^2$  but fixing  $\varphi_a = \varphi_y = 0$ . m3 estimates the full 2D AR1 structure for  $M$  deviations.

### 2.1.1.3 Selectivity ( $s$ )

As in ASAP and many other SCAA assessment frameworks, WHAM assumes separability in the fishing mortality rate by age and year, e.g.  $F_{a,y} = F_y s_a$ , where  $F_y$  is the “fully selected” fishing mortality rate in year  $y$  and  $s_a$  is the selectivity at age. We note that this differs from the approach in SAM (Nielsen and Berg, 2014), where the  $F_{a,y}$  are estimated directly as multivariate random effects without the separability assumption. Three parametric forms are available (logistic, double-logistic, and decreasing-logistic), as well as a non-parametric option to estimate each  $s_a$  individually (“age-specific”). To allow for temporal changes in selectivity as in ASAP, WHAM can estimate selectivity in user-specified time blocks. WHAM estimates selectivity parameters on the logit scale to avoid boundary problems during estimation.

WHAM estimates annual full  $F_y$  and mean selectivity parameters as fixed effects. Deviations in selectivity parameters can be estimated as random effects,  $\zeta_{p,y}$ , with autocorrelation by parameter ( $p$ ), year ( $y$ ), both, or neither. This is done similarly to Xu et al. (2019), except that the deviations are placed on the parameters instead of the mean  $s_{a,y}$  in order to guarantee that  $0 < s_{a,y} < 1$ . For example, logistic selectivity with two parameters  $a_{50}$  and  $k$  is estimated as:

$$\begin{aligned}s_{a,y} &= \frac{1}{1+e^{-(a-a_{50})/k_y}} \\ a_{50,y} &= l_{a_{50}} + \frac{u_{a_{50}} - l_{a_{50}}}{1+e^{-(\nu_1+\zeta_{1,y})}} \\ k_y &= l_k + \frac{u_k - l_k}{1+e^{-(\nu_2+\zeta_{2,y})}} \\ \text{Cov}(\zeta_{1,y}, \zeta_{2,\tilde{y}}) &= \frac{\sigma_s^2 \phi_p \phi_y^{|y-\tilde{y}|}}{(1-\phi_p^2)(1-\phi_y^2)}\end{aligned}\tag{4}$$

where  $\nu_1$  is the logit-scale mean  $a_{50}$  parameter with lower and upper bounds  $l_{a_{50}}$  and  $u_{a_{50}}$ ,  $\nu_2$  is the logit-scale mean  $k$  parameter with lower and upper bounds  $l_k$  and  $u_k$ ,  $\sigma_s^2$  is the AR1 variance, and  $\phi_p$ , and  $\phi_y$  are the AR1 correlation coefficients by parameter and year.



Below, we demonstrate three models with random effect deviations on logistic selectivity, akin to those for  $M$ . m1 treats all numbers at age as independent random effects (i.e.  $NAA$  m3) but with no random effects on  $s$  ( $\sigma_s^2 = \phi_p = \phi_y = 0$  and not estimated). m2 allows IID  $s$  deviations, estimating  $\sigma_s^2$  but fixing  $\phi_p = \phi_y = 0$ . m3 estimates the full 2D AR1 structure for  $s$  deviations.

#### 2.1.1.4 Environmental covariates (*Ecov*)

WHAM models environmental covariate data using state-space models with process and observation components. The true, unobserved values (or “latent states”,  $X_y$ ) are then linked to the population dynamics equations with user-specified lag. For example, recruitment in year  $y$  may be influenced by  $X_{y-1}$  (lag 1), while natural mortality in year  $y$  may be influenced by  $X_y$  (lag 0). Multiple environmental covariates may be included, but only as independent processes. The *Ecov* and population model years do not need to match, and missing years are allowed. In particular, including *Ecov* data in the projection period can be useful.

##### 2.1.1.4.1 Process model

There are currently two options in WHAM for the *Ecov* process model: a normal random walk and AR1. We model the random walk as in Miller et al. (2016):

$$X_{y+1}|X_y \sim \mathcal{N}(X_y, \sigma_X^2)$$

where  $\sigma_X^2$  is the process variance and  $X_1$  is estimated as a fixed effect parameter. One disadvantage of the random walk is that its variance is nonstationary. In short-term projections,  $\hat{X}_y$  will be equal to the last estimate with an observation and the uncertainty of  $\hat{X}_y$  will increase over time. If  $\hat{X}_y$  influences reference points, this leads to increasing uncertainty in stock status over time as well (Miller et al., 2016).

For this reason, we generally prefer to model  $X_y$  as a stationary AR1 process as in Miller et al. (2018):

$$\begin{aligned} X_1 &\sim \mathcal{N}\left(\mu_X, \frac{\sigma_X^2}{1-\phi_X^2}\right) \\ X_y &\sim \mathcal{N}\left(\mu_X(1-\phi_X) + \phi_X X_{y-1}, \sigma_X^2\right) \end{aligned} \tag{5}$$

where  $\mu_X$ ,  $\sigma_X^2$ , and  $|\phi_X| < 1$  are the marginal mean, variance, and autocorrelation parameters. In addition to having stationary variance, another important difference between the random walk and AR1 in short-term projections is that the AR1 will gradually revert to the mean over time, unless environmental covariate observations are included in the projection period.

#### 2.1.1.4.2 Observation model

The environmental covariate observations,  $x_y$ , are assumed to be normally distributed with mean  $X_y$  and variance  $\sigma_{x_y}^2$ :

$$x_y|X_y \sim \mathcal{N}\left(X_y, \sigma_{x_y}^2\right)$$

The observation variance in each year,  $\sigma_{x_y}^2$ , can be treated as known with year-specific values (as in Miller et al., 2016) or one overall value shared among years. They can also be estimated as parameters, likewise either as yearly values or one overall value. If included, WHAM estimates yearly  $\sigma_{x_y}^2$  as random effects with parameters  $\mu_{\sigma_x}$  and  $\sigma_{\sigma_x}$ :

$$\sigma_{x_y}^2 \sim \mathcal{N}\left(\mu_{\sigma_x}, \sigma_{\sigma_x}^2\right)$$

#### 2.1.1.4.3 Link to population

WHAM currently provides options to link the modeled environmental covariate,  $X_y$ , to the population dynamics via recruitment or natural mortality. It is also sometimes useful to fit the *Ecov* model without a link to the population dynamics so that models with and without environmental effects have the same data in the likelihood and can be compared via AIC.

In the case of recruitment, the options follow Fry (1971) and Iles and Beverton (1998): “controlling” (density-independent mortality), “limiting” (carrying capacity effect, e.g.  $X_y$  determines the amount of suitable habitat), “lethal” (threshold effect, i.e.  $R_t$  goes to 0 at some  $X_y$  value), “masking” ( $X_y$  decreases  $dR/dSSB$ , as expected if  $X_y$  affects metabolism or growth), and “directive” (e.g. behavioral). Of these, WHAM currently allows controlling, limiting, or masking effects in the Beverton-Holt stock-recruit function, and controlling or masking effects in the Ricker function. For natural mortality, environmental effects are placed on  $\mu_M$ , shared across ages.

Regardless of where the environment-population link is, the effect can be either linear or polynomial. Nonlinear effects of environmental covariates are common in ecology, and quadratic effects are to be expected in cases where intermediate values are optimal (Agostini et al., 2008; Brett, 1971). WHAM includes a function to calculate orthogonal polynomials in TMB, akin to the `poly()` function in R.

In this analysis, we compare five models with limiting effects on Beverton-Holt recruitment:

$$\hat{R}_{y+1} = \frac{\alpha \text{SSB}_y}{1 + e^{\beta_0 + \beta_1 X_y + \beta_2 X_y^2} \text{SSB}_y}$$

where  $\text{SSB}_y$  is spawning stock biomass in year  $y$ ,  $\alpha$  and  $\beta_0$  are the standard parameters of the Beverton-Holt function, and  $\beta_1$  and  $\beta_2$  are polynomial effect terms that modify  $\beta_0$  based on the value of the estimated environmental covariate,  $X_y$ .

m1 treats  $X_y$  as a random walk ( $\phi_X = 1$ ) but does not include an effect on recruitment ( $\beta_1 = \beta_2 = 0$ ). We include m1 in order to compare AIC of the original model to those with effects on recruitment. m2 and m3 also treat  $X_y$  as a random walk, but m2 estimates  $\beta_1$  and m3 estimates both  $\beta_1$  and  $\beta_2$ . m4 and m5 estimate  $X_y$  as an AR1 process, and m4 estimates  $\beta_1$  and m5 estimates both  $\beta_1$  and  $\beta_2$ .

### 2.1.2 Population observation model

Like ASAP, there are observation likelihood components for aggregate catch and abundance index for each fleet and index, and age composition for each fleet and index.

#### 2.1.2.1 Aggregate catch and indices

The predicted catch at age for fleet  $i$ ,  $\hat{C}_{a,y,i}$ , is a function of  $N_{a,y}$ ,  $M_{a,y}$ ,  $F_{y,i}$ ,  $s_{a,y,i}$ , and empirical weight at age,  $W_{a,y,i}$ :

$$\hat{C}_{a,y,i} = N_{a,y} W_{a,y,i} (1 - e^{-Z_{a,y}}) \frac{F_{a,y,i}}{Z_{a,y}}.$$

The log-aggregate catch  $\hat{C}_{y,i} = \sum_a \hat{C}_{a,y,i}$  observation is assumed to have a normal distribution

$$\log(C_{y,i}) \sim \mathcal{N}\left(\log(\hat{C}_{y,i}) - \frac{\sigma_{\hat{C}_{y,i}}^2}{2}, \sigma_{\hat{C}_{y,i}}^2\right). \quad (6)$$

where the standard deviation

$$\sigma_{C_{y,i}} = e^{\eta_i} \sigma_{\hat{C}_{y,i}}$$

is a function of an input standard deviation  $\sigma_{\hat{C}_{y,i}}$  and a fleet-specific parameter  $\eta_i$  that is fixed at 0 by default, but may be estimated. The bias correction term,  $-\frac{\sigma_{\hat{C}_{y,i}}^2}{2}$ , is included by default based on Aldrin et al. (2020) but can be turned off.

Observations of aggregate indices of abundance are handled identically to the aggregate catch as in Eqn. 6

except that for index  $i$  the predicted index at age is

$$\hat{I}_{a,y,i} = q_i s_{a,y,i} N_{a,y} W_{a,y,i} e^{-Z_{a,y} f_{y,i}}$$

where  $q_i$  is the catchability and  $f_{y,i}$  is the fraction of the annual time step elapsed when the index is observed.

There are options for indices to be in terms of abundance (numbers) or biomass and  $W_{a,y,i} = 1$  for the former.

### 2.1.2.2 Catch and index age composition

WHAM includes several options for the catch and index age compositions including multinomial (default), Dirichlet, Dirichlet-multinomial, logistic normal and a zero-one inflated logistic normal. In all of the applications and simulation studies here, we assumed age composition observations were multinomial distributed.

### 2.1.3 Projections

The default settings for short-term projections follow common practice for stock assessments in the U.S. Northeast: the population is projected three years using the average selectivity, maturity, weight, and natural mortality at age from the last five model years to calculate reference points (NEFSC, 2020a). WHAM implements similar options as ASAP for specifying  $F_y$  in the projection years: terminal year  $F_y$ , average  $F$  over specified years,  $F_{X\%}$  ( $F$  at  $X\%$  SPR, where  $X$  is specified and 40 by default), *should we quickly include an option for projecting  $F_{MSY}$* , user-specified  $F_y$ , or  $F$  derived from user-specified catch. For all options except user-specified  $F_y$ , the uncertainty in projected  $F$  is propagated into the uncertainty of projected population attributes. For models with random effects on  $NAA$ ,  $M$ , or  $Ecov$ , the default is to continue the stochastic process into the projection years. WHAM does not currently do this for selectivity because, like  $F$ , it is a function of management. Instead, selectivity is taken as the average of recent model years.

If the  $Ecov$  data extend beyond the population model years, WHAM will fit the  $Ecov$  model to all available data and use the estimated  $X_y$  in projections. This may often be the case because lags can exist in both the physical-biological mechanism and the assessment process. As an example, a model with an  $Ecov$  effect on recruitment may link physical conditions in year  $t$  to recruitment in year  $t + 1$ , and the assessment conducted in year  $t$  may only use population data through year  $t - 1$ . In this case, 3-year population projections only need the  $Ecov$  model to be projected one year. While the default handling of  $Ecov$  projections is to continue the stochastic process, WHAM includes options to use terminal year  $x_y$ ,  $x_y$  averaged over specified years, or specified  $x_y$ . The option to specify  $x_y$  allows users to investigate how alternative climate projections may affect the stock.

## 2.2 Fits to original datasets

We fit the models described above to data from five stocks with a broad range of life history, status, and model dimension (number of ages and years): Southern New England-Mid Atlantic (SNEMA) yellowtail flounder, butterfish, North Sea cod, Icelandic herring, and Georges Bank (GB) haddock (Tables 1 and 2). We fit the *NAA* random effects models to all five stocks since these represent core WHAM functionality. We chose to highlight one stock each for the *M*, *s*, and *Ecov* processes because all models did not converge for all stocks and processes: butterfish (*M*), GB haddock (*s*), and SNEMA yellowtail flounder (*Ecov*). We fixed  $\mu_{M_a}$  at the values used or estimated in the original assessments. Except for the GB haddock *s* models, we used the same selectivity parameterization and time blocks as in the original assessments. We used the m3 *NAA* model (all *NAA* deviations are IID random effects) in the *s* and *Ecov* demonstrations, but the m1 *NAA* model (only recruitment deviations are IID random effects) for the *M* demonstrations, because the *NAA* transitions can be interpreted as survival and including random effect deviations on both *NAA* and *M* can lead to model non-convergence (Stock et al., n.d.). Within each process, we compared the performance of the different random effects models using AIC.

We fit all models using the open-source statistical software R (R Core Team, 2020) and TMB (Kristensen et al., 2016), as implemented in the R package WHAM (Miller and Stock, 2020). Documentation and tutorials for how to specify additional random effect structures in WHAM are available at <https://timjmiller.github.io/wham/>. Code and data files to run the analysis are available at <https://github.com/brianstock-NOAA/wham-sim>.

## 2.3 Simulation tests

After fitting each model to the original datasets, we used the simulation feature of TMB to conduct self- and cross-tests (*sensu* Deroba et al., 2015). For each model, we generated 100 sets of new data and random effects, keeping the fixed effect parameters constant at values estimated in original fits. We then re-fit all models to datasets simulated under each as an operating model. We calculated the relative error in parameters constraining random effects (Table 1) and quantities of interest, such as spawning stock biomass (SSB),  $F$ ,  $\frac{B}{B_{40\%}}$ ,  $\frac{F}{F_{40\%}}$ , and  $R$ . We calculated relative error as  $\frac{\hat{\theta}_i}{\theta_i} - 1$ , where  $\theta_i$  is the true value for simulated data set  $i$  and  $\hat{\theta}_i$  is the value estimated from fitting the model to the simulated data. To estimate bias of each estimation model for a given operating model, we calculated 95% confidence intervals of the median relative error using the binomial distribution (Thompson, 1936). To summarize the bias across simulations and years for each model, we calculated the median quantity across years and took the mean of the medians across simulations. Finally, for each operating model we calculated the proportion of simulations in which each

estimation model converged and had the lowest AIC, aggregated across stocks.

## 3 Results

### 3.1 Original datasets

The 2D AR1 covariance structure for random effects on  $NAA$ ,  $M$ , and selectivity performed well across stocks and processes, evidenced by lower AIC than the IID random effects models (Fig. 1). This AIC difference was larger for selectivity than  $NAA$  or  $M$ , but the differences for  $NAA$  and  $M$  were also non-trivial, ranging from 11.1–53.2.

#### 3.1.1 Numbers-at-age ( $NAA$ )

Treating all ages as random effects was strongly supported by AIC, compared to treating only recruitment deviations as random effects (Fig. 1). Including autoregressive  $NAA$  random effects was also generally supported by AIC, either the AR1 when only recruitment deviations were random effects or the 2D AR1 when all ages were random effects. The estimated AR and IID random effects were similar, with the AR random effects slightly smoothed compared to the IID random effects (e.g. for Icelandic herring in Fig. 2).

#### 3.1.2 Natural mortality ( $M$ )

As for the  $NAA$  models, including random effect deviations on  $M$  was supported by AIC (Fig. 1). Although the 2D AR1  $M$  model did not converge for North Sea cod, it had the lowest AIC for butterfish and SNEMA yellowtail flounder. In contrast to the  $NAA$  models, the patterns in estimated 2D AR1 and IID  $M$  random effects differed noticeably (e.g. for butterfish in Fig. 3). The butterfish IID  $M$  model estimated elevated  $M$  for age 5+ fish early and late in the time-series, with lower  $M$  in intervening years. The butterfish 2D AR1  $M$  model estimated a similar, but much exaggerated, pattern for age 5+ fish and reduced  $M$  for ages 1 and 4. Since  $\mu_M$  was held constant, the  $M$  random effects models had to decrease  $M$  in some ages and years in order to increase  $M$  in others.

#### 3.1.3 Selectivity ( $s$ )

For Georges Bank haddock, including 2D AR1 random effect deviations on selectivity was strongly supported by AIC (Fig. 1). The 2D AR1  $s$  model estimated similar patterns in selectivity compared to the IID  $s$  model,

except with smoother variations by age and year (Fig. 4). Compared to the model with constant selectivity, the IID and 2D AR1 models estimated lower  $s$  for age 3 in recent years and higher  $s$  for age 3 before 1990. They also estimated higher  $s$  for age 2 before 1990, especially from 1973 to 1976.

### 3.1.4 Environmental covariate effect on recruitment (*Ecov*)

As in previous analyses (Miller et al., 2016; Xu et al., 2018), including an effect of the CPI on recruitment for SNEMA yellowtail flounder was clearly supported by AIC ( $\Delta$ AIC of 20.3-33.0 between m1 and m2-m4, Fig. 1). The AR1-linear model (m4) had the lowest AIC—fitting the CPI using an AR1 model was preferred over the random walk ( $\Delta$ AIC of 12.7 between m2 and m4), and the quadratic term,  $\beta_2$  was deemed unnecessary ( $\Delta$ AIC of 1.3 between m5 and m4). As expected, the AR1 process model estimated higher uncertainty in years with higher observation error (e.g. 1982–1992) and missing observations (2017, Fig. 5A). The CPI negatively influenced recruitment, i.e.  $\hat{R}$  was higher following years with lower CPI (lower fall bottom temperature, Fig. 5C). Including the CPI-recruitment link noticeably changed the estimates of SSB and  $R$  in some years (compare locations of points between Fig. 5B-C).

## 3.2 Simulation tests

Several findings were consistent across processes and stocks. When the estimation and operating model were consistent, bias of SSB,  $F$ ,  $\frac{B}{B_{40\%}}$ ,  $\frac{F}{F_{40\%}}$ ,  $R$ , and the variance and correlation parameters controlling random effects was generally small and not significant based on confidence intervals. Bias was also generally small when more complex models were fitted to less complicated operating model simulations. In contrast, the bias was often non-trivial when the estimation model was less complex than the operating model, especially when models without random effects were fit to data simulated from models with random effects. Bias in SSB and  $F$  were always opposite; when SSB was biased high,  $F$  was biased low, and vice-versa. Predicted catch was never biased. Bias of variance parameters was either negligible or negative, as expected. Restricted maximum likelihood (REML) should be used if more accurate estimation of these parameters is a priority. In cross-tests, the percentage of simulations in which AIC selected the correct model varied between 62–99% by process, stock, and operating model (Fig. 6). Estimation models more complex than the operating model were more likely to be chosen for  $NAA$  models than for  $M$  or selectivity.

### 3.2.1 Numbers-at-age

For four of the five stocks,  $\sigma_R^2$  was estimated without bias in self-tests (Fig. 7). The exception was butterfish, for which  $\sigma_R^2$  was negatively biased in m1, m2, and m4 (but not m3). In contrast, both m3 and m4 estimated  $\sigma_a^2$  with negative bias for all five stocks. There was no consistent pattern in bias for the correlation parameters  $\rho_a$  and  $\rho_y$ .

Across the five stocks, all NAA models estimated SSB,  $F$ ,  $\frac{B}{B_{40\%}}$ ,  $\frac{F}{F_{40\%}}$ , and  $R$  with very minimal bias in self-tests. An exception was the estimation of  $F$  for Icelandic herring, particularly for the SCAA models (median relative error with 95% CI for m1: -0.060 (-0.073, -0.047), m2: -0.060 (-0.073, -0.047), m3: 0.015 (-0.004, 0.034), and m4: 0.049 (0.017, 0.080); Fig. 8). The convergence rate for most models and stocks was above 95%, again with the exception of the SCAA models for Icelandic herring (ref supp figure). In cross-tests, SCAA models exhibited non-trivial bias when estimating data simulated from models that treated numbers at all ages as random effects. The degree of bias varied by quantity and stock between -25% and 25% (ref supp figures). The more complex NAA models estimated all quantities without bias regardless of operating model.

### 3.2.2 Natural mortality

As for the NAA models,  $\sigma_R^2$  was estimated without bias in the  $M$  model self-tests (Fig. 9).  $\sigma_M^2$  was biased low,  $\varphi_y$  had no bias, and the direction of bias in  $\varphi_a$  was inconsistent.

SSB,  $F$ ,  $\frac{B}{B_{40\%}}$ ,  $\frac{F}{F_{40\%}}$ , and  $R$  were estimated without significant bias in self-tests, although m1 and m2 exhibited bias in SSB and  $\frac{F}{F_{40\%}}$  when fit to data simulated from m3 (e.g. for butterfish, Fig. 10).

### 3.2.3 Selectivity

For Georges Bank haddock, the variance and correlation parameters  $\sigma_a^2$ ,  $\sigma_s^2$ ,  $\phi_y$ , and  $\phi_a$  were all estimated with slight negative bias in self-tests of models with  $s$  random effects (Fig. 11).

Models with IID or 2D AR1  $s$  random effects estimated SSB,  $F$ ,  $\frac{B}{B_{40\%}}$ ,  $\frac{F}{F_{40\%}}$ , and  $R$  with little to no bias across operating models (Fig. 12). The model without  $s$  random effects showed substantial bias when fit to data simulated from m2 or m3.



### 3.2.4 Ecov-Recruitment

The random walk CPI models estimated  $\sigma_X^2$  with less bias than the AR1 CPI models (Fig. 13).  $\beta_0$  was biased high in all models, although this was not significant for the model with lowest AIC (m4, AR1-linear). All models estimated the parameters  $\phi_X$ ,  $\alpha$ ,  $\beta_1$ , and  $\beta_2$  without significant bias.

## 4 Discussion

### 4.1 Overview

Our results suggest that the WHAM package can be a useful tool for stock assessment when environmental effects on recruitment or natural mortality, variation in selectivity or natural mortality, or stochastic changes in the numbers at age are of interest. The simulation tests showed negligible or no bias in estimation of important assessment outputs (SSB,  $F$ , stock and harvest status) when the operating and estimation models matched. The less complex models, without random effects or autoregressive structure, exhibited some bias in cross-tests, while the more complex models did not. In these cases, bias in SSB and  $F$  were opposite such that predicted catch was unbiased. The WHAM models with IID or 2D AR1 random effect deviations performed well across stocks and processes, which suggests that they warrant consideration in future stock-specific studies.

### 4.2 Differences from other assessment model frameworks

WHAM can estimate interannual transitions in NAA as random walk processes like SAM, or as a function of autoregressive deviations. Alternatively (or simultaneously, Stock et al., n.d.) WHAM can estimate deviations in natural mortality as autoregressive processes like NCAM (Cadigan, 2016). Uniquely, WHAM can model multiple environmental covariate time series as state-space processes and include their effects (nonlinearly) in various ways on recruitment or natural mortality.

WHAM assumes separability in  $F_{a,y}$ , i.e.  $F_{a,y} = F_y s_a$ , and estimates annual full  $F_y$  as fixed effects and selectivity of surveys or fisheries as constant or time-varying random effects with various autoregressive assumptions possible. Most other assessment frameworks in the U.S, e.g. ASAP, Stock Synthesis (SS), an Assessment Model for Alaska (AMAK), and the Beaufort Assessment Model (BAM), make this same separability assumption because oftentimes we want to specify  $F$  in projections to calculate a reference point, as opposed to continuing a  $F$  time-series process. Estimating selectivity in time blocks is common practice

in these frameworks. In contrast, SAM estimates  $F_{a,y}$  directly as a multivariate stochastic process (Nielsen and Berg, 2014). The approach in WHAM of estimating  $F_y$ ,  $\sigma_s$ ,  $\phi_y$ , and  $\phi_p$  as fixed effect parameters and allowing 2D AR1 random effects on the selectivity parameters should allow for similar selectivity patterns as in SAM. The method in Stock Synthesis is similar: allow 2D AR1 random effects on  $s_{a,y}$  instead of the parameters, and then estimate  $\sigma_s$  by an iterative tuning algorithm (Xu et al., 2019). It is unclear whether estimating time-varying selectivity as random effects produces different results or an improvement over using time blocks, or if there are advantages to using any of the three approaches for estimating time-varying selectivity used by WHAM, SAM, or Stock Synthesis.

Likewise, WHAM and SAM also make different separability assumptions for the catch and index observations. WHAM treats aggregate and composition observations separately for fisheries and indices, as do ASAP, SS, AMAK, and BAM. This observation model is natural for landings data that are measured as total weight plus age composition sampling, with survey age composition sampling often done separately. This differs from SAM where  $C_{a,y}$  and  $I_{a,y}$  observations are treated directly as multivariate log-normal.

### 4.3 Future use of WHAM

The productivity of several groundfish stocks in the U.S. Northeast has declined in recent decades, and conducting assessments in WHAM with time-varying processes via random effects or environment-productivity links could account for these trends and potentially reduce retrospective bias (Perretti et al., 2017; Stock et al., n.d.; Tableau et al., 2018). WHAM can be configured to fit SCAA models very similar to ASAP and therefore bridge between the two frameworks. Including random effects or environmental covariates in an assessment model is a large structural change, and simulation cross-tests such as we have demonstrated here should be conducted through the research track process (Lynch et al., 2018). If comparisons against status quo SCAA models prove favorable, WHAM could transition to being used in operational assessments. However, because the details of how to include time-varying processes, as well as the effect on the assessment output, will vary by stock, this evaluation likely needs to be conducted on a stock-by-stock basis.

Random effects allow for changing productivity and, if they are considered as an autoregressive process, can propagate the effect of these changes on assessment output in short-term projections (Stock et al., n.d.). However, the AR1 process demonstrated here reverts to the mean in projections and will not predict values beyond extremes in observed time-series. This is an issue worth examining because many marine ecosystems are changing to such extent that recent conditions are time-series extremes, and conditions in the near future may have never been observed (e.g., sea surface temperatures over the U.S. Northeast Shelf in the last decade;

Chen et al., 2020). More complex nonstationary time-series models such as autoregressive integrated moving average (ARIMA) that can forecast beyond observed values could be implemented. Nonlinear (e.g. splines) or double linear integration (Di Lorenzo and Ohman, 2013) techniques could also be worth pursuing, as well as time-delay embedding methods (Munch et al., 2018, 2017), which not only allow for nonstationary dynamics but also do not require a specified functional form. Still, stochastic projections of productivity responses to environmental conditions that are rapidly changing and at time-series extremes should be done with care.

In addition to the empirical (i.e. random effects) approach, it is worth considering the mechanistic approach (i.e. explicitly modeled links to environmental covariates) for a given stock whose recruitment or  $M$  is suspected to have shifted due to a clear, plausible hypothesized external influence. Several factors may result in a higher likelihood of determining that the mechanistic approach is useful in an assessment: a history of overfishing (Free et al., 2019; but high  $F$  can also swamp the signal of an environmental influence, Haltuch and Punt, 2011), more rapid environmental change, stocks at the edge of species' range, opportunistic (short-lived) species (*sensu* Winemiller and Rose, 1992), lower trophic level species, longer time series, periodic signals in which more than once cycle has been recorded (e.g. Pacific Decadal Oscillation and sardine), and stronger signals (wider ranges of observed stock status and environmental conditions) (Free et al., 2019; Haltuch et al., 2019; Haltuch and Punt, 2011; Marshall et al., 2019).

Although including external drivers of productivity may increase model uncertainty without improving accuracy, stock assessments that do not allow for time-varying productivity underestimate model uncertainty and implicitly assume that conditions will return to average in forecasts. In lieu of operationalizing an environment-linked WHAM model to set catch advice, WHAM can still be useful in two main ways. First, WHAM can serve as an operating model to simulate realistic environmentally-driven changes in recruitment or  $M$  in order to estimate the sensitivity of status quo models. Second, WHAM could account for uncertainty in environment-productivity relationships by constructing ensembles of alternative states of nature (Anderson et al., 2017; Möllmann et al., 2014).

## 4.4 Conclusion

A major present-day challenge in fisheries is to assess and manage stocks in a changing environment. We foresee environment-linked stock assessments increasingly becoming feasible as fisheries and oceanographic time-series lengthen and our ecological understanding deepens. We have developed WHAM with this in mind. Finally, we note that the development of TMB has been a critical advance for fisheries assessment models such as WHAM, allowing us to treat population and environmental processes as time-varying random effects

452 in a state-space framework.

## 453 **Acknowledgements**

454 This research was performed while BCS held an NRC Research Associateship award at the NEFSC. NOAA  
455 Fish & Climate grant number ??.

<sup>456</sup> **Supplementary material**

<sup>457</sup> More figures.

## References

- Aeberhard, W.H., Mills Flemming, J., Nielsen, A., 2018. Review of State-Space Models for Fisheries Science. *Annu. Rev. Stat. Appl.* 5, 215–235. <https://doi.org/10.1146/annurev-statistics-031017-100427>
- Agostini, V., Hendrix, A., Hollowed, A., Wilson, C., Pierce, S., Francis, R., 2008. Climate-ocean variability and Pacific hake: A geostatistical modeling approach. *Journal of Marine Systems* 71, 237–248. <https://doi.org/10.1016/j.jmarsys.2007.01.010>
- Aldrin, M., Tvete, I., Aanes, S., Subbey, S., 2020. The specification of the data model part in the SAM model matters. *Fisheries Research* 229, 105585. <https://doi.org/10.1016/j.fishres.2020.105585>
- Anderson, S.C., Cooper, A.B., Jensen, O.P., Minto, C., Thorson, J.T., Walsh, J.C., Afflerbach, J., Dickey-Collas, M., Kleisner, K.M., Longo, C., Osio, G.C., Ovando, D., Mosqueira, I., Rosenberg, A.A., Selig, E.R., 2017. Improving estimates of population status and trend with superensemble models. *Fish and Fisheries* 18, 732–741. <https://doi.org/10.1111/faf.12200>
- Berg, C.W., Nielsen, A., 2016. Accounting for correlated observations in an age-based state-space stock assessment model. *ICES J Mar Sci* 73, 1788–1797. <https://doi.org/10.1093/icesjms/fsw046>
- Brett, J.R., 1971. Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus Nerka*). *Am Zool* 11, 99–113. <https://doi.org/10.1093/icb/11.1.99>
- Brooks, E.N., Legault, C.M., 2016. Retrospective forecasting evaluating performance of stock projections for New England groundfish stocks. *Can. J. Fish. Aquat. Sci.* 73, 935–950. <https://doi.org/10.1139/cjfas-2015-0163>
- Cadigan, N.G., 2016. A state-space stock assessment model for northern cod, including under-reported catches and variable natural mortality rates. *Canadian Journal of Fisheries and Aquatic Sciences* 73, 296–308. <https://doi.org/10.1139/cjfas-2015-0047>
- Chen, Z., Kwon, Y.-O., Chen, K., Fratantoni, P., Gawarkiewicz, G., Joyce, T.M., 2020. Long-Term SST Variability on the Northwest Atlantic Continental Shelf and Slope. *Geophys. Res. Lett.* 47. <https://doi.org/10.1029/2019GL085455>
- De Oliveira, J., Butterworth, D., 2005. Limits to the use of environmental indices to reduce risk and/or increase yield in the South African anchovy fishery. *African Journal of Marine Science* 27, 191–203. <https://doi.org/10.2989/18142320509504078>

Deroba, J.J., Butterworth, D.S., Methot, R.D., De Oliveira, J.a.A., Fernandez, C., Nielsen, A., Cadrin, S.X., Dickey-Collas, M., Legault, C.M., Ianelli, J., Valero, J.L., Needle, C.L., O'Malley, J.M., Chang, Y.-J., Thompson, G.G., Canales, C., Swain, D.P., Miller, D.C.M., Hintzen, N.T., Bertignac, M., Ibaibarriaga, L., Silva, A., Murta, A., Kell, L.T., de Moor, C.L., Parma, A.M., Dichmont, C.M., Restrepo, V.R., Ye, Y., Jardim, E., Spencer, P.D., Hanselman, D.H., Blaylock, J., Mood, M., Hulson, P.-J.F., 2015. Simulation testing the robustness of stock assessment models to error: Some results from the ICES strategic initiative on stock assessment methods. *ICES J Mar Sci* 72, 19–30. <https://doi.org/10.1093/icesjms/fst237>

Di Lorenzo, E., Ohman, M.D., 2013. A double-integration hypothesis to explain ocean ecosystem response to climate forcing. *Proceedings of the National Academy of Sciences* 110, 2496–2499. <https://doi.org/10.1073/pnas.1218022110>

Free, C.M., Thorson, J.T., Pinsky, M.L., Oken, K.L., Wiedenmann, J., Jensen, O.P., 2019. Impacts of historical warming on marine fisheries production. *Science* 363, 979–983. <https://doi.org/10.1126/science.aau1758>

Fry, F., 1971. The effect of environmental factors on the physiology of fish, in: *Fish Physiology*. Elsevier, pp. 1–98. [https://doi.org/10.1016/S1546-5098\(08\)60146-6](https://doi.org/10.1016/S1546-5098(08)60146-6)

Garstang, W., 1900. The Impoverishment of the Sea. A Critical Summary of the Experimental and Statistical Evidence bearing upon the Alleged Depletion of the Trawling Grounds. *Journal of the Marine Biological Association of the United Kingdom* 6, 1–69. <https://doi.org/10.1017/S0025315400072374>

Gudmundsson, G., 1994. Time series analysis of catch-at-age observations. *Applied Statistics* 43, 117–126.

Haltuch, M.A., Brooks, E., Brodziak, J., Devine, J., Johnson, K., Klibansky, N., Nash, R., Payne, M., Shertzer, K., Subbey, S., Wells, B., 2019. Unraveling the recruitment problem: A review of environmentally-informed forecasting and management strategy evaluation. *Fisheries Research* 217, 198–216. <https://doi.org/10.1016/j.fishres.2018.12.016>

Haltuch, M.A., Punt, A.E., 2011. The promises and pitfalls of including decadal-scale climate forcing of recruitment in groundfish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* 68, 912–926. <https://doi.org/10.1139/f2011-030>

Hare, J.A., Borggaard, D.L., Friedland, K.D., Anderson, J., Burns, P., Chu, K., Clay, P.M., Collins, M.J., Cooper, P., Fratantoni, P.S., Johnson, M.R., Manderson, J.P., Milke, L., Miller, T.J., Orphanides, C.D., Saba, V.S., 2016. Northeast Regional Action Plan - NOAA Fisheries Climate Science Strategy (No. NMFS-NE-239). NOAA Fisheries, Northeast Fisheries Science Center, Woods Hole, MA.

Hill, K.T., Crone, P.R., Zwolinski, J.P., 2018. Assessment of the Pacific sardine resource in 2018 for U.S.

Management in 2018-2019 (No. NOAA Technical Memorandum NMFS-SWFSC-600). US Department of  
Commerce.

Hjort, J., 1914. Fluctuations in the great fisheries of Northern Europe viewed in the light of biological  
research. *Rapports et Procès-Verbaux des Réunions du Conseil Permanent International Pour L'Exploration  
de la Mer* 20, 1–228.

ICES, 2020. Workshop on the review and future of state space stock assessment models in ICES (WKRFSAM).  
ICES Scientific Reports 2, 23p. <https://doi.org/10.17895/ices.pub.6004>

ICES, 2017a. Report of the working group on assessment of demersal stocks in the North Sea and Skagerrak  
(2017). ICES CM 2017/ACOM:21, 26 April-5 May 2017, ICES HQ.

ICES, 2017b. Report of the North Western Working Group (NWWG). ICES CM 2017/ACOM:08, 27 April-4  
May 2017, Copenhagen, Denmark.

Iles, T.C., Beverton, R.J.H., 1998. Stock, recruitment and moderating processes in flatfish. *Journal of  
Sea Research, Proceedings of the Third International Symposium on Flatfish Ecology, Part II* 39, 41–55.  
[https://doi.org/10.1016/S1385-1101\(97\)00022-1](https://doi.org/10.1016/S1385-1101(97)00022-1)

Kristensen, K., Nielsen, A., Berg, C., Skaug, H., Bell, B.M., 2016. TMB: Automatic differentiation and  
Laplace approximation. *Journal of Statistical Software* 70, 1–21. <https://doi.org/10.18637/jss.v070.i05>

Larkin, P., 1996. Concepts and issues in marine ecosystem management. *Reviews in Fish Biology and  
Fisheries* 6, 139–164. <https://doi.org/10.1007/BF00182341>

Legault, C.M., Restrepo, V.R., 1998. A Flexible Forward Age-Structured Assessment Program (No. 49).

Link, J.S., 2002. What Does Ecosystem-Based Fisheries Management Mean? *Fisheries* 27, 5.

Lorenzen, K., 1996. The relationship between body weight and natural mortality in juvenile and adult  
fish: A comparison of natural ecosystems and aquaculture. *Journal of Fish Biology* 49, 627–642. <https://doi.org/10.1111/j.1095-8649.1996.tb00060.x>

Lynch, P.D., Methot, R.D., Link, J.S. (Eds.), 2018. Implementing a Next Generation Stock Assessment  
Enterprise. An Update to the NOAA Fisheries Stock Assessment Improvement Plan, in: U.S. Dep. Commer.,  
NOAA Tech. Memo. NMFS-F/ SPO-183, p. 127. <https://doi.org/10.7755/TMSPO.183>

Marshall, K.N., Koehn, L.E., Levin, P.S., Essington, T.E., Jensen, O.P., 2019. Inclusion of ecosystem  
information in US fish stock assessments suggests progress toward ecosystem-based fisheries management.  
*ICES J Mar Sci* 76, 1–9. <https://doi.org/10.1093/icesjms/fsy152>



546 McClatchie, S., Goericke, R., Auad, G., Hill, K., 2010. Re-assessment of the stockRecruit and tempera-  
547 tureRecruit relationships for Pacific sardine (*Sardinops sagax*). *Can. J. Fish. Aquat. Sci.* 67, 1782–1790.  
548 <https://doi.org/10.1139/F10-101>

549 Mendelssohn, R., 1988. Some problems in estimating population sizes from catch-at-age data. *Fishery*  
550 *Bulletin* 86, 617–630.

551 Methot, R.D., Taylor, I.G., 2011. Adjusting for bias due to variability of estimated recruitments in fishery  
552 assessment models. *Can. J. Fish. Aquat. Sci.* 68, 1744–1760. <https://doi.org/10.1139/f2011-092>

553 Methot, R.D., Wetzel, C.R., 2013. Stock synthesis: A biological and statistical framework for fish stock  
554 assessment and fishery management. *Fisheries Research* 142, 86–99. [https://doi.org/10.1016/j.fishres.2012.10.](https://doi.org/10.1016/j.fishres.2012.10.012)  
555 012

556 Miller, T.J., Hare, J.A., Alade, L.A., 2016. A state-space approach to incorporating environmental effects on  
557 recruitment in an age-structured assessment model with an application to southern New England yellowtail  
558 flounder. *Canadian Journal of Fisheries and Aquatic Sciences* 73, 1261–1270. [https://doi.org/10.1139/cjfas-](https://doi.org/10.1139/cjfas-2015-0339)  
559 2015-0339

560 Miller, T.J., Hyun, S.-Y., 2018. Evaluating evidence for alternative natural mortality and process error  
561 assumptions using a state-space, age-structured assessment model. *Canadian Journal of Fisheries and Aquatic*  
562 *Sciences* 75, 691–703. <https://doi.org/10.1139/cjfas-2017-0035>

563 Miller, T.J., Legault, C.M., 2015. Technical details for ASAP version 4 (No. Ref Doc. 15-17). US Dept  
564 Commer, Northeast Fish Sci Cent.

565 Miller, T.J., O'Brien, L., Fratantoni, P.S., 2018. Temporal and environmental variation in growth and  
566 maturity and effects on management reference points of Georges Bank Atlantic cod. *Can. J. Fish. Aquat.*  
567 *Sci.* 1–13. <https://doi.org/10.1139/cjfas-2017-0124>

568 Miller, T.J., Stock, B.C., 2020. The Woods Hole Assessment Model (WHAM).

569 Möllmann, C., Lindegren, M., Blenckner, T., Bergström, L., Casini, M., Diekmann, R., Flinkman, J.,  
570 Müller-Karulis, B., Neuenfeldt, S., Schmidt, J.O., Tomczak, M., Voss, R., Gårdmark, A., 2014. Implementing  
571 ecosystem-based fisheries management: From single-species to integrated ecosystem assessment and advice  
572 for Baltic Sea fish stocks. *ICES J Mar Sci* 71, 1187–1197. <https://doi.org/10.1093/icesjms/fst123>

573 Munch, S.B., Giron-Nava, A., Sugihara, G., 2018. Nonlinear dynamics and noise in fisheries recruitment: A  
574 global meta-analysis. *Fish and Fisheries* 19, 964–973. <https://doi.org/10.1111/faf.12304>

Munch, S.B., Poynor, V., Arriaza, J.L., 2017. Circumventing structural uncertainty: A Bayesian perspective on nonlinear forecasting for ecology. *Ecological Complexity* 32, 134–143. <https://doi.org/10.1016/j.ecocom.2016.08.006>

Myers, R.A., 1998. When do environment-recruitment correlations work? *Reviews in Fish Biology and Fisheries* 8, 285–305.

NEFSC, 2020a. Operational assessment of 14 Northeast groundfish stocks, updated through 2018. U.S. Dept. Commer., NOAA, NMFS, NEFSC, Woods Hole, MA.

NEFSC, 2020b. Butterfish 2020 assessment update report. U.S. Dept. Commer., NOAA, NMFS, NEFSC, Woods Hole, MA.

Nielsen, A., Berg, C.W., 2014. Estimation of time-varying selectivity in stock assessments using state-space models. *Fisheries Research* 158, 96–101. <https://doi.org/10.1016/j.fishres.2014.01.014>

O’Leary, C.A., Miller, T.J., Thorson, J.T., Nye, J.A., 2019. Understanding historical summer flounder (*Paralichthys Dentatus*) abundance patterns through the incorporation of oceanography-dependent vital rates in Bayesian hierarchical models. *Can. J. Fish. Aquat. Sci.* 76, 1275–1294. <https://doi.org/10.1139/cjfas-2018-0092>

Patrick, W.S., Link, J.S., 2015. Myths that Continue to Impede Progress in Ecosystem-Based Fisheries Management. *Fisheries* 40, 155–160. <https://doi.org/10.1080/03632415.2015.1024308>

Perretti, C.T., Fogarty, M.J., Friedland, K.D., Hare, J.A., Lucey, S.M., McBride, R.S., Miller, T.J., Morse, R.E., O’Brien, L., Pereira, J.J., Smith, L.A., Wuenschel, M.J., 2017. Regime shifts in fish recruitment on the Northeast US Continental Shelf. *Marine Ecology Progress Series* 574, 1–11. <https://doi.org/10.3354/meps12183>

Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Bris, A.L., Mills, K.E., Nye, J.A., Record, N.R., Scannell, H.A., Scott, J.D., Sherwood, G.D., Thomas, A.C., 2015. Slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery. *Science* 350, 809–812. <https://doi.org/10.1126/science.aac9819>

Punt, A.E., A’mar, T., Bond, N.A., Butterworth, D.S., de Moor, C.L., De Oliveira, J.A.A., Haltuch, M.A., Hollowed, A.B., Szuwalski, C., 2014. Fisheries management under climate and environmental uncertainty: Control rules and performance simulation. *ICES J Mar Sci* 71, 2208–2220. <https://doi.org/10.1093/icesjms/fst057>

R Core Team, 2020. R: A Language and Environment for Statistical Computing.

605 Rose, G.A., Rowe, S., 2015. Northern cod comeback. *Can. J. Fish. Aquat. Sci.* 72, 1789–1798. <https://doi.org/10.1139/cjfas-2015-0346>

607 Shelton, P.A., Sinclair, A.F., Chouinard, G.A., Mohn, R., Duplisea, D.E., 2006. Fishing under low productivity  
608 conditions is further delaying recovery of Northwest Atlantic cod (*Gadus morhua*). *Can. J. Fish. Aquat. Sci.*  
609 63, 235–238. <https://doi.org/10.1139/f05-253>

610 Shotwell, S.K., Hanselman, D.H., Belkin, I.M., 2014. Toward biophysical synergy: Investigating advection  
611 along the Polar Front to identify factors influencing Alaska sablefish recruitment. *Deep Sea Research Part II: Topical Studies in Oceanography* 107, 40–53. <https://doi.org/10.1016/j.dsr2.2012.08.024>

613 Stock, B.C., Xu, H., Miller, T.J., Thorson, J.T., Nye, J.A., n.d. Implementing a 2-dimensional smoother on  
614 either survival or natural mortality improves a state-space assessment model for Southern New England-Mid  
615 Atlantic yellowtail flounder.

616 Stock, C.A., Alexander, M.A., Bond, N.A., Brander, K.M., Cheung, W.W., Curchitser, E.N., Delworth, T.L.,  
617 Dunne, J.P., Griffies, S.M., Haltuch, M.A., Hare, J.A., Hollowed, A.B., Lehodey, P., Levin, S.A., Link, J.S.,  
618 Rose, K.A., Rykaczewski, R.R., Sarmiento, J.L., Stouffer, R.J., Schwing, F.B., Vecchi, G.A., Werner, F.E.,  
619 2011. On the use of IPCC-class models to assess the impact of climate on Living Marine Resources. *Progress*  
620 *in Oceanography* 88, 1–27. <https://doi.org/10.1016/j.pocean.2010.09.001>

621 Sullivan, P.J., 1992. A Kalman filter approach to catch-at-length analysis. *Biometrics* 48, 237–257.

622 Tableau, A., Collie, J.S., Bell, R.J., Minto, C., 2018. Decadal changes in the productivity of New England  
623 fish populations. *Can. J. Fish. Aquat. Sci.* 76, 1528–1540. <https://doi.org/10.1139/cjfas-2018-0255>

624 Thompson, W.R., 1936. On Confidence Ranges for the Median and Other Expectation Distributions for  
625 Populations of Unknown Distribution Form. *Ann. Math. Statist.* 7, 122–128. <https://doi.org/10.1214/aoms/1177732502>

627 Thorson, J.T., 2019. Perspective: Let’s simplify stock assessment by replacing tuning algorithms with  
628 statistics. *Fisheries Research* 217, 133–139. <https://doi.org/10.1016/j.fishres.2018.02.005>

629 Walters, C.J., Collie, J.S., 1988. Is Research on Environmental Factors Useful to Fisheries Management?  
630 *Can. J. Fish. Aquat. Sci.* 45, 1848–1854. <https://doi.org/10.1139/f88-217>

631 Winemiller, K.O., Rose, K.A., 1992. Patterns of Life-History Diversification in North American Fishes:  
632 Implications for Population Regulation. *Canadian Journal of Fisheries and Aquatic Sciences* 49, 2196–2218.  
633 <https://doi.org/10.1139/f92-242>

- 634 Xu, H., Miller, T.J., Hameed, S., Alade, L.A., Nye, J.A., 2018. Evaluating the utility of the Gulf Stream Index  
635 for predicting recruitment of Southern New England-Mid Atlantic yellowtail flounder. *Fisheries Oceanography*  
636 27, 85–95. <https://doi.org/10.1111/fog.12236>
- 637 Xu, H., Thorson, J.T., Methot, R.D., 2020. Comparing the performance of three data weighting methods when  
638 allowing for time-varying selectivity. *Can. J. Fish. Aquat. Sci.* 77, 247–263. [https://doi.org/10.1139/cjfas-](https://doi.org/10.1139/cjfas-2019-0107)  
639 2019-0107
- 640 Xu, H., Thorson, J.T., Methot, R.D., Taylor, I.G., 2019. A new semi-parametric method for autocorrelated  
641 age- and time-varying selectivity in age-structured assessment models. *Can. J. Fish. Aquat. Sci.* 76, 268–285.  
642 <https://doi.org/10.1139/cjfas-2017-0446>
- 643 Zwolinski, J.P., Demer, D.A., 2012. A cold oceanographic regime with high exploitation rates in the Northeast  
644 Pacific forecasts a collapse of the sardine stock. *Proceedings of the National Academy of Sciences* 109,  
645 4175–4180. <https://doi.org/10.1073/pnas.1113806109>

Table 1: Model descriptions and estimated parameters. Parameter descriptions and equations are given in text. Note that the base model in the  $M$  module is NAA m1, and the base model in the Selectivity and Ecov-Recruitment modules is NAA m3. Ecov m1 fits the Cold Pool Index data and estimates  $\sigma_x$  in order to allow comparison to m2-m5 using AIC (same data needed in likelihood).

| Model                                     | Description   | Estimated parameters                                     |
|---|---|--|
| <b>Numbers-at-age (NAA)</b>               |   |  |
| m1: SCAA (IID)                            | Recruitment deviations are IID random effects                                       | $\sigma_R$   |
| m2: SCAA (AR1)                            | Recruitment deviations are autocorrelated (AR1) random effects                      | $\sigma_R, \rho_y$                                       |
| m3: NAA (IID)                             | All NAA deviations are IID random effects   | $\sigma_R, \sigma_a$                                     |
| m4: NAA (2D AR1)                          | All NAA deviations are random effects with correlation by year and age (2D AR1)     | $\sigma_R, \sigma_a, \rho_y, \rho_a$                     |
| <b>Natural mortality (<math>M</math>)</b> |   |  |
| m1: none                                  | No random effects on $M$  | $\sigma_R$   |
| m2: IID                                   | $M$ deviations are IID random effects   | $\sigma_R, \sigma_M$                                     |
| m3: 2D AR1                                | $M$ deviations are random effects with correlation by year and age (2D AR1)         | $\sigma_R, \sigma_M, \varphi_y, \varphi_a$               |
| <b>Selectivity (Sel)</b>                  |   |  |
| m1: none                                  | No random effects on selectivity  | $\sigma_R, \sigma_a$                                     |
| m2: IID                                   | Selectivity deviations are IID random effects                                       | $\sigma_R, \sigma_a, \sigma_{Sel}$                       |
| m3: 2D AR1                                | Selectivity deviations are random effects with correlation by year and age (2D AR1) | $\sigma_R, \sigma_a, \sigma_{Sel}, \phi_y, \phi_a$       |
| <b>Ecov-Recruitment (Ecov)</b>            |   |  |
| m1: RW-none                               | Ecov: random walk (RW), effect on $\beta$ : none                                    | $\sigma_R, \sigma_a, \sigma_x$                           |
| m2: RW-linear                             | Ecov: random walk (RW), effect on $\beta$ : linear                                  | $\sigma_R, \sigma_a, \sigma_x, \beta_1$                  |
| m3: RW-poly                               | Ecov: random walk (RW), effect on $\beta$ : 2nd order polynomial (poly)             | $\sigma_R, \sigma_a, \sigma_x, \beta_1, \beta_2$         |
| m4: AR1-linear                            | Ecov: autocorrelated (AR1), effect on $\beta$ : linear                              | $\sigma_R, \sigma_a, \sigma_x, \phi_x, \beta_1$          |
| m5: AR1-poly                              | Ecov: autocorrelated (AR1), effect on $\beta$ : 2nd order polynomial (poly)         | $\sigma_R, \sigma_a, \sigma_x, \phi_x, \beta_1, \beta_2$ |

Table 2: Stocks used in simulation tests.

| Stock                     | Modules tested |   |     |      | Model dim |      | Biol. par. |       | Stock status |            | Source |                    |                    |
|---------------------------|----------------|---|-----|------|-----------|------|------------|-------|--------------|------------|--------|--------------------|--------------------|
|                           | NAA            | M | Sel | Ecov | #         | Ages | #          | Years | $M$          | $\sigma_R$ |        | $\frac{B}{B_{40}}$ | $\frac{F}{F_{40}}$ |
| SNEMA yellowtail flounder | x              | x |     | x    |           | 6    |            | 49    | 0.2-0.4      | 1.67       | 0.01   | 0.44               | NEFSC (2020a)      |
| Butterfish                | x              | x |     |      |           | 5    |            | 31    | 1.3          | 0.23       | 2.57   | 0.03               | NEFSC (2020b)      |
| North Sea cod             | x              | x |     |      |           | 6    |            | 54    | 0.2-1.2      | 0.87       | 0.14   | 2.00               | ICES (2017a)       |
| Icelandic herring         | x              |   |     |      |           | 11   |            | 30    | 0.1          | 0.55       | 0.40   | 1.81               | ICES (2017b)       |
| Georges Bank haddock      | x              |   | x   |      |           | 9    |            | 86    | 0.2          | 1.65       | 5.16   | 0.12               | NEFSC (2020a)      |




Figure 1: AIC differences by model and stock when fit to original datasets. Stock abbreviations: SNEMA yellowtail flounder (SNEMAYT), North Sea cod (NScod), Icelandic herring (ICEherring), and Georges Bank haddock (GBhaddock).




Figure 2: Abundanc-at-age deviations estimated for Icelandic herring using four models of numbers-at-age (NAA) random effects. m1 = only recruitment deviations are random effects (most similar to traditional statistical catch-at-age, SCAA), and deviations are independent and identically distributed (IID). m2 = as m1, but with autocorrelated recruitment deviations (AR1<sub>y</sub>). m3 = all NAA deviations are IID random effects. m4 = as m3, but deviations are correlated by age and year (2D AR1).






Figure 3: Natural mortality ( $M$ ) estimated for butterfish using three random effects models. m1 = no random effects on  $M$ . m2 =  $M$  deviations are independent and identically distributed (IID). m3 =  $M$  deviations are correlated by age and year (2D AR1).




Figure 4: Selectivity estimated for Georges Bank haddock using three random effects models. m1 = no random effects (constant logistic selectivity). m2 = selectivity deviations are independent and identically distributed (IID). m3 = selectivity deviations are correlated by parameter and year (2D AR1).




Figure 5: Beverton-Holt stock-recruit relationships fit for Southern New England-Mid Atlantic yellowtail flounder, with and without effects of the Cold Pool Index (CPI). A) CPI estimated from the model with lowest AIC (m4, AR1-linear). Points are observations with 95% CI, and the line with shading is the model-estimated CPI with 95% CI. Note the increased uncertainty surrounding the CPI estimate in 2017 (no observation). B) Estimates of spawning stock biomass (SSB), recruitment, and the stock-recruit function from the model without a CPI effect, m1. C) Estimates of SSB and recruitment from m4, with an effect of the CPI on  $\beta$ . Lines depict the expected stock-recruit relationship in each year  $t$ , given the CPI in year  $t - 1$  (color).




Figure 6: Proportion of simulations in which each model had the lowest AIC. A) Numbers-at-age (NAA), aggregated across all five stocks. B) Natural mortality ( $M$ ), aggregated over two stocks (SNEMAYT and NScod). C) Selectivity (GBhaddock). Not all estimation models converged for each simulation, even when the operating model matched.




Figure 7: Relative error of parameters constraining numbers-at-age (NAA) random effects. Four models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. m1 = only recruitment deviations are random effects (most similar to traditional statistical catch-at-age, SCAA), and deviations are independent and identically distributed (IID). m2 = as m1, but with autocorrelated recruitment deviations (AR1<sub>y</sub>). m3 = all NAA deviations are IID random effects. m4 = as m3, but deviations are correlated by age and year (2D AR1). Relative error was calculated as  $\frac{\hat{\theta}_i}{\theta} - 1$ , where  $\hat{\theta}_i$  was the estimate in simulation  $i$  for parameter  $\theta$ , and  $\theta$  was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI.




Figure 8: Relative error of key quantities estimated for Icelandic herring using four models of numbers-at-age (NAA) random effects. m1 = only recruitment deviations are random effects (most similar to traditional statistical catch-at-age, SCAA), and deviations are independent and identically distributed (IID). m2 = as m1, but with autocorrelated recruitment deviations (AR1<sub>y</sub>). m3 = all NAA deviations are IID random effects. m4 = as m3, but deviations are correlated by age and year (2D AR1).




Figure 9: Relative error of parameters constraining natural mortality ( $M$ ) random effects. Three models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. m1 = no random effects on  $M$ . m2 =  $M$  deviations were independent and identically distributed (IID). m3 =  $M$  deviations were correlated by age and year (2D AR1). Relative error was calculated as  $\frac{\hat{\theta}_i}{\theta} - 1$ , where  $\hat{\theta}_i$  was the estimate in simulation  $i$  for parameter  $\theta$ , and  $\theta$  was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI. Stock abbreviations: SNEMAYT yellowtail flounder (SNEMAYT) and North Sea cod (NScod, m3 did not converge).




Figure 10: Relative error of key quantities estimated for butterfish using three models of natural mortality ( $M$ ) random effects. m1 = no random effects on  $M$ . m2 =  $M$  deviations are independent and identically distributed (IID). m3 =  $M$  deviations are correlated by age and year (2D AR1).






Figure 11: Relative error of parameters constraining selectivity random effects for Georges Bank haddock (GBhaddock). Three models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. m1 = no random effects (constant selectivity). m2 = selectivity deviations were independent and identically distributed (IID). m3 = selectivity deviations were correlated by parameter and year (2D AR1). Relative error was calculated as  $\frac{\hat{\theta}_i}{\theta} - 1$ , where  $\hat{\theta}_i$  was the estimate in simulation  $i$  for parameter  $\theta$ , and  $\theta$  was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI.




Figure 12: Relative error of key quantities estimated for Georges Bank haddock using three models of selectivity random effects. m1 = no random effects (constant logistic selectivity). m2 = selectivity deviations are independent and identically distributed (IID). m3 = selectivity deviations are correlated by parameter and year (2D AR1).




Figure 13: Relative error of parameters constraining variation in recruitment for Southern New England-Mid Atlantic yellowtail flounder (SNEMAYT). Five models were used to simulate 100 datasets keeping fixed effect parameters constant, and then re-fit to each simulated dataset. All models estimated recruitment using the Beverton-Holt function and included CPI effects on  $\beta$ :  $\hat{R}_{t+1} = \frac{\alpha S_t}{1 + e^{\beta_0 + \beta_1 x_t + \beta_2 x_t^2 S_t}}$ . m1 = Cold Pool Index (CPI) modeled as a random walk (RW) with no effect on recruitment ( $\beta_1 = \beta_2 = 0$ ). m2 = CPI as RW, linear effect on  $\beta$ . m3 = CPI as RW, 2nd order polynomial effect on  $\beta$ . m4 = CPI as AR1, linear effect. m5 = CPI as AR1, polynomial effect. Relative error was calculated as  $\frac{\hat{\theta}_i}{\theta} - 1$ , where  $\hat{\theta}_i$  was the estimate in simulation  $i$  for parameter  $\theta$ , and  $\theta$  was the true value (estimate from original dataset). Red points and lines show median relative error with 95% CI.