

Cartesian Coordinates

BRIAN ZHANG

August 23, 2022

§1 Theorems

Theorem 1.1 (Distance Formula)

The distance between points (x_1, y_1) and (x_2, y_2) is

$$\sqrt{((x_1 - x_2)^2 + (y_1 - y_2)^2)}.$$

Theorem 1.2 (Point slope form)

The equation of the line with slope m that passes through point (a, b) is

$$y - b = m(x - a).$$

Theorem 1.3 (Parallel lines)

Two lines are parallel if and only if their slope is the same.

Theorem 1.4 (Perpendicular lines)

Two lines with slopes m and n are perpendicular if and only if $m \cdot n = -1$.

Theorem 1.5 (Circle)

The circle with center (h, k) and radius r has equation

$$(x - h)^2 + (y - k)^2 = r^2.$$

Theorem 1.6 (Point to Line)

The distance between the point (x_0, y_0) to the line $Ax + By + C = 0$ can be written as

$$\frac{|Ax_0 + By_0 + C|}{\sqrt{A^2 + B^2}}.$$

Theorem 1.7 (Shoelace)

In a polygon with coordinates $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$, the area \mathbf{A} is:

$$\begin{aligned}\mathbf{A} &= \frac{1}{2} \left| \left(\sum_{i=1}^{n-1} x_i y_{i+1} \right) + x_n y_1 - \left(\sum_{i=1}^{n-1} x_{i+1} y_i \right) - x_1 y_n \right| \\ &= \frac{1}{2} \cdot |x_1 y_2 + x_2 y_3 + \dots + x_{n-1} y_n + x_n y_1 - x_2 y_1 - x_3 y_2 - \dots - x_n y_{n-1} - x_1 y_n|.\end{aligned}$$

Theorem 1.8 (Midpoint)

The midpoint of points (x_1, y_1) and (x_2, y_2) is

$$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right).$$

Theorem 1.9 (Coordinates of the Centroid)

In a triangle with coordinates $A = (x_a, y_a), B = (x_b, y_b), C = (x_c, y_c)$, the coordinates of the centroid is

$$\left(\frac{x_a + x_b + x_c}{3}, \frac{y_a + y_b + y_c}{3} \right).$$

§2 Heuristics

Here are a couple of tips and some philosophy about coordinate bashes.

1. Many people say that coordinate bashing is a “no brain” technique, and is for when you don’t know how to synthetic the geometry. This is false. When coordinate bashing, the most important step is your set up. This means what you define as the origin, and the equations of circles, lines, etc. Having the right setup will save you lots of time.
2. Have a plan. Whenever I use coordinates on a problem, I plan out how I would calculate each point, and the method of calculating each point. This helps you get a sense of how long it will take you to solve the problem, and whether it’s worth it to just move onto another problem, or to try to look for a synthetic solution.
3. Know when to apply coordinates! Right angles, lots of lines, problems where you are given triangle side lengths are your friend. Some things that are harder to deal with are incenters, multiple circles, etc.
4. Angle conditions are usually pretty hard to deal with, but if they can be reduced to the angle bisector theorem, cyclic quadrilaterals, or tangents, then it is much easier to deal with them.
5. When you have a nice central circle, it’s a good idea to set that circle at the origin, so the equation of the circle is easier to work with.

6. For obvious reasons, knowledge of standard geometry theorems is also very helpful. If you can use synthetic techniques to help you simplify the problem, then coordinate bashing can be much less computationally heavy and much faster.
7. When given a triangle with three side lengths, then you can use Heron's to calculate the area, and then calculate the altitude of one of the sides, so you can place the triangle onto the coordinate plane.

§3 Examples

Example 3.1 (MATHCOUNTS 2008)

Triangle $\triangle ABC$ is a right triangle with $\angle C = 90^\circ$ and $AC = 4$ and $BC = 7$. Points E and F are on sides CA and CB , respectively with $CE = 2$ and $CF = 3$. Given that D is the intersection of lines AF and BE , and the area of $\triangle ABD$ can be written as $\frac{m}{n}$ with $\gcd(m, n) = 1$, then find the value of $m + n$.

Walkthrough:

- (a) Let C be the origin. Then, what are the coordinates of E and F ?
- (b) Find the equations of lines AF and BE , and then find their intersection point. This is point D .
- (c) Finish by finding the area of ABD with an area formula.

Example 3.2 (AMC 10A 2020/20)

Quadrilateral $ABCD$ satisfies $\angle ABC = \angle ACD = 90^\circ$, $AC = 20$, and $CD = 30$. Diagonals \overline{AC} and \overline{BD} intersect at point E , and $AE = 5$. What is the area of quadrilateral $ABCD$?

Walkthrough:

- (a) We could set the origin to be C , as we have CD and AC , but there actually is a better point to use. (If you can't figure out where this is, then think about how we know AC , and $\angle ABC = 90^\circ$.)
You should have set the origin to be the midpoint of segment AC . We are motivated to do this because this allows us to draw a circle ω centered at the origin that passes through A and C , and note that B also passes through the circle.
- (b) Find the equation to the circle ω , and calculate B by defining it as the second intersection of ω with DE .
- (c) You should get a quadratic. Eliminate the "wrong" solution.
- (d) Finish.

Example 3.3 (AMC 10B 2009/18)

Let rectangle $ABCD$ have side lengths $AB = 8$ and $BC = 6$. Let M be the midpoint of diagonal \overline{AC} , and E be on AB with $\overline{ME} \perp \overline{AC}$. What is the area of $\triangle AME$?

Walkthrough:

- (a) Let A be the origin. Then, what are the coordinates of the other points?
- (b) We need to use the perpendicular condition to write an equation about point E . If we let E be $(e, 0)$, then what equation can we write?
- (c) You should find that E is $(\frac{25}{4}, 0)$. Use this to find the desired area.

Example 3.4 (AMC 10A 2016/19)

In rectangle $ABCD$, $AB = 6$ and $BC = 3$. Point E between B and C , and point F between E and C are such that $BE = EF = FC$. Segments \overline{AE} and \overline{AF} intersect \overline{BD} at P and Q , respectively. The ratio $BP : PQ : QD$ can be written as $r : s : t$, where the greatest common factor of r, s and t is 1. What is $r + s + t$?

Walkthrough:

- (a) Let B be the origin. Then, what are the equations for lines BD , AE , and AF ?
- (b) Calculate the locations of points P and Q by solving for the intersections of the lines.
- (c) Find the desired ratio by comparing the x -coordinates of B , P , Q , and D .
- (d) Why do we only have to look at the x -coordinate?

Example 3.5 (AIME I 2021/9)

Let $ABCD$ be an isosceles trapezoid with $AD = BC$ and $AB < CD$. Suppose that the distances from A to the lines BC, CD , and BD are 15, 18, and 10, respectively. Let K be the area of $ABCD$. Find $\sqrt{2} \cdot K$.

Walkthrough:

- (a) Note how we have a bunch of distances from a point to several lines, with the point being on one of 2 parallel lines. What are we motivated to use?
- (b) Find a nice point that we can set as the origin, so that the coordinates of A, B, C, D are all relatively clean.

What I found to be the cleanest was to set the foot of the altitude from A to CD to be the origin, although some other solutions used different origins.

- (c) Use point to line from A to BC and BD to get two equations.

You should have gotten the quadratics

$$\begin{aligned}\sqrt{324 + (b + d)^2} &= \frac{9}{5}b \\ \sqrt{324 + d^2} &= \frac{6}{5}b\end{aligned}$$


or some equivalent expression. This may be slightly different based on how you chose your variables and the origin.

- (d) Solve for b and d , by squaring and subtracting.
- (e) Finish.


§4 Problems

Problem 4.1 (AMC 10B 2004/18). In the right triangle $\triangle ACE$, we are given that $AC = 12$, $CE = 16$, and $EA = 20$. Let the points B , D , and F be located on AC , CE , and EA , respectively, so that $AB = 3$, $CD = 4$, and $EF = 5$. If the value of $\frac{[\triangle DBF]}{[\triangle ACE]}$ is equal to $\frac{m}{n}$, where $\gcd(m, n) = 1$, then find the value of $m + n$.

Problem 4.2 (AMC 10A 2018/23). Farmer Kevin has a field of corn in the shape of a right triangle. The right triangle's legs have lengths 3 and 4 units. In the corner where those sides meet at a right angle, he leaves a small unplanted square S so that from the air it looks like the right angle symbol. The rest of the field is planted. The shortest distance from S to the hypotenuse is 2 units. Given that the fraction of the field that is planted can be written as $\frac{m}{n}$ where $\gcd(m, n) = 1$, then find the value of $m + n$.



Problem 4.3 (AMC 10A 2014/16). In rectangle $ABCD$, $AB = 1$, $BC = 2$, and points E , F , and G are midpoints of \overline{BC} , \overline{CD} , and \overline{AD} , respectively. Point H is the midpoint of \overline{GE} . What is the area of the shaded region?




Problem 4.4 (CIME I 2021/5). In rectangle $ABCD$, suppose $AD = 20$ and $AB = 21$. The circle centered at A passing through D intersects the circle centered at C passing through D at a point $P \neq D$. Then the length BP can be written in the form $\frac{p}{q}$, where p and q are relatively prime positive integers. Find $p + q$.

Problem 4.5 (AIME I 2015/4). Point B lies on line segment \overline{AC} with $AB = 16$ and $BC = 4$. Points D and E lie on the same side of line AC forming equilateral triangles $\triangle ABD$ and $\triangle BCE$. Let M be the midpoint of \overline{AE} , and N be the midpoint of \overline{CD} . The area of $\triangle BMN$ is x . Find x^2 .

Problem 4.6 (AIME II 2018/9). Octagon $ABCDEFGH$ with side lengths $AB = CD = EF = GH = 10$ and $BC = DE = FG = HA = 11$ is formed by removing four $6-8-10$ triangles from the corners of a 23×27 rectangle with side \overline{AH} on a short side of the

rectangle, as shown. Let J be the midpoint of \overline{HA} , and partition the octagon into 7 triangles by drawing segments \overline{JB} , \overline{JC} , \overline{JD} , \overline{JE} , \overline{JF} , and \overline{JG} . Find the area of the convex polygon whose vertices are the centroids of these 7 triangles.



Problem 4.7 (AIME II 2017/10). Rectangle $ABCD$ has side lengths $AB = 84$ and $AD = 42$. Point M is the midpoint of \overline{AD} , point N is the trisection point of \overline{AB} closer to A , and point O is the intersection of \overline{CM} and \overline{DN} . Point P lies on the quadrilateral $BCON$, and \overline{BP} bisects the area of $BCON$. Find the area of $\triangle CDP$.

Problem 4.8 (AIME II 2003/11). Triangle ABC is a right triangle with $AC = 7$, $BC = 24$, and right angle at C . Point M is the midpoint of AB , and D is on the same side of line AB as C so that $AD = BD = 15$. Given that the area of triangle CDM may be expressed as $\frac{m\sqrt{n}}{p}$, where m , n , and p are positive integers, m and p are relatively prime, and n is not divisible by the square of any prime, find $m + n + p$.

§A Solutions to Examples

§A.1 Mathcounts

Let $C = (0, 0)$, $A = (0, 4)$, $B = (7, 0)$, $E = (0, 2)$, and $F = (3, 0)$. Then, we have line AF be equal to $y = -\frac{4}{3}x + 4$ and BE be $y = -\frac{2}{7}x + 2$. These two equations have solution $(\frac{21}{11}, \frac{16}{11})$. Then, we may apply shoelace to get an answer of $\boxed{\frac{56}{11}}$.

§A.2 AMC 10A 2020/20

We set the origin to be the midpoint of AC . Now, construct (ABC) , and note that the equation of (ABC) is $x^2 + y^2 = 100$. We have the coordinates as

$$A = (-10, 0)$$

$$C = (10, 0)$$

$$D = (10, 30)$$

$$E = (-5, 0).$$

It remains to calculate B . Now, note that B is simply the intersection of $x^2 + y^2 = 100$ and line DE . But line DE can be easily calculated to be $y = -2(x + 5)$ from point slope formula. It remains to solve

$$\begin{aligned} x^2 + (-2(x + 5))^2 &= 100 \\ \implies 5x^2 + 40x &= 0 \\ \implies x &= -8, 0 \end{aligned}$$

Obviously, $x = 0$ is the first intersection of DE with (ABC) , so we can toss that out. Then, plugging $x = -4$ back in we get $y = -6$. Finally, we finish with shoelace on $ABCD$, and get an answer of $\boxed{360}$.




Diagram from AoPS Wiki.

§A.3 AIME I 2021/9

Instead of taking the effort to find similar triangles, we use the method of coordinate bashing. We do this by setting the origin to be the foot from A to CD , and noting that

we can use point to line formula and isosceles trapezoid properties to get the locations of the points B, C, D .

We let X, Y, Z be the feet of the altitudes from A to CD, DB , and BC , respectively.

Now, setting X as the origin, we can set

$$\begin{aligned} A &= (0, 18) \\ B &= (b, 18) \\ C &= (b + d, 0) \\ D &= (-d, 0) \\ X &= (0, 0) \end{aligned}$$

We want to use point to line, so we need the equations of BC and BD . It's pretty easy to get

$$\begin{aligned} BD : y &= \frac{18}{b + d}(x + d) \\ BC : y &= \frac{18}{-d}(x - b - d) \end{aligned}$$

Now, point to line on A to BD and BC gives

$$\begin{aligned} 10 &= \frac{-18(b + d) + 18d}{\sqrt{324 + (b + d)^2}} \\ 15 &= \frac{18d + 18(-b - d)}{\sqrt{18^2 + d^2}} \end{aligned}$$

respectively. Now, rearranging, we get

$$\begin{aligned} \sqrt{324 + (b + d)^2} &= \frac{9}{5}b \\ \sqrt{324 + d^2} &= \frac{6}{5}b \end{aligned}$$

This is equivalent to

$$\begin{aligned} 324 + b^2 + 2bd + d^2 &= \frac{81}{25}b^2 \\ 324 + d^2 &= \frac{36}{25}b^2 \end{aligned}$$

Upon subtracting the two equations, we get $d = \frac{2}{5}b$. Now plugging this back into $324 + d^2 = \frac{36}{25}b^2$, we get

$$\begin{aligned} 324 + \frac{4}{25}b^2 &= \frac{36}{25}b^2 \\ 324 &= \frac{32}{25}b^2 \\ b &= \frac{45\sqrt{2}}{4} \\ d &= \frac{9\sqrt{2}}{2}. \end{aligned}$$

From here it is not hard to get the answer of $\boxed{567}$.

Remark A.1 ((Brian Zhang)). On contest, I calculated the area of the trapezoid to be $\frac{1}{2} \cdot (b + d) \cdot b \cdot 18 = 486$. Oops.

§A.4 AMC 10B 2009/18

Let $A = (0, 0)$, $B = (8, 0)$, $C = (8, 6)$, and $D = (0, 6)$. Then M is $(4, 3)$. If the coordinates of E is $(e, 0)$, then the perpendicular condition implies that

$$\frac{3}{4-e} \cdot \frac{3}{4} = -1 \implies e = \frac{25}{4}.$$

Thus, the area of $\triangle AME$ is $\frac{1}{2} \cdot \frac{25}{4} \cdot 3 = \boxed{\frac{75}{8}}$.

§A.5 AMC 10A 2016/19

We let $A = (6, 0)$, $B = (0, 0)$, $C = (0, 3)$, and $D = (6, 3)$. Then, E and F will be $(0, 1)$ and $(0, 2)$, respectively. It is easy to see that

$$\text{Line } BD: y = \frac{1}{2}x$$

$$\text{Line } AE: y = -\frac{1}{6}x + 1$$

$$\text{Line } AF: y = -\frac{1}{3}x + 2$$

Thus, we may calculate the locations of points P and Q by intersecting the two equations, so $P = (\frac{3}{2}, \frac{3}{4})$ and $Q = (\frac{12}{5}, \frac{6}{5})$. Thus, the desired ratio is

$$BP : PQ : QD = \frac{3}{2} : \frac{9}{10} : \frac{18}{5} = 5 : 3 : 12,$$

hence our answer is $\boxed{20}$.