

¹ bw_timex: A Python Package for Time-Explicit Life Cycle Assessment

³ **Timo Diepers**  ¹, **Amelie Müller**  ^{2,3}, and **Arthur Jakobs**  ⁴

⁴ 1 Institute of Technical Thermodynamics, RWTH Aachen University, Germany 2 Institute of Environmental Sciences (CML), Leiden University, The Netherlands 3 Flemish Institute for Technology Research (VITO), EnergyVille, Belgium 4 Technology Assessment Group, Laboratory for Energy Analysis, Center for Nuclear Engineering and Sciences & Center for Energy and Environmental Sciences, Paul Scherrer Institut (PSI), Villigen PSI, Switzerland

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

⁹ Summary

¹⁰ bw_timex is a Python package for time-explicit Life Cycle Assessment (LCA). Unlike conventional LCA, time-explicit LCA allows the quantification of environmental impacts of products and processes *over time*, considering their temporal distribution and evolution. As such, bw_timex enables simultaneously accounting for:

- the timing of processes throughout the supply chain (e.g., end-of-life treatment occurs 20 years after production),
- variable and/or evolving supply chains and technologies (e.g., increasing shares of renewable electricity or higher process efficiencies in the future), and
- the timing of emissions (enabling dynamic characterization).

To achieve this, bw_timex uses graph traversal to convolve process-relative temporal distributions through the supply chain. From the resulting timeline of technosphere exchanges, Life Cycle Inventories (LCIs) are automatically linked across time-specific background databases. The resulting time-explicit LCI reflects the current technology status within the product system at the actual time of each process. Moreover, bw_timex preserves the timing of emissions, enabling both dynamic and static Life Cycle Impact Assessment.

²⁵ Statement of need

²⁶ LCA traditionally assumes a static system, where all processes occur simultaneously and do not change over time ([Heijungs & Suh, 2002](#)). To add a temporal dimension to LCA, the fields of dynamic LCA (dLCA) and prospective LCA (pLCA) have emerged. While dLCA focuses on when processes and emissions occur and how impacts are distributed over time (*temporal distribution*), it typically assumes that the underlying product system remains the same ([Beloin-Saint-Pierre et al., 2020](#)). Conversely, while pLCA tracks how processes evolve (*temporal evolution*) using future scenarios, it generally only assesses a single (future) point in time, ignoring that processes occur at different times across a product's life cycle ([Arvidsson et al., 2024](#)).

³⁵ bw_timex provides a framework for time-explicit LCA calculations within the Brightway ecosystem ([Mutel, 2017](#)). It combines considerations of temporal distribution and evolution by accounting for both the timing of processes and emissions as well as the state of the product system at the respective points in time. This makes bw_timex particularly useful for studies involving variable or strongly evolving product systems, long-lived products, biogenic carbon, and scenario analyses.

41 State of the field

42 Existing dLCA tools such as Temporalis (Cardellini et al., 2018) handle temporal distributions
 43 but not system evolution. Conversely, pLCA tools like premise (Sacchi et al., 2022), Futura
 44 (Joyce & Björklund, 2022), and pathways (Sacchi & Hahn-Menacho, 2024) model evolving
 45 systems but not temporal distributions within the supply chain. Two recent tools combine
 46 both temporal distribution and evolution: ProsperDyn (Lang-Quantzendorff & Beermann,
 47 2025) and TRAILS (Sacchi, 2026). ProsperDyn is presently provided as a collection of research
 48 notebooks with limited documentation and without a consolidated, performance-oriented
 49 software architecture suitable for broader reuse. TRAILS, although methodologically advanced,
 50 currently relies on annual discretization and sequential year-specific calculations rather than a
 51 unified matrix-based integration of both dimensions.

52 bw_timex uniquely embeds the time dimension directly into the technosphere and biosphere
 53 matrices, enabling flexible temporal resolution within a single matrix-based framework. This
 54 allows efficient computation and seamless integration with the broader Brightway ecosystem.

55 Workflow

56 A time-explicit LCA with bw_timex follows four main steps, as illustrated in Figure 1. First,
 57 a conventional product system model is temporalized by adding process-relative temporal
 58 distributions (rTDs) to the exchanges (cf. Cardellini et al. (2018)). These rTDs describe how
 59 the amount of a technosphere or biosphere exchange is distributed over time, relative to the
 60 consuming or emitting process. In addition, temporal evolution of foreground processes can be
 61 defined through time-specific parameters. In Step 2, a timeline of technosphere exchanges is
 62 constructed by convolving rTDs along the supply chain, starting from the absolute reference
 63 time for the demand, which is defined by the user. In Step 3, the exchanges in the timeline
 64 are re-linked to time-specific background databases that reflect the technology landscape at
 65 specific points in time. Based on the temporally re-linked product system, a time-explicit LCI
 66 is calculated, preserving the timing of processes and emissions. The inventory is calculated
 67 following the conventional matrix-based LCA formulation (Heijungs & Suh, 2002), with the
 68 time dimension embedded in the matrices through additional row/column pairs. In Step 4,
 69 these emissions are characterized, either using standard characterization factors or by applying
 70 dynamic characterization functions that take the emissions' timing into account.




Figure 1: Workflow for a time-explicit LCA with bw_timex.

71 Further reading

72 The documentation of the `bw_timex` package, including installation instructions, extensive
73 example notebooks and detailed API reference, can be found at <https://docs.brightway.dev/>
74 `projects/bw-timex`. For a detailed explanation of the methodological basis of time-explicit
75 LCA, please refer to our accompanying publication (Müller et al., 2025).

76 Acknowledgements

77 We thank Chris Mutel for his help in adapting the graph traversal algorithm. This work received
78 funding from ForestPaths, which is funded by European Union's Horizon Europe Research and
79 Innovation Programme (101036849) and United Kingdom Research and Innovation Council
80 (UKRI) (101056755). This work received funding from the ETH Board in the framework of
81 the Joint Initiative SCENE, Swiss Center of Excellence on Net Zero Emissions.

82 References

- 83 Arvidsson, R., Svanström, M., Sandén, B. A., Thonemann, N., Steubing, B., & Cucurachi, S.
84 (2024). Terminology for future-oriented life cycle assessment: Review and recommendations.
85 *The International Journal of Life Cycle Assessment*, 29(4), 607–613. <https://doi.org/10.1007/s11367-023-02265-8>
- 86 Beloin-Saint-Pierre, D., Albers, A., Hélias, A., Tiruta-Barna, L., Fantke, P., Levasseur, A.,
87 Benetto, E., Benoit, A., & Collet, P. (2020). Addressing temporal considerations in life
88 cycle assessment. *Science of The Total Environment*, 743, 140700. <https://doi.org/10.1016/j.scitotenv.2020.140700>
- 89 Cardellini, G., Mutel, C. L., Vial, E., & Muys, B. (2018). Temporalis, a generic method and
90 tool for dynamic Life Cycle Assessment. *Science of The Total Environment*, 645, 585–595.
<https://doi.org/10.1016/j.scitotenv.2018.07.044>
- 91 Heijungs, R., & Suh, S. (2002). *The Computational Structure of Life Cycle Assessment* (A.
92 Tukker, Ed.; Vol. 11). Springer Netherlands. <https://doi.org/10.1007/978-94-015-9900-9>
- 93 Joyce, P. J., & Björklund, A. (2022). Futura: A new tool for transparent and shareable scenario
94 analysis in prospective life cycle assessment. *Journal of Industrial Ecology*, 26(1), 134–144.
<https://doi.org/10.1111/jiec.13115>
- 95 Lang-Quantzendorff, L., & Beermann, M. (2025). Prosperdyn—a tool to describe dynamic
96 transitions in prospective life cycle assessment. *The International Journal of Life Cycle
97 Assessment*. <https://doi.org/10.1007/s11367-025-02515-x>
- 98 Müller, A., Diepers, T., Jakobs, A., Cardellini, G., von der Assen, N., Guinée, J., & Steubing, B.
99 (2025). Time-explicit life cycle assessment: A flexible framework for coherent consideration
100 of temporal dynamics. *The International Journal of Life Cycle Assessment*. <https://doi.org/10.1007/s11367-025-02539-3>
- 101 Mutel, C. (2017). Brightway: An open source framework for Life Cycle Assessment. *Journal
102 of Open Source Software*, 2(12), 236. <https://doi.org/10.21105/joss.00236>
- 103 Sacchi, R. (2026). *TRAILS: Temporal routing and aggregation of impacts across life-cycle
104 systems* (Version v1.0.0). <https://trails.readthedocs.io/en/latest/>
- 105 Sacchi, R., & Hahn-Menacho, A. J. (2024). Pathways: Life cycle assessment of energy
106 transition scenarios. *Journal of Open Source Software*, 9(103), 7309. <https://doi.org/10.21105/joss.07309>
- 107 Sacchi, R., Terlouw, T., Siala, K., Dirnachner, A., Bauer, C., Cox, B., Mutel, C., Daioglou,

- ¹¹⁴ V., & Luderer, G. (2022). PRospective EnvironMental Impact asSEment (*premise*):
¹¹⁵ A streamlined approach to producing databases for prospective life cycle assessment
¹¹⁶ using integrated assessment models. *Renewable and Sustainable Energy Reviews*, 160.
¹¹⁷ <https://doi.org/10.1016/j.rser.2022.112311>

DRAFT