

Braids group

Escuela simetrías
FECHA

Briseida Trejo
Ernesto Vázquez

Instituto de Matemáticas, UNAM



Instituto de
Matemáticas



Título

puras mamadas



Puras mamadas

The **n-strand braid groups** B_n has the presentation:

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_j = \sigma_j \sigma_i \text{ for } |i-j| \geq 2, \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \rangle.$$



Pure braids

Two words β_1 and β_2 represent the same braid $\Leftrightarrow \beta_1\beta_2^{-1} = 1$.



Pure braids

Two words β_1 and β_2 represent the same braid $\Leftrightarrow \beta_1\beta_2^{-1} = 1$.

Definition

Dibujo de pure braid



Pure braids


Two words β_1 and β_2 represent the same braid $\Leftrightarrow \beta_1\beta_2^{-1} = 1$.


Definition

Dibujo de pure braid

Theorem


The pure braid group P_n is biorderable for all $n \geq 1$.





strand 3

strand 3 (from bottom)



strand 3

strand 3 (from bottom)



Mapping class groups

Definition

Let \mathcal{S} an oriented compact surface, possibly with boundary, and \mathcal{P} be a finite set of distinguished interior points of \mathcal{S} .



Mapping class groups

Definition

Let \mathcal{S} an oriented compact surface, possibly with boundary, and \mathcal{P} be a finite set of distinguished interior points of \mathcal{S} .

The **mapping clas group** $\mathcal{MCG}(\mathcal{S}, \mathcal{P})$ of the surface \mathcal{S} relative to \mathcal{P} is the group of all isotopy classes of orientation-preserving homeomorphisms $\psi : \mathcal{S} \rightarrow \mathcal{S}$ satisfying $\psi|_{\partial\mathcal{S}} = id$ and $\psi(\mathcal{P}) = \mathcal{P}$.



Mapping class groups

Definition

Let \mathcal{S} an oriented compact surface, possibly with boundary, and \mathcal{P} be a finite set of distinguished interior points of \mathcal{S} .

The **mapping class group** $\mathcal{MCG}(\mathcal{S}, \mathcal{P})$ of the surface \mathcal{S} relative to \mathcal{P} is the group of all isotopy classes of orientation-preserving homeomorphisms $\psi : \mathcal{S} \rightarrow \mathcal{S}$ satisfying $\psi|_{\partial\mathcal{S}} = id$ and $\psi(\mathcal{P}) = \mathcal{P}$.

Proposition

There is an isomorphism $B_n \cong \mathcal{MCG}(D_n)$, where D_n is the disk D^2 with n regularly spaced points.



Order in B_n

σ -positive
subword property

Proposition

Thanks.