

# Braid groups

Briseida Trejo  
Ernesto Vázquez

*Instituto de Matemáticas, UNAM*

Summer Graduate School  
MSRI-CMO: Geometric Group  
Theory  
FECHA





# Braids

## Definition

An  $n$ -braid is a collection of  $n$  disjoint strings.

# Braids

## Definition

An  $n$ -braid is a collection of  $n$  disjoint strings.

## Examples



3-braid





## Braid groups

The **n-strand braid group**  $B_n$  has the presentation:

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_j = \sigma_j \sigma_i \text{ for } |i-j| \geq 2, \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \rangle.$$

## Braid groups

The **n-strand braid group**  $B_n$  has the presentation:

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_j = \sigma_j \sigma_i \text{ for } |i-j| \geq 2, \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \rangle.$$

### Generators of $B_4$



$\sigma_1$



$\sigma_2$



$\sigma_3$



$\sigma_1^{-1}$




$\sigma_2^{-1}$



$\sigma_3^{-1}$


## Relations



$$\sigma_1\sigma_3 = \sigma_3\sigma_1$$

$$\sigma_i\sigma_j = \sigma_j\sigma_i$$


if  $|i - j| \geq 2$




$$\sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2$$

$$\sigma_i\sigma_{i+1}\sigma_i = \sigma_{i+1}\sigma_i\sigma_{i+1}$$

## Multiplication (Concatenation)



## Multiplication (Concatenation)






## The word problem

Two words  $\beta_1, \beta_2 \in B_n$  represent the same braid  $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$ .

## The word problem


Two words  $\beta_1, \beta_2 \in B_n$  represent the same braid  $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$ .




$\sigma_1 \sigma_3 \quad \quad \quad \sigma_3 \sigma_1$

## The word problem

Two words  $\beta_1, \beta_2 \in B_n$  represent the same braid  $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$ .




$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$




$$\sigma_1 \sigma_3$$

## The word problem

Two words  $\beta_1, \beta_2 \in B_n$  represent the same braid  $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$ .




$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$



$$\sigma_1 \sigma_3 * \sigma_3^{-1} \sigma_1^{-1}$$


## The word problem

Two words  $\beta_1, \beta_2 \in B_n$  represent the same braid  $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$ .



$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$


$\Leftrightarrow$



$$\sigma_1 \sigma_3 * \sigma_3^{-1} \sigma_1^{-1} = \sigma_1 \sigma_3 \sigma_3^{-1} \sigma_1^{-1}$$


## The word problem

Two words  $\beta_1, \beta_2 \in B_n$  represent the same braid  $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$ .




$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$

$\Leftrightarrow$




$$\sigma_1 \sigma_3 * \sigma_3^{-1} \sigma_1^{-1} = \sigma_1 \sigma_3 \sigma_3^{-1} \sigma_1^{-1} = id$$

## Pure braids




Pure braid

## Pure braids




Pure braid




Not pure



## Pure braids



Pure braid



Not pure

### Theorem

The pure braid group  $P_n$  is biorderable for all  $n \geq 1$ .



## Mapping class groups

### Definition

Let  $\mathcal{S}$  an oriented compact surface, possibly with boundary, and  $\mathcal{P}$  be a finite set of distinguished interior points of  $\mathcal{S}$ .

## Mapping class groups

### Definition

Let  $\mathcal{S}$  an oriented compact surface, possibly with boundary, and  $\mathcal{P}$  be a finite set of distinguished interior points of  $\mathcal{S}$ .

The **mapping clas group**  $\mathcal{MCG}(\mathcal{S}, \mathcal{P})$  of the surface  $\mathcal{S}$  relative to  $\mathcal{P}$  is the group of all isotopy classes of orientation-preserving homeomorphisms  $\psi : \mathcal{S} \rightarrow \mathcal{S}$  satisfying  $\psi|_{\partial\mathcal{S}} = id$  and  $\psi(\mathcal{P}) = \mathcal{P}$ .

## Mapping class groups

### Definition

Let  $\mathcal{S}$  an oriented compact surface, possibly with boundary, and  $\mathcal{P}$  be a finite set of distinguished interior points of  $\mathcal{S}$ .

The **mapping class group**  $\mathcal{MCG}(\mathcal{S}, \mathcal{P})$  of the surface  $\mathcal{S}$  relative to  $\mathcal{P}$  is the group of all isotopy classes of orientation-preserving homeomorphisms  $\psi : \mathcal{S} \rightarrow \mathcal{S}$  satisfying  $\psi|_{\partial\mathcal{S}} = id$  and  $\psi(\mathcal{P}) = \mathcal{P}$ .

### Proposition


There is an isomorphism  $B_n \cong \mathcal{MCG}(D_n)$ , where  $D_n$  is the disk  $D^2$  with  $n$  regularly spaced points.




## Order in $B_n$

$\sigma$ -positive  
subword property


Proposition





strand 3

strand 3 (from bottom)



strand 3

strand 3 (from bottom)



Thanks.