

Braid groups

Briseida Trejo
Ernesto Vázquez

Instituto de Matemáticas, UNAM

Summer Graduate School
MSRI-CMO: Geometric Group
Theory
FECHA





Braids

Definition

An *n -braid* is a collection of n disjoint strings.

Braids


Definition

An **n -braid** is a collection of n disjoint strings.

Examples



3-braid



id



Braid groups

The **n-strand braid group** B_n has the presentation:

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_j = \sigma_j \sigma_i \text{ for } |i-j| \geq 2, \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \rangle.$$

Braid groups

The n -strand braid group B_n has the presentation:

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_j = \sigma_j \sigma_i \text{ for } |i-j| \geq 2, \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \rangle.$$

Generators of B_4



σ_1



σ_2



σ_3



σ_1^{-1}




σ_2^{-1}



σ_3^{-1}


Relations



$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i$$


if $|i - j| \geq 2$




$$\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2$$

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$$

Multiplication (Concatenation)



Multiplication (Concatenation)






The word problem

Two words $\beta_1, \beta_2 \in B_n$ represent the same braid $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$.

The word problem


Two words $\beta_1, \beta_2 \in B_n$ represent the same braid $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$.



$\sigma_1 \sigma_3$ $=$ $\sigma_3 \sigma_1$


The word problem

Two words $\beta_1, \beta_2 \in B_n$ represent the same braid $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$.



$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$


$$\Leftrightarrow$$



$$\sigma_1 \sigma_3$$


The word problem

Two words $\beta_1, \beta_2 \in B_n$ represent the same braid $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$.



$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$


$$\Leftrightarrow$$



$$\sigma_1 \sigma_3 \quad * \quad \sigma_3^{-1} \sigma_1^{-1}$$


The word problem

Two words $\beta_1, \beta_2 \in B_n$ represent the same braid $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$.



$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$


\Leftrightarrow



$$\sigma_1 \sigma_3 * \sigma_3^{-1} \sigma_1^{-1} = \sigma_1 \sigma_3 \sigma_3^{-1} \sigma_1^{-1}$$


The word problem

Two words $\beta_1, \beta_2 \in B_n$ represent the same braid $\Leftrightarrow \beta_1 \beta_2^{-1} = 1$.




$$\sigma_1 \sigma_3 = \sigma_3 \sigma_1$$

\Leftrightarrow




$$\sigma_1 \sigma_3 * \sigma_3^{-1} \sigma_1^{-1} = \sigma_1 \sigma_3 \sigma_3^{-1} \sigma_1^{-1} = id$$

Pure braids




Pure braid

Pure braids




Pure braid




Not pure

Pure braids



Pure braid



Not pure

Theorem

The pure braid group P_n is biorderable for all $n \geq 1$.



Mapping class groups

Definition

Let \mathcal{S} an oriented compact surface, possibly with boundary, and \mathcal{P} be a finite set of distinguished interior points of \mathcal{S} .

Mapping class groups

Definition

Let \mathcal{S} an oriented compact surface, possibly with boundary, and \mathcal{P} be a finite set of distinguished interior points of \mathcal{S} .

The **mapping class group** $\mathcal{MCG}(\mathcal{S}, \mathcal{P})$ of the surface \mathcal{S} relative to \mathcal{P} is the group of all isotopy classes of orientation-preserving homeomorphisms $\psi : \mathcal{S} \rightarrow \mathcal{S}$ satisfying $\psi|_{\partial\mathcal{S}} = id$ and $\psi(\mathcal{P}) = \mathcal{P}$.

Mapping class groups

Definition

Let \mathcal{S} an oriented compact surface, possibly with boundary, and \mathcal{P} be a finite set of distinguished interior points of \mathcal{S} .

The **mapping class group** $\mathcal{MCG}(\mathcal{S}, \mathcal{P})$ of the surface \mathcal{S} relative to \mathcal{P} is the group of all isotopy classes of orientation-preserving homeomorphisms $\psi : \mathcal{S} \rightarrow \mathcal{S}$ satisfying $\psi|_{\partial\mathcal{S}} = id$ and $\psi(\mathcal{P}) = \mathcal{P}$.


Proposition


There is an isomorphism $B_n \cong \mathcal{MCG}(D_n)$, where D_n is the disk D^2 with n regularly spaced points.

Order in B_n

σ -positive
subword property


Proposition





strand 3

strand 3 (from bottom)



strand 3

strand 3 (from bottom)

Thanks.