

# Complex Dynamics

Lexi Reed and Brian Zhang

May 2021

## 1 Julia and Mandelbrot Set Definitions

Let  $P_c(z) = z^2 + c$

Julia set for  $P_c$ , fixes  $c$

$$J_c = \{z \mid |P_c^n(z)|_n \text{ bounded}\}$$

Mandelbrot set instead fixes  $z$

$$M = \{c \mid |P_c^n(0)|_n \text{ bounded}\}$$

To be more precise

- Def above is the "filled-in" Julia set
- "True" Julia set is the boundary
- Above is a Quadratic Julia set, can use other rational functions (ratio of complex polynomials)

## 2 Julia Set Fractal Nature

Sequence  $z_n = P_c^n(z_0)$ , iterating  $P_c$  for a given  $z_0$  in the Julia set.

Easier to visualize the pre-image (single iteration) of a point  $z_n = P_c^{-1}(z_{n+1})$

$$z_{n+1} = \left(|z_n| e^{i\text{Arg}z}\right)^2 + c$$
$$z_{n+1} - c = |z_n|^2 e^{i2\text{Arg}z}$$

Solving for  $z_n$  in terms of  $z_{n+1} - c$

$$2\text{Arg}z_n = \text{Arg}(z_{n+1} - c) + 2\pi k$$
$$\text{Arg}z_n = .5\text{Arg}(z_{n+1} - c) + \pi k, \quad k \in \{0, 1\}$$
$$|z_n| = \sqrt{|z_{n+1} - c|}$$

We can iterate taking the pre-image of a set containing  $J_c$  to approach its shape,

- 
- Take nbhd of infinity  $S$  s.t.  $J_c \subset \mathbb{C} - S$
  - $z_0 \notin J_c \iff \exists N \ni \forall n > N, z_n \in S$
  - $\mathbb{C} - J_c = \bigcup_n P_c^{-n}(S)$
  - DeMorgan's, pre-image of complement:  $J_c = \bigcap_n P_c^{-n}(\mathbb{C} - S)$




Figure 1: Julia set, bands around set indicate "escape" iterations [3]

Karl Sims gives a visualization for a disk in Fig 2. For one iteration,

- Translate the disk by  $-c$ , halve angle for each point, shift radius towards 1 for each point
- Mirror across real axis (pre-image maps one point to two with  $\pi$  arg offset)




Figure 2: Approaching Julia set by pre-images of disk [9]

### 3 Julia Set Escape Radius

How big does the disk need to be such that  $J_c \subset D[0, R]$ ?

- Called "Escape Radius"
- Find  $R$  where  $z_0 \notin D[0, R] \implies |z_{n+1}| > |z_n|$

For  $P_c(z) = z^2 + 1$ , take  $|z_n| > |R| > 1$ ,

$$\begin{aligned} |z_{n+1}| &\geq \left| z_n^2 \right| - |c| && \text{Reverse Triangle Inequality} \\ &\geq \left| z_n^2 \right| - |z_n| \frac{|z_n|}{R} |c| \\ &= |z_n| \frac{|z_n|}{R} (R - |c|) \end{aligned}$$

Thus  $R > 1 + |c| \implies |z_{n+1}| > |z_n|$  [7].

Plotting algorithm: take points in disk, iterate  $P_c$ , remove points that "escape" the disk.

### 4 Julia Set Connectedness

- Julia Set is either connected (Fatou Set)...
  - Start from  $D[0, R]$ , each pre-image iteration preserves connectedness
- Or something that's like the Cantor set (Fatou Dust)
  - Start from  $D[0, R]$ , each pre-image iteration cuts the previous pieces in two
- If  $0 \in J_c$ , then it's connected, otherwise Fatou Dust
- $J_c$  connected  $\iff 0 \in J_c \iff c \in M$  (M is Mandelbrot set)



Figure 3: Cantor set interval visualization [1]

TODO something about Julia set fixed points, 0 is circle, -2 is chaotic, Mandelbrot explore the transition




Figure 4: Fatou dust,  $c = .45 + .1428i$

## 5 Mandelbrot Set Properties

- Mandelbrot set is quasi-fractal, whereas Julia is fractal
  - "Copies" are deformed (see Figure 5)
  - "Index" of (connected) Julia sets ( $c \notin M$  for disconnected  $J_c$ )
  - Mandelbrot at  $c$  resembles  $J_c$  near 0
- Mandelbrot set is connected
  - Mandelbrot conjectured it was not from initial images!
  - Conformal (angle-preserving) isomorphism to complement of a closed disk
- Open question: Is Mandelbrot locally connected (MLC)?
  - $\forall x \in M, x \in \text{open } U \implies \exists \text{open, connected } W, x \in W \subset U$
  - Example: Topologist's sine curve connected, not locally connected

## 6 Mandelbrot Bound / Escape Radius

Let  $z_n = P_c^n(0)$ ,

$$c \in M \iff \exists n \ni |z_n| > 2 \text{ [10]}$$

Given  $|z_n| > 2$ , for  $|c| \leq 2$

$$\begin{aligned} |z_{n+1}| &\geq |z_n|^2 - |c| && \text{Reverse Triangel Inequality} \\ &\geq 2|z_n| - 2 = |z_n| + (|z_n| - 2) \\ &> |z_n| \end{aligned}$$

For  $|c| > 2$ , inductively  $|z_n| > |c|$ , so

$$|z_{n+1}| \geq |z_n|^2 - |c| > |c||z_n| - |c| > 2|z_n| - |c| > |z_n|$$




Figure 5: Typical Mandelbrot [4] and a deformed "Mini-Mandelbrot" [8]




Figure 6: Topologist's Sine Curve [5]

---

## 7 Mandelbrot Fixed Points and Cycles

- "Main" Cardioid contains points whose sequence (orbit) approach attractive fixed point
  - For fixed point  $x^*$ ,  $x_{n+1} = f(x_n) \approx f(x^*) + f'(x^*)(x_n - x^*)$
  - $|x_{n+1} - x^*| = |x_{n+1} - f(x^*)| \approx |f'(x^*)| |x_n - x^*|$
- Other cardioids have cycles, periods labeled in Fig 7

For  $c$  with an attractive fixed point  $z^*$  of  $M$ , we must have

$$z^* = (z^*)^2 + c$$

and

$$\left| \frac{d}{dz} \Big|_{z^*} [z^2 + c] \right| < 1, |z^*| < \frac{1}{2}$$

For  $c$  on the boundary of those with fixed points, we have  $|z^*| = \frac{1}{2}$ . Combining with the first condition, the solutions are the main cardioid curve.




Figure 7: Cardioids labeled with period [6]

## 8 Visualizations

## 9 General Julia Sets

Definitions

- In general, iterating function  $f(x) = p(x)/q(x)$  for complex polynomials  $p, q$ , no shared roots
  - $f$  maps Riemann sphere onto itself
  - $f$  is holomorphic
- $f$  is invariant on open sets, Fatou Domains

- 
- Fatou Set is union of Fatou Domains, it's dense!
  - "True" Julia Set is the complement of the Fatou Set (boundary), also invariant

In the context of the Quadratic Julia Set described earlier...

- "True" Julia Set is boundary of filled-in Julia Set
- Fatou Domains are...
  - interior of filled-in Julia Set
  - other side of Julia Set boundary, with unbounded orbit

## 10 Newton Fractals

- Newton method
  - Iteratively approximate zeroes using tangent lines
  - $x_{n+1} = x_n - p(x_n)/p'(x_n)$
  - Works with complex functions
- $f(x) = x - p(x)/p'(x)$  can be used as the iteration function for a Julia Set
- One Fatou Domain for each zero of  $f$
- True Julia Set is the boundary where the method fails!




Figure 8: Newton's Method

---

## References

- [1] Wikimedia Commons. File:cantor set in seven iterations.svg — wikimedia commons, the free media repository, 2020. [Online; accessed 30-May-2021].
- [2] Wikimedia Commons. File:julia set for the rational function.png — wiki-media commons, the free media repository, 2020. [Online; accessed 30-May-2021].
- [3] Wikimedia Commons. Time escape Julia set from coordinate ( $\phi-2$ , 0).jpg — wikimedia commons, the free media repository, 2020. [Online; accessed 30-May-2021].
- [4] Wikimedia Commons. File:mandel zoom 00 mandelbrot set.jpg — wikimedia commons, the free media repository, 2021. [Online; accessed 30-May-2021].
- [5] Wikimedia Commons. File:topologist's sine curve.svg — wikimedia commons, the free media repository, 2021. [Online; accessed 31-May-2021].
- [6] Michael Frame. The Mandelbrot set and Julia sets. [Online; accessed 30-May-2021].
- [7] Lutz Lehmann (<https://math.stackexchange.com/users/115115/lutz-lehmann>). The escape radius of a polynomial and its filled julia set. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/3839776> (version: 2020-09-25).
- [8] mateuszica (<https://math.stackexchange.com/users/86329/mateuszica>). Mini mandelbrots, are they exact copies? Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/595839> (version: 2017-03-05).
- [9] Karl Sims. Understanding Julia and Mandelbrot sets. [Online; accessed 30-May-2021].
- [10] user147263. Mandelbrot sets and radius of convergence. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/890448> (version: 2014-12-19).