

# SARS-CoV-2 sequencing report

The Sequencing Lab is sequencing SARS-CoV-2 from patients.


This report, generated on 2021-03-18, summarizes our sequencing activity for patient samples provided by **Broad Institute Clinical Research Sequencing Platform**. The samples included in this report were sequenced between 2021-02-06 and 2021-03-12. This report is current as of the epiweek ending **2021-03-13**.


## Weekly summary

	week ending 2021-03-13	cumulative total
samples sequenced	184	486
genomes assembled	179	480
genomes submittable	173	454
Variants of Interest (Vols)	5	6
Variants of Concern (VoCs)	11	21

CDC definitions of Variants of Concern (VoCs) and Variants of Interest (Vols) are available at: <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/variant-surveillance/variant-info.html>

## Sequencing activity over time and space






This describes sequencing attempts over time, broken down by geography.

## Sequencing performance over time

### Samples sequenced weekly



This describes the total number of patient samples sequenced in this data set, plotted by the date of the sequencing run in our lab.


**Submittable** genomes pass all QC checks and are quickly released to public genome repositories. **Failed sequencing** are samples that failed to produce at least 24000 unambiguous base pairs of viral genome. Raw data from these samples are submitted to NCBI's SRA database, but the genomes are not used for any analyses. **Failed annotation** are samples that produced a sufficiently complete genome, but did not pass NCBI's VADR quality checks.

### Tabular view by CDC epiweek

epiweek ending	samples sequenced	genomes assembled	genomes submittable
2021-02-06	91	91	87
2021-02-13	31	31	31
2021-02-27	180	179	163
2021-03-13	184	179	173

## Timeliness of surveillance


### Sequencing date vs collection date



This plot describes the “timeliness” of the sequencing run for the purpose of real-time surveillance of circulating lineages and variants of interest. Note that this plot likely includes many samples that were sequenced for non-surveillance purposes.

## Evolutionary Clock

### Genetic distance root-to-tip vs sample collection date



A "root-to-tip plot" plots the genetic distance of each sample from Wuhan Hu-1 against the date it was collected. It is generally somewhat linear. Outliers on this plot may be indicative of laboratory or metadata errors, or of evolutionarily unusual lineages (such as B.1.1.7).

## Phylogenetic Clades and Variants

### Reportable Variants of Concern (VoCs) by CDC epiweek of sample collection


Collection epiweek ending	B.1.1.7	B.1.351	B.1.427	B.1.429
2021-01-09	0	0	1	0
2021-01-16	1	0	0	0
2021-01-30	4	3	0	1
2021-02-06	3	0	0	0
2021-02-13	3	0	0	2
2021-02-20	1	0	0	1
2021-02-27	1	0	0	0

### Variants of Interest (VoIs) by CDC epiweek of sample collection

Collection epiweek ending	B.1.526	P.2
2021-01-09	0	1
2021-02-13	2	0
2021-02-20	1	0
2021-02-27	2	0

CDC definitions of Variants of Concern (VoCs) and Variants of Interest (VoIs) are available at:  
<https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/variant-surveillance/variant-info.html>

### Nextclade classifications vs sample collection date



This shows the breakdown of major phylogenetic clades over time, using the Nextclade naming system. Variants of Concern (VoCs) are highlighted as specially named Nextclade clades. Nextclade clade 20I/501Y.V1 corresponds to PANGO lineage B.1.1.7, 20H/501Y.V2 corresponds to B.1.351, and 20J/501Y.V3 corresponds to P.1. At this time, no other VoC or VoI lineages have dedicated Nextclade definitions.