

## ***0.a. Goal***

3 GUARANTEE ACCESS TO QUALITY HEALTH AND PROMOTE WELFARE FOR ALL

## ***0.b. Target***

3.2 By 2030, end preventable deaths of newborns and children under 5 in all countries with the aim of reducing neonatal mortality to at least 12 per 1,000 live births and the mortality of children under 5 to at least 25 per 1,000 live births

## ***0.c. Indicator***

### **3.2.1 Mortality rate of children under 5 years old (Deaths per 1000 live births)**

---

## ***0.d. Series***

Mortality of children under 5 years old (Deaths per 1000 live births)

## ***0.e. Metadata update***

10/9/2020

## ***1.a. Organisation***

NATIONAL INSTITUTE OF STATISTICS

## ***1.b. Contact person(s)***

Teixeira Mandlate, Maria Alfeu and João Manguê

## ***1.c. Contact organisation unit***

Directorate of Demographic, Vital and Social Statistics

## ***1.d. Contact person function***

Technicians

## ***1.e. Contact phone***

+258 844386629; +258 823032619; +258 827679480

## ***1.f. Contact mail***

www.ine.gov.mz

## ***1.g. Contact email***

[Teixeira.mandlate@ine.gov.mz](mailto:Teixeira.mandlate@ine.gov.mz); [maria.alfeu@ine.gov.mz](mailto:maria.alfeu@ine.gov.mz); [joao.mangue@ine.gov.mz](mailto:joao.mangue@ine.gov.mz)

## ***2.a. Definition and concepts***

Infant mortality (below five years) is the probability that a child born in a specific year or period will die before reaching 5 years of age, subject to age-specific mortality rates for that period, expressed per 1,000 births alive.

## ***2.b. Unit of measure***

Percentage

## ***3.a. Data sources***

Statistics Portugal, Demographic and Health Survey (IDS); Statistics Portugal, General Population and Housing Census

## ***3.b. Data collection method***

The recommended data sources for calculating this indicator are Household Surveys. The Demographic and Health Survey data collection questionnaire was designed and subsequently tested in the field during the training of interviewers.

For data collection, the methodology of interviews was applied face to face to the households, applying three types of questionnaires:

- Household Questionnaire
- Women's Questionnaire
- Men's Questionnaire.

The Sample Design

The Demographic and Health Survey comprises a probabilistic, stratified and multi-stage sample, selected from the Data and Cartography of the III General Census of Population and Housing, carried out by INE in 2007.

The data collection lasted for five months starting in June 2011, ending in November 2011.

Response rate

Of the 13,964 households interviewed in the survey, a total of 13,871 eligible women were identified. Interviews were conducted with 13,718 of these women, which resulted in a response rate of 99%.

Introduction • 13 In one third of the IDS household sample, interviews were also conducted with all eligible men found. Thus, of the 4,130 eligible men identified in the subsample of households selected for the men survey, 4,027 were successfully interviewed, giving a response rate of 98%

### ***3.c. Data collection calendar***

2021

### ***3.d. Data release calendar***

2022

### ***3.e. Data providers***

National Institute of Statistics

### ***3.f. Data compilers***

National Institute of Statistics

### ***3.g. Institutional mandate***

Law 7/96 of 5 July

## ***4.a. Rationale***

Mortality rates among children under five are indicators of public health follow-up because it reflects the access of children and communities to basic health interventions, such as vaccination, medical treatment of infectious diseases and adequate nutrition.

The indicator shows the number of deaths under the age of five that occur in every 1000 children under the age of five

## ***4.b. Comment and limitations***

Mortality rates among young children are a key indicator of measuring child health and well-being and, more broadly, for social and economic development. It is an indicator of public health monitoring because it reflects the access of children and communities to basic health interventions, such as vaccination, medical treatment of infectious diseases and adequate nutrition.

## ***4.c. Method of computation***

The indicator is calculated by dividing the number of deaths of children aged 0 to 59 months (< de 5 anos) num determinado período e o número médio de crianças com idade entre os 0 a 59 meses (< 5 anos) durante o mesmo período de tempo

#### ***4.d. Validation***

The field work had close supervision and quality control by the central and provincial technicians, both from INE, MISAU and ICF International staff. In addition, during the data collection, a strict control was established at the level of each team over the collection process, by detecting errors by the field critics, which allowed for immediate correction still on the ground. At the level of central coordination, the data critics carried out a further review of the base data and the problems encountered were communicated to the respective teams.

The interactive and batch processing of information through the CSPro program also allowed, at central level, the periodic obtaining of partial results, for analysis of the data collected until a given moment, through the production of tables for monitoring and quality control. The results of these tabulations were reported in feedback to the interviewers, ensuring data quality.

#### ***4.h. Methods and guidance available to countries for the compilation of the data at the national level***

The Demographic and Health Survey (IDS) in Mozambique is part of an international survey program (MEASURE DHS) developed by ICF International through a contract with USAID, with the purpose of supporting governments and private institutions in developing countries in conducting national surveys by sampling, in the areas of population and health. The MEASURE DHS Program aims to:

- Support the formulation of policies and implementation of programs in the areas of population and health;
- Increase the international population and health data base for monitoring and evaluation;
- Improve the survey methodology by sampling, and
- Consolidate, in the survey area, the technical capacity of the executing institution in the country participating in the Program.

#### ***4.i. Quality management***

Quality Management Instrument still to be approved

#### ***4.j. Quality assurance***

The interactive and batch processing of information through the CSPro program also allowed, at central level, the periodic obtaining of partial results, for analysis of the data collected until a given moment, through the production of tables for monitoring and quality control. The results of these tabulations were reported in feedback to the interviewers, ensuring data quality.

#### ***4.k. Quality assessment***

Quality Assessment Instrument still to be approved

### ***5. Data availability and disaggregation***

Data are available every 5 years and can be disaggregated by sex, rural and urban area of residence, province and country

## ***6. Comparability/deviation from international standards***

Survey estimates are based on standardized methodology, using the WHO Child Growth Standards, as described in (Ref: Anthro software manual). Global and regional estimates are based on the methodology described at UNICEF-WHO, World Bank: Joint estimates of child malnutrition - Levels and trends (UNICEF / WHO / BM)

## ***7. References and Documentation***

Ministry of Health (MISAU), [www.misau.gov.mz](http://www.misau.gov.mz); National Statistics Institute (INE), [www.misau.gov.mz](http://www.misau.gov.mz); ICF International (ICFI), [www.measuredhs.com](http://www.measuredhs.com)