

T-Mobile Park's Effect on Major League Baseball in Seattle

Brodie Monohan
University of Washington
monohanb@uw.edu

August 7, 2026

Abstract

The Seattle Mariners are the only Major League Baseball team to never appear in a World Series. Despite being near the middle of the pack in terms of payroll, Mariner teams since the early 2000s have been disappointing to say the least, with historically bad offenses in the 2010s. Since the team's move from their old stadium into what would become T-Mobile Park (formerly Safeco Field), some analysts have noticed that batters seem to perform poorly in the stadium. Former sluggers who were traded to the Mariners would constantly under-perform with their new team, or so it seemed. Some analysts began to theorize there was some external factor causing batters to lose production when being traded to the Mariners. The most prominent theory is that the Seattle climate and the stadium itself that was causing hitters to under-perform. This theory is reinforced by the pitching statistics for Seattle which, since 2000, have been historically good. While a "cursed" stadium is a romantic way to explain the Mariner's historic woes, there is also a much more ordinary explanation: that Mariners have just been bad hitters. This project examines if hitting in Seattle is significantly "harder" or if players are worse when playing in Seattle by looking at historical batting numbers and investigating the physical conditions of T-Mobile Park.

GitHub repo: <https://github.com/brodiemonohan/MLB-in-Seattle>

1 Research Questions

I set out to answer three questions about the supposed disadvantages for hitters in T-Mobile Park. The first question was if hitter performance demonstrated a systematic dip as a Seattle Mariner or at T-Mobile Park by looking at traded player performance and historic home vs away splits respectively. The results indicate no systematic trends in loss of production as a Seattle Mariner or in T-Mobile Park. The average

change in performance remained within 3% of zero for both home/away splits and season performance on either side of a trade with Seattle for a range of stats including On-Base Percentage (OBP), Slugging (SLG), Home Run rate (HR rate), and singles (1B). The second question was if the environment of T-Mobile Park (and Seattle in general) was a significant factor in the flight of a baseball such that playing at T-Mobile Park is disadvantageous in terms of how far a ball travels

for given hit parameters. The results did show a significant difference in trajectories between stadiums (as was expected). Based on temperature, elevation, and coordinate data, a ball hit in the range of typical home-run parameters can travel up to 10 meters farther in Colorado compared to Seattle. The third question was if these results indicated an overall competitive disadvantage for the Seattle Mariners. The results for this question were mixed. Historical team batting numbers suggest that some Mariner teams, especially in the past few years, have relied on hard-hit balls, slugging, and home-runs. The first question indicated that historically, SLG and HR rate were seemingly unaffected by T-Mobile Park or the Mariners organization, however, the second question indicated that there is a physical limitation in T-Mobile Park for hard-hit and high hang-time balls, like home-runs. Possible explanations are explored in the results section.

2 Motivation

My motivation for this research has stayed pretty much the same which is described in the abstract (I put it up there because I think it is better to read first, and it goes nicely into a README). I wanted to investigate the ad nauseum argument that hitting is harder in Seattle. I made no updates since the EDA assignment.

3 Dataset(s)

My main dataset comes from `pybaseball`. The MLB and third parties have tracked statistics since far before the 1999 season when the Mariners changed stadiums, however, the MLB started to take advanced statistics after the 2014 season with StatCast which includes things like swing speed, launch angle, launch velocity, and other advanced metrics which will be useful for my analysis. An open source Python package already exists for these analytics called `pybaseball` which scrapes Baseball Savant (built on StatCast), as well as Baseball Reference which are two more proprietary baseball statistic trackers. Documentation can be found at pypi.org/project/pybaseball/. This package works with `pandas` and can be indexed and manipulated as a `DataFrame`.

My second dataset comes from `openmeteo-requests` which collects hourly weather data from Open Mateo as a Python package. Documentation can be found at <https://pypi.org/project/openmeteo-requests/>. To access this data I had to use a new library (`requests`) which is my other challenge goal.

My third dataset is `MLB-StatsAPI`. Documentation can be found at <https://pypi.org/project/MLB-StatsAPI/>. `pybaseball` did not have direct access to MLB game-time data, so I

had to use this instead which had a very similar format to `openmateo-requests` and used the `requests` library.

My fourth dataset is a small `.csv` file I manually scraped from Google Maps and MLB.com which has stadium longitude and latitude data as well as venue IDs for MLB-StatsAPI.

4 Method

4.1 Historical Batting

I look at systematic drops in batter performance in T-Mobile Park by comparing home and away splits as well as season performance on either side of a trade with the Seattle Mariners. I pull batting averages (BA), on-base percentages (OBP), home-run rate, and slugging percentages (SLG) for players traded to or from Seattle since 2008 and plot them as a one-dimensional histogram.

4.2 Atmosphere Effects

For the second question I can find the average drag on the ball for each MLB stadium by calculating the air density based on `open-mateo` data and the ideal gas physics then numerically calculate a trajectory through x-z space using Euler's Method. There are three main forces that influence the trajectory of a ball: gravity, drag, and the Magnus force. Fig 1 shows an ideal example free-body diagram of these forces and the direc-




Figure 1: A free-body diagram of a baseball in-flight illustrating the three main forces influencing the ball's trajectory through x-z space. The magnitudes are not to scale. The ball's direction of motion is at an angle of 45° ($\pi/4$ rad) from parallel with the ground with pure backspin. The optimal angle is actually shallower, between 20 and 30 degrees. The spin axis is usually not purely in the y dimension (in or out of the screen), however, in the absence of real spin axis data, this is a decent assumption. "w" is standing for ω here because I haven't figured out how to type Greek letters in Photoshop yet.

tion of motion. These forces are described parametrically in Adair (2002), Sawicki et al. (2003), and Nathan (2008b) as

$$F_g = -mg \quad (1)$$

$$F_d = \frac{1}{2}C_d\rho Av^2 \quad (2)$$

$$F_m = \frac{1}{2}C_l\rho Av^2 \quad (3)$$

where m is the mass of the ball, g is the acceleration due to gravity, v is the velocity vector, ρ is the air density, A is the cross-sectional area of the ball, C_d is the drag coefficient, and C_l is the lift coefficient. The drag and lift coefficients for an MLB baseball were experimentally determined in Sawicki et al. (2003) and Nathan (2008b) as

$$C_l = \begin{cases} 0.09 + (0.6S) & \text{if } S > 0.1 \\ 1.5S & \text{if } S \leq 0.1 \end{cases} \quad (4)$$

$$C_d = 0.3 + 0.03S \quad (5)$$

where $S = R\omega/v$. The radius and mass of an MLB baseball are also well established; I will use $R = 0.0364\text{m}$ and $m = 0.145\text{kg}$. F_d and F_m depend on ω and v in addition to R which will be input parameters to the model (hit variables). This leave just the air density which I can esti-

mate using the ideal gas law

$$PV = nRT \rightarrow n/V = \rho = P/RT \quad (6)$$

where P is pressure (Pa), V is volume (m^3), n is the number of moles, R is the ideal gas constant ($\text{J} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$), and T is the temperature (K), and ρ is the air density ($\text{mol} \cdot \text{m}^{-3}$). This equation will yield the number density of moles of air particles per cubic meter which I can convert to a mass density (which the above models expect) by multiplying the number density by the mass of dry air $m_{\text{air}} = 0.0289647 \text{ kg} \cdot \text{mol}^{-1}$. To get pressure I can pull elevation data from **open-mateo** using stadium coordinate data. The elevation can translate to pressure via barometric formula.

$$P = P_0(1 - 2.25577 \cdot 10^{-5} \cdot e)^{5.25588} \quad (7)$$

where $P_0 = 101325 \text{ Pa}$ is the pressure at sea level, and e is the elevation (m). This leaves just the temperature unresolved. For this I can pull hourly temperature data from **openmateo-requests** from the start time of each game during the 2025 MLB season for each team to find the average air density during the average game at each current MLB stadium. All together, I can combine equations 1-7 with Euler's Method to numerically calculate a trajectory for

a given set of hit parameters (v , ω , θ_{angle} (launch angle), and θ_{axis} (spin axis)) for the average game at each current MLB stadium.

4.3 Competitive Advantages

For the third question I look at if there has been any major competitive impact to Mariners players or seasons based on player archetype. This question is specifically motivate by this year's team which is built as a "slugging" team which some criticize for a place like Seattle for not playing into the strengths of the park. The New York Yankees, for example, tend to buy or trade for left-handed power hitters because right field in Yankee Stadium is significantly shorter than in other parks, turning pop-outs into home-runs for 81 games a year. I look at team batting stats like SLG, HR rate, and isolated power (ISO) and compare them to the league average.

5 EDA Results

For the EDA I built one of the functions for question 1: how does player performance change when traded to or from the Mariners? I also made significant progress on question 2: does the air density, wind, temperature, and/or humidity in Seattle cause baseballs to travel differently compared to other major league stadiums and is it disadvantageous to hitters?

5.1 Question 1

I attacked this one first because seemed the hardest to implement. To do this, I used the `batting_stats_bref` function from `pybaseball` which takes a year argument and returns a DataFrame of all the batting statistics for that season (seasons lineup with calendar years in MLB baseball). Each row is a unique player with the headers

```
'Name', 'Age', '#days', 'Lev', 'Tm',
'G', 'PA', 'AB', 'R', 'H', '2B', '3B',
'HR', 'RBI', 'BB', 'IBB', 'SO', 'HBP',
'SH', 'SF', 'GDP', 'SB', 'CS', 'BA',
'OBP', 'SLG', 'OPS', 'mlbID'
```

For my function, I wanted the inputs to be a team (actually a location of a team because the 'Tm' parameter returns 'Seattle' for Mariner players for example), as well as a range of seasons which will default to all the available data if not specified (2008 – 2026). I also wanted an optional minimum plate appearances or at-bats so that players who only played a small amount can be filtered out. The output from this file is shown in Fig 2.

5.2 Question 2

Using the equations described in Sec 4.2, I built the numerical calculator that traces a ball's trajectory in x-z space. This time-stepping scheme




Figure 2: The change in performance of all players traded to or from the Seattle Mariners between 2008 and 2026 in the seasons on either side of the trade. Negative values indicate worse performance as a Mariner and vice versa. There are no cuts for minimum play (AB or PA).

is a very classing structure to follow in Physics models. It is commonly called Euler’s Method. The basic idea is at each timestep, you take the conditions from the previous step (like position and velocity) and apply the forces to the object based on those conditions, updating them for the next time step. When the time steps are small enough, you can trace out smooth arcs like how a baseball travels through the air. Any undergraduate physicist will have done this at some point, the challenge here was just moving it to Python. The output plot from main is shown in Fig 3 with some fun environments that test the extremes.

during the EDA I was worried about the effect of spin-down on the ball as friction with the air causes the ball so slow its angular velocity, however, Nathan (2008a) suggests this effect will be trivial in all real-world conditions.

6 Results

6.1 Historical Batting

The stats I chose to look at for batting statistics were 1B rate (single rate), OBP, HR rate, and SLG. I had initially planned on using batting average as well but Slowinski and Weinberg (2010) suggests that this stat requires a very large sample size to converge to statistically significant (920 plate appearances (PA) to be exact which is




Figure 3: The trajectory of a baseball hit at 45m/s at an angle of $\pi/4$ rad (45°) with pure backspin and a spin rate of 150 rad/s ($\approx 1,500$ rpm) in different environments. The environmental data on air density each came from a quick google search but will eventually be replaced by a calculated air density based on the elevation and temperature data from my second dataset.

way too big to be useful for any of the analysis I'm doing here). Instead I focused on OBP (460 PA), 1B rate (290 PA), HR rate (170 PA), SLG (320 AB). For the traded player stats, single rate did not end up being easily calculable and was not worth re-adding considering the results.

There was little to no difference in overall hitting splits for players traded to or from the Mariners for OBP, HR rate, and SLG. Additionally, the sample sizes were too small to resemble a clean gaussian distribution, indicating that they are not statistically significant at this scale. This can be seen in Fig 4. HR rate, which had the lowest minimum PA threshold, does resem-

ple a gaussian distribution with 20 counts at the peak but deviates by less than 0.5% from 0. This suggests no overall effect on player's home-run rates when changing home fields to or from T-Mobile Park.

There was also little to no difference when comparing home or away splits for Mariner hitters. These sample sizes were much larger and produced much more gaussian distributions, however, while the splits for three of the four tracked stats were negative, they deviated from zero only $\sim 3\%$ at the most. HR rate did not deviate. This is shown in Fig 5

6.2 Atmosphere Effects

The analysis here did yield some significant results, however, this was not unexpected. The average temperature in Seattle was low compared to other low elevation areas like Houston or Miami which is inversely related to air density in the ideal gas law. This means the air density in Seattle during the average game was the second highest (behind San Francisco). This translated to a $\sim 6\%$ reduction in travel distance as compared to the lowest air density stadium, Colorado. This is shown in Fig 6.

This does not consider, however, ballpark dimensions which is an important factor for competitive advantages. Would a ball hit at T-Mobile Park have been a home-run in other




Figure 4: The statistical splits for OBP, HR rate, and SLG for traded players between 2008 and 2025. More negative values indicate a worse performance as a Mariner.

parks?

6.3 Competitive Disadvantages

To see how Mariner teams have historically been built I look at team batting in SLG, HR rate, and isolated power (ISO). This analysis showed that teams in the late 2000s and early 2010s were below league average in terms of power hitting compared to league average, leveled out in the mid to late 2010s, then became hard-hit focused in the 2020s. The full splits can be found in Table 1. If the first two questions had yielded significant results, then this could have had some implications. For example, if I found that hard-hit metrics dipped at T-Mobile then I could have argued that slugging teams would not be a good build for a team in Seattle. While the second question did show that balls will travel less-far

in Seattle, I could not account for stadium dimensions which is a significant factor in if a hit becomes a home run and could potentially explain the discrepancy between questions 1 and 2. This leaves question 3 with little insight into competitive advantages or disadvantages in T-Mobile Park.

7 Impact and Limitations

These results from question 1 show little systematic production changes due to players playing in Seattle or for the Mariners. This would suggest that the poor team performances are due to errors in recruiting and player evaluation by the organization. It also could suggest that perhaps Mariner teams intentionally use capital in areas besides hitting. The results from question

Home and Away Splits




Figure 5: The home and away splits for OBP, HR rate, single rate, and SLG for Mariner players between 2008 and 2025. More Negative indicates a worse performance at home.




Figure 6: The simulated trajectories for all 30 current MLB stadiums (except Tampa Bay because of stadium change between 2025 and 2026) for a ball with hit parameters $v = 45 \text{ m} \cdot \text{s}^{-1}$, $\omega = 150 \text{ rad} \cdot \text{s}^{-1}$, $\theta_{\text{axis}} = 0 \text{ rad}$ (pure backspin) and $\theta_{\text{angle}} = \pi/6 \text{ rad}$.

2 show that hard-hit, high-angle balls do not result in as far trajectories in Seattle compared to other stadiums. This would suggest that sluggers who rely on home-runs for run production likely aren't the best fit for a team playing in Seattle for half of their games. This is a particularly important implication given the current team's reliance on home-runs, however, as discussed in the results this may be misleading. There are clear limitations to this analysis. The most pressing are the field dimensions which weren't accounted for here and the assumptions about spin rate on batted balls which is not publicly tracked.

8 Challenge Goals

My challenge goals have been a bit of a mess because I am doing a lot of things that could be a challenge goal. In the end I used multiple datasets and used a second library which were my proposal goals. I used `openmateo-requests`, `MLBStats-API` and the small manually entered `stadium_data.csv` for the weather and elevation data in combination with the trajectory calculator function. I also used a second library `requests` which can be found at <https://pypi.org/project/requests/>.

Mariner Power Team Stats			
Season	ISO	SLG	HR_rate
2008	-0.030271	-0.026491	-0.006508
2009	-0.001676	0.004956	0.001869
2010	-0.051938	-0.073512	-0.011271
2011	-0.028755	-0.050709	-0.006699
2012	-0.014335	-0.036833	-0.001535
2013	0.013471	0.003789	0.005902
2014	0.005362	0.003563	0.001665
2015	0.018928	0.012333	0.007053
2016	0.010887	0.015869	0.005797
2017	-0.003506	0.003784	-0.000172
2018	-0.006523	-0.000680	-0.000572
2019	-0.002593	-0.014516	-0.000441
2020	-0.031300	-0.055703	-0.006881
2021	-0.000508	-0.018492	0.002118
2022	0.005848	-0.002957	0.002449
2023	0.009706	0.008433	0.002932
2024	0.000294	-0.025402	0.002757
2025	0.016226	0.017155	0.006635

Table 1: Seattle Mariners power metrics relative to league average (2008–2025)

9 Work Plan Evaluation

I followed my work plan almost exactly.

1. Install and figure out `pybaseball` indexing and manipulation ✓
2. Build function for Euler’s Method trajectory calculator and main method ✓
3. Build function and main method for traded player stats ✓
4. Build function for pulling weather data from `open-mateo` to use in trajectory calculator function ✓
5. Build function and main method for home/away hitting splits ✓

6. Go back and produce any more visualizations I will need ✓
7. Write docs for each function using markdown ✓
8. Write-up findings in $\text{\LaTeX}$ ✓

I followed these steps in this order and did not have to make any major changes.

10 Testing

I built testing files for the trajectory calculator which compares estimated hits based on input parameters to real hits by Julio Rodríguez. I implemented this function as a part of my EDA. The percent difference from the model to the the actual hit data was at the highest 1.4%. Since spin data is not reported publicly on batted ball data, I ran the model for a range of -100 to 200 rad/s which covers all of the acceptable range of spin on batted balls which should not exceed the spin on pitches (this range covers mid topspin to high backspin). From there I keep the model with the lowest percent error and that is what gets printed in main.

For testing the batting data code, I found and manually calculated all the data for Ken Griffey Jr., J. P. Crawford, and the 2025 Seattle Mariners using FanGraphs (not the same place as I am getting the data via `pybaseball`) just




Figure 7: This is the graph of all the simulated home-runs for Julio Rodríguez for 2026 that corresponds with the table. This plot also suffers from the axis being at different scales so I'm going to need to fix that at some point. It makes the hits look shallower than they actually are.

to make sure all the stats agree. I then tested it with assert statements in a testing.py file. For the calculated stats like SLG, BA, and OBP, I had to give the statements a range of values that were acceptable (the third decimal place) because these values often get rounded in computation at different points. It all checked out.

11 Citations

Aside from the references I have listed below which I can cite properly, I also consulted the `matplotlib` and `numpy` documentation briefly for things like `np.mean` and the trigonometry functions as well as `fig.suptitle()` and `plt.hist()`. I also consulted the `pybaseball` documentation which I listed above in the

dataset section. I consulted Google Maps and `MLB.com` for stadium information, specifically coordinate data. When pulling real data to test my functions, I consulted `baseballsavant.mlb.com` and `baseball-reference.com` which are both part of my data but to check my functions I had to actually go to their websites. For my second dataset I consulted the `open-meteo requests` documentation at <https://pypi.org/project/openmeteo-requests/> from which I got most of the structure for the `openmeteo-requests` syntax and structure in `air_densities.py`. I also looked at the `datetime` documentation because Open Meteo uses `datetime` objects <https://docs.python.org/3/library/datetime.html>. I got the venue IDs manually from <https://statsapi.mlb.com/api/v1/venues>. I also consulted <https://pypi.org/project/MLB-StatsAPI/> for the game start times data I got using `requests`. To do the `datetime` rounding I looked at the `datetime` docs and <https://www.geeksforgeeks.org/python/datetime-replace-function-in-python/>

References

Adair, R. K. (2002). *The Physics of Baseball: Third Edition, Revised, Updated, and Expanded*. HarperCollins.

Nathan, A. M. (2008a). The effect of spin-down on the flight of a baseball.

Nathan, A. M. (2008b). The effect of spin on the flight of a baseball. *American Journal of Physics*, 76(2):119–124.

Sawicki, G. S., Hubbard, M., and Stronge, W. J. (2003). How to hit home runs: Optimum baseball bat swing parameters for maximum range trajectories. *American Journal of Physics*, 71(11):1152–1162.

Slowinski, P. and Weinberg, N. (2010). Sample size. FanGraphs Sabermetrics Library. Originally published February 18, 2010; updated August 2017.