

Dynamical Matching in a Caldera Potential Energy surface

Matthaios Katsanikas

November 5, 2019

1 The Model

Dynamical matching is an interesting mechanism originally proposed in [2, 3] that arises in Caldera-type potential energy surfaces (PES). These potentials are relevant in Chemistry since they provide good approximations for many organic chemical reactions such as those that occur in the vinylcyclopropane-cyclopentene rearrangement [1, 7], the stereomutation of cyclopropane [4], the degenerate rearrangement of bicyclo[3.1.0]hex-2-ene [5, 6] or that of 5-methylene bicyclo[2.1.0]pentane [8]. The potential energy landscape of a Caldera is similar to that of a collapsed region in an erupted volcano. It is characterized by a shallow potential well region (a central minimum) surrounded by four entrance/exit channels in the form of index-1 saddles. Two of these saddles have low energy values and correspond to the formation of chemical products, while the other two are higher in energy and represent reactants. The dynamical matching phenomenon occurs when the Newtonian momentum conservation will favor that a trajectory entering the Caldera from a channel corresponding to a high energy index-1 saddle (reactant) exits through the diametrically opposing low energy index-1 saddle (product). Consequently, this mechanism will determine to a considerable extent the outcome of the chemical reaction.

References

- [1] J.E Baldwin. Thermal rearrangements of vinylcyclopropanes to cyclopentenenes. *Chemical reviews*, 103(4):1197–1212, 2003.
- [2] Barry K. Carpenter. Trajectories through an intermediate at a fourfold branch point. implications for the stereochemistry of biradical reactions. *Journal of the American Chemical Society*, 107(20):5730–5732, 1985.
- [3] Barry K. Carpenter. Dynamic matching: The cause of inversion of configuration in the [1,3] sigmatropic migration? *Journal of the American Chemical Society*, 117(23):6336–6344, 1995.
- [4] C. Doubleday, K. Bolton, and W.L. Hase. Direct dynamics study of the stereomutation of cyclopropane. *Journal of the American Chemical Society*, 119(22):5251–5252, 1997.




Figure 1: Caldera potential energy surface given by Eq. (??) for the model parameters $c_1 = 5$, $c_2 = 3$, $c_3 = -3/10$ and $\lambda = 1$.

- [5] C. Doubleday, M. Nendel, K.N. Houk, D. Thweatt, and M. Page. Direct dynamics quasiclassical trajectory study of the stereochemistry of the vinylcyclopropane - cyclopentene rearrangement. *Journal of the American Chemical Society*, 121(19):4720–4721, 1999.
- [6] C. Doubleday, C.P. Suhrada, and K.N. Houk. Dynamics of the degenerate rearrangement of bicyclo[3.1.0]hex-2-ene. *Journal of the American Chemical Society*, 128(1):90–94, 2006.
- [7] Z. Goldschmidt and B. Crammer. Vinylcyclopropane rearrangements. *Chem. Soc. Rev*, 17:229–267, 1988.
- [8] M.B Reyes, E.B Lobkovsky, and B.K Carpenter. Interplay of orbital symmetry and nonstatistical dynamics in the thermal rearrangements of bicyclo[n.1.0]polyenes. *J Am Chem Soc*, 124:641–651, 2002.