

Fundamentals of Ecology

Week 9, Ecology Lecture 7

Cara Brook

February 28, 2023

Let's recap a bit!

Let's recap a bit!

Disease Ecology

- We learned about extensions to the SIR model that can be made to better represent diverse types of pathogens (e.g. dynamics different from classic measles).

Let's recap a bit!

Disease Ecology

- We learned about extensions to the SIR model that can be made to better represent diverse types of pathogens (e.g. dynamics different from classic measles).
- These extensions included incorporating inherited maternal immunity into a disease model (important in the case of natural vs. vaccinated immunity in mothers), as well as modeling different types of pathogen transmission.

Let's recap a bit!

Disease Ecology

- We learned about extensions to the SIR model that can be made to better represent diverse types of pathogens (e.g. dynamics different from classic measles).
- These extensions included incorporating inherited maternal immunity into a disease model (important in the case of natural vs. vaccinated immunity in mothers), as well as modeling different types of pathogen transmission.
- Environmental disease transmission occurs when the pathogen is maintained outside the host in an environmental reservoir. We learned a few approaches to modeling this type of pathogen. Example: cholera, with water-borne transmission.

Let's recap a bit!

Disease Ecology

- We learned about extensions to the SIR model that can be made to better represent diverse types of pathogens (e.g. dynamics different from classic measles).
- These extensions included incorporating inherited maternal immunity into a disease model (important in the case of natural vs. vaccinated immunity in mothers), as well as modeling different types of pathogen transmission.
- Environmental disease transmission occurs when the pathogen is maintained outside the host in an environmental reservoir. We learned a few approaches to modeling this type of pathogen. Example: cholera, with water-borne transmission.
- Vertical transmission occurs when a pathogen is passed mother-to-infant *in utero*. We learned how to model this in the case of HIV- and also learned a few details about this pathogen.

Let's recap a bit!

Disease Ecology

- We learned about extensions to the SIR model that can be made to better represent diverse types of pathogens (e.g. dynamics different from classic measles).
- These extensions included incorporating inherited maternal immunity into a disease model (important in the case of natural vs. vaccinated immunity in mothers), as well as modeling different types of pathogen transmission.
- Environmental disease transmission occurs when the pathogen is maintained outside the host in an environmental reservoir. We learned a few approaches to modeling this type of pathogen. Example: cholera, with water-borne transmission.
- Vertical transmission occurs when a pathogen is passed mother-to-infant *in utero*. We learned how to model this in the case of HIV- and also learned a few details about this pathogen.
- Vector-borne transmission occurs when a pathogen is transmitted via blood-feeding arthropod (tick, flea, mosquito). We learned the specifics of how vector-borne disease are modeled in the case of malaria.

Let's recap a bit!

Disease Ecology

- We explored advances in malaria modeling, starting with the simplest Ross model, then adding a mosquito incubation period in the MacDonald model, then adding a human incubation period in the Anderson and May model.

Let's recap a bit!

Disease Ecology

- We explored advances in malaria modeling, starting with the simplest Ross model, then adding a mosquito incubation period in the MacDonald model, then adding a human incubation period in the Anderson and May model.
- Incorporation of the mosquito incubation period in the MacDonald model led to the insight that the survival of the female mosquito is a weak point in the malaria transmission cycle, which inspired decades of WHO malaria control via spraying of the insecticide DDT.

Let's recap a bit!

Disease Ecology

- We explored advances in malaria modeling, starting with the simplest Ross model, then adding a mosquito incubation period in the MacDonald model, then adding a human incubation period in the Anderson and May model.
- Incorporation of the mosquito incubation period in the MacDonald model led to the insight that the survival of the female mosquito is a weak point in the malaria transmission cycle, which inspired decades of WHO malaria control via spraying of the insecticide DDT.
- We learned terminology for zoonotic diseases, those transmitted from wildlife reservoirs to human spillover hosts.

Let's recap a bit!

Disease Ecology

- We explored advances in malaria modeling, starting with the simplest Ross model, then adding a mosquito incubation period in the MacDonald model, then adding a human incubation period in the Anderson and May model.
- Incorporation of the mosquito incubation period in the MacDonald model led to the insight that the survival of the female mosquito is a weak point in the malaria transmission cycle, which inspired decades of WHO malaria control via spraying of the insecticide DDT.
- We learned terminology for zoonotic diseases, those transmitted from wildlife reservoirs to human spillover hosts.
- Spillback occurs when pathogens transmit from the spillover host back to the reservoir that sourced the infection, as in the case of mink transmission of SARS-CoV-2 to humans.

Let's recap a bit!

Disease Ecology

- We explored advances in malaria modeling, starting with the simplest Ross model, then adding a mosquito incubation period in the MacDonald model, then adding a human incubation period in the Anderson and May model.
- Incorporation of the mosquito incubation period in the MacDonald model led to the insight that the survival of the female mosquito is a weak point in the malaria transmission cycle, which inspired decades of WHO malaria control via spraying of the insecticide DDT.
- We learned terminology for zoonotic diseases, those transmitted from wildlife reservoirs to human spillover hosts.
- Spillback occurs when pathogens transmit from the spillover host back to the reservoir that sourced the infection, as in the case of mink transmission of SARS-CoV-2 to humans.
- We can classify pathogens into stage I-V based on their R_0 in the human population. Zoonotic pathogens encompass stages II-IV.


Modeling **zoonotic pathogens** typically necessitates **modeling epidemic dynamics in multiple host populations.**

Modeling **zoonotic pathogens** typically necessitates **modeling epidemic dynamics in multiple host populations.**

Plague has added complexity as a vector-borne pathogen!


Modeling **zoonotic pathogens** typically necessitates **modeling epidemic dynamics in multiple host populations**.

Plague has added complexity as a vector-borne pathogen!



Modeling zoonotic pathogens typically necessitates modeling epidemic dynamics in multiple host populations.


Plague has added complexity as a vector-borne pathogen!



Humans get infected from contact with free-living infectious fleas!


Modeling zoonotic pathogens typically necessitates modeling epidemic dynamics in multiple host populations.

Plague has added complexity as a vector-borne pathogen!



How many equations would we need to model this system?

A simple plague model




$$\left. \begin{aligned} \frac{dS_R}{dt} &= b_R(S_R + I_R + R_R) - \beta_R S_R I_{FR} - \mu_R S_R \\ \frac{dI_R}{dt} &= \beta_R S_R I_{FR} - \mu_R I_R - \alpha_R I_R - \sigma_R I_R \\ \frac{dR_R}{dt} &= \sigma_R I_R - \mu_R R_R \end{aligned} \right\} \text{Rats}$$

$$\left. \begin{aligned} \frac{dS_F}{dt} &= b_F(S_F + I_{FR} + I_{FF}) - \beta_F S_F I_R - \mu_F S_F \\ \frac{dI_{FR}}{dt} &= \beta_F S_F I_R - \delta I_{FR} - \mu_F I_{FR} \\ \frac{dI_{FF}}{dt} &= \delta I_{FR} - \mu_F I_{FF} \end{aligned} \right\} \text{Fleas}$$

$$\left. \begin{aligned} \frac{dS_H}{dt} &= b_H(S_H + I_H + R_H) - \beta_H S_H I_{FF} - \mu_H S_H \\ \frac{dI_H}{dt} &= \beta_H S_H I_{FF} - \mu_H I_H - \alpha_H I_H - \sigma_H I_H \\ \frac{dR_H}{dt} &= \sigma_H I_H - \mu_H R_H \end{aligned} \right\} \text{Humans}$$


Fleas get infected from rats.
Humans get infected from free-living infected fleas!



“...from April 18 onwards, quantities of dead or dying rats were found in factories and warehouses...From the outer suburbs to the center of the town, in all the byways where the doctor's duties took him, in every thoroughfare, rats were piled up in garbage cans or lying in long lines in the gutters...On the fourth day the rats began to come out and die in batches...”

--*La Peste*, Albert Camus (1948)

Humans get infected from free-living infected fleas!




“...from April 18 onwards, quantities of dead or dying rats were found in factories and warehouses...From the outer suburbs to the center of the town, in all the byways where the doctor's duties took him, in every thoroughfare, rats were piled up in garbage cans or lying in long lines in the gutters...On the fourth day the rats began to come out and die in batches...”

--*La Peste*, Albert Camus (1948)


The force of infection (FOI), λ , is the rate at which susceptibles become infected

- For a single host pathogen: $\lambda = R_0 \frac{I}{N}$
- (FOI = $R_0 * \text{proportion infected}$)




The force of infection (FOI), λ , is the rate at which susceptibles become infected

- For a single host pathogen: $\lambda = R_0 \frac{I}{N}$
- (**FOI = $R_0 * \text{proportion infected}$**)
- \approx mirror image of $R_E = R_0 \frac{S}{N}$
- Also a finger on the epidemic pulse




The force of infection (FOI), λ , is the rate at which susceptibles become infected

- For a single host pathogen: $\lambda = R_0 \frac{I}{N}$
- (**FOI = $R_0 * \text{proportion infected}$**)
- \approx mirror image of $R_E = R_0 \frac{S}{N}$
- Also a finger on the epidemic pulse
- For multi-host pathogens, we can define the **spillover force of infection**



The force of infection (FOI), λ , is the rate at which susceptibles become infected


- For a single host pathogen: $\lambda = R_0 \frac{I}{N}$
- (**FOI = $R_0 * \text{proportion infected}$**)
- \approx mirror image of $R_E = R_0 \frac{S}{N}$
- Also a finger on the epidemic pulse
- For multi-host pathogens, we can define the **spillover force of infection**
 - Comprised of:
 - Infectiousness** of the reservoir
 - Reservoir-human **contact rates**
 - Susceptibility** in the human host



The force of infection (FOI), λ , is the rate at which susceptibles become infected

- For a single host pathogen: $\lambda = R_0 \frac{I}{N}$
- (**FOI = $R_0 * \text{proportion infected}$**)
- \approx mirror image of $R_E = R_0 \frac{S}{N}$
- Also a finger on the epidemic pulse
- For multi-host pathogens, we can define the **spillover force of infection**
 - Comprised of:
 - Infectiousness** of the reservoir
 - Reservoir-human **contact rates**
 - Susceptibility** in the human host

Keeling & Gilligan model λ_H for plague as proportional to the abundance of free-living infected fleas.



Zoonosis is a series of improbable events.


Animal ecology, population biology, biogeography, behavioural ecology, landscape ecology, agricultural sciences


Disease ecology, animal epidemiology, infectious disease dynamics, immunology, microbiology, veterinary medicine

Microbiology, disease ecology, vector ecology, epidemiology, spatial ecology, infectious disease dynamics

Human epidemiology, medical anthropology, vector ecology, social sciences, behavioural ecology, infectious disease dynamics


Microbiology, innate and adaptive immunology, cell biology of pathogen–host interactions, pathology, genetics, evolutionary biology





Bottlenecks to spillover.

Plowright et al. 2017. *Nature Reviews Microbiology*.




What's so special about bats?


Plowright et al. 2017. *Nature Reviews Microbiology*.


It is **debated** whether bats are reservoir hosts for **more zoonotic viruses** than other mammals.

Bats host a higher proportion of zoonotic viruses than other mammalian orders.




The number of zoonotic viruses hosted by bats scales with the total number of bat species in existence (>1400 spp.; 20% of mammalian diversity).






But it is known that bats do act as natural **reservoirs** and **maintenance hosts** for the majority of the world's **most virulent zoonotic viruses**.


Bats are natural reservoir hosts for several subgenera of *Alpha*- and *Betacoronaviruses*. *Rhinolophus* spp. bats are reservoirs for the SARS-like *Sarbecoviruses*.



Many CoVs transmit (and adapt) via **intermediate hosts** prior to infecting humans.
An intermediate host for SARS-CoV-2 has **not yet been identified**.








Bats likely sourced infections to both humans and pangolins in disparate spillover events (Scenario III)



Phylogenetic analysis suggests multiple spillover origins of SARS-CoV-2 to humans. Both lineages appear to have emerged from the Huanan Seafood Market.

Polytomy: a node on a phylogeny where more than 2 lineages descend from an ancestral lineage






But it is known that bats do act as natural **reservoirs** and **maintenance hosts** for the majority of the world's **most virulent zoonotic viruses**.

Virulence is the extent of a pathogen's harm to the host.

But it is known that bats do act as natural **reservoirs and maintenance hosts** for the majority of the world's **most virulent zoonotic viruses**.


Virulence is the extent of a pathogen's harm to the host.



Why do pathogens make us sick?

A virus will evolve to
maximize its capacity for
between-host infections (R_0).


*Why do pathogens
make us sick?*



Alizon et al. 2008. *J Evolutionary Biology*
Anderson and May 1982. *Parasitology*.


A virus will evolve to
maximize its capacity for
between-host infections (R_0).

*Why do pathogens
make us sick?*



A virus will evolve to
maximize its capacity for
between-host infections (R_0).


*Why do pathogens
make us sick?*



Mechanisms that promote
transmission may also enhance
virulence to the host.

Alizon et al. 2008. *J Evolutionary Biology*
Anderson and May 1982. *Parasitology*.

A virus will evolve to **maximize** its capacity for **between-host infections** (R_0).




Why do pathogens make us sick?

Virulence, then, is a by-product of a pathogen's need to transmit for reproduction!

Mechanisms that promote **transmission** may also enhance **virulence** to the host.

Alizon et al. 2008. *J Evolutionary Biology*
Anderson and May 1982. *Parasitology*.


A virus will evolve to **maximize** its capacity for **between-host infections** (R_0).



Mechanisms that promote **transmission** may also enhance **virulence** to the host.

Why do pathogens make us sick?


As a result, we predict the evolution of "**optimal virulence**."



$$R_0 = \frac{\beta(\text{virus density})}{\gamma + \mu + \alpha(\text{virus density})}$$

} infections created
} infections lost

A virus will evolve to **maximize** its capacity for **between-host infections** (R_0).



Mechanisms that promote **transmission** may also enhance **virulence** to the host.


Why do pathogens make us sick?

As a result, we predict the evolution of "**optimal virulence**."


Note that originally Anderson and May (1982) represented this link to virus density as acting on the disease recovery rate, though it is now more commonly expressed as a function of virulence!

$$R_0 = \frac{\beta(\text{virus density})}{\gamma + \mu + \alpha(\text{virus density})}$$


} infections created
} infections lost



For zoonoses,
**virulence and
transmission
tradeoff** to result in
total death burden.



For zoonoses,
**virulence and
transmission
tradeoff** to result in
total death burden.



The **virulence case study** of rabbit Myxoma virus

The **virulence case study** of rabbit Myxoma virus

- 1788: European rabbits brought to Australia as a food source
- Rabbits quickly became feral and numbers soared.
- 1901: Australia constructed the famous “rabbit-proof fence” to attempt to keep rabbits out of agriculture in the West.



The **virulence case study** of rabbit Myxoma virus

- 1788: European rabbits brought to Australia as a food source
- Rabbits quickly became feral and numbers soared.
- 1901: Australia constructed the famous “rabbit-proof fence” to attempt to keep rabbits out of agriculture in the West.
- Government looked to control measures, including biological controls in the 1930s.
- Tried Myxoma virus, a highly virulent European poxvirus infecting rabbits. with a CFR >99%.



Myxoma virus evolved to **intermediate virulence** in just a single year.

TABLE 4. THE VIRULENCE OF STRAINS OF MYXOMA VIRUS RECOVERED FROM THE FIELD IN AUSTRALIA BETWEEN 1951 AND 1981, EXPRESSED AS PERCENTAGES

virulence grade	I >99	II 95–99	III 70–95	IV 50–70	V <50	number of samples
case fatality rate (%)						
mean survival time/day	< 13	14–16	17–28	29–50	—	
1950–51†	100					1
1952–55†	13.3	20.0	53.3	13.3	0	60
1955–58†	0.7	5.3	54.6	24.1	15.5	432
1959–63‡	1.7	11.1	60.6	21.8	4.7	449
1964–66‡	0.7	0.3	63.7	34.0	1.3	306
1967–69‡	0	0	62.4	35.8	1.7	229
1970–74‡	0.6	4.6	74.1	20.7	0	174
1975–81§	1.9	3.3	67.0	27.8	0	212

† Data from Marshall & Fenner (1960).

‡ Data from Edmonds *et al.* (1975).


§ Data from J. W. Edmonds and R. C. H. Shepherd (personal communication, 1982).

|| Although only one strain was tested, the very high mortality rates in the initial outbreaks justify this extrapolation.

For Myxoma virus, **intermediate virulence evolution** resulted from **optimization of the tradeoffs between virulence and transmission**.



Rabbits around a waterhole in the myxomatosis trial site on Wardang Island, Australia, 1938



A SIMULATION MODEL OF THE POPULATION DYNAMICS
AND EVOLUTION OF MYXOMATOSIS¹

GREG DWYER

Department of Zoology, University of Washington, Seattle, Washington 98195 USA

SIMON A. LEVIN

Section of Ecology and Systematics, Corson Hall, Cornell University,
Ithaca, New York 14853 USA



LINDA BUTTEL

Ecosystems Research Center, Corson Hall, Cornell University,
Ithaca, New York 14853 USA

Dwyer, Levin, and Buttel. 1990.
Ecological Monographs.




Another classic **transmission-virulence tradeoff**: parasites of monarch butterflies

- Monarch butterflies infected with the protozoan parasite, *Ophryocystis elektroscirrha*, demonstrate reduced emergence and mating probabilities at higher parasite spore load (**virulence**).




Another classic **transmission-virulence tradeoff**: parasites of monarch butterflies

- Monarch butterflies infected with the protozoan parasite, *Ophryocystis elektroscirrha*, demonstrate reduced emergence and mating probabilities at higher parasite spore load (**virulence**).
- On the flip side, higher parasite loads also led to increased **transmission** through higher proportions of monarch eggs that acquired spores and higher numbers of parasites per egg and milkweed leaf, which increased the probability of infection.




Another classic **transmission-virulence tradeoff**: parasites of monarch butterflies

- Monarch butterflies infected with the protozoan parasite, *Ophryocystis elektroscirrha*, demonstrate reduced emergence and mating probabilities at higher parasite spore load (**virulence**).
- On the flip side, higher parasite loads also led to increased **transmission** through higher proportions of monarch eggs that acquired spores and higher numbers of parasites per egg and milkweed leaf, which increased the probability of infection.
- **Parasite fitness is calculated to be maximized at intermediate spore load.**




Why then are bat
viruses so virulent [to
humans]?





Why then are bat viruses so virulent [to humans]?

Mismatch of optimal virulence evolved in the **reservoir host** and a lack of **viral tolerance** in the spillover (human) host.



Bats appear **highly tolerant** of virus infection!
(They host viruses without experiencing disease)




Virus **tolerance** is linked to the **evolution of flight**.

Bats are the only **flying mammals**.



Flight is **more metabolically expensive** than
any form of terrestrial locomotion.


Bats have **elongated lifespans** despite **high metabolic rates**.



We hypothesize that:


unique **anti-**
inflammatory
pathways

evolved to
enable flight



We hypothesize that:


unique **anti-**
inflammatory
pathways




evolved to
enable flight




bat
longevity




anti-
cancer
defenses



virus
tolerance




A virus will evolve to
maximize its capacity for
between-host infections (R_0).




Mechanisms that promote
transmission may also enhance
virulence to the host.

A virus will evolve to **maximize** its capacity for **between-host infections** (R_0).




Mechanisms that promote **transmission** may also enhance **virulence** to the host.

Bats accrue **less virulence** for a given level of transmission.




A virus will evolve to **maximize** its capacity for **between-host infections** (R_0).




Mechanisms that promote **transmission** may also enhance **virulence** to the host.

Bats accrue **less virulence** for a given level of transmission.




Bat reservoirs may evolve **virus traits** that cause **virulence in non-bats**.

A virus will evolve to **maximize** its capacity for **between-host infections** (R_0).



Mechanisms that promote **transmission** may also enhance **virulence** to the host.

Bats accrue **less virulence** for a given level of transmission.




Bat reservoirs may evolve **virus traits** that cause **virulence in non-bats**.

Bat reservoirs may evolve **virus traits** that cause **virulence in non-bats**.




r^* within-host
virus growth
rate



Bat reservoirs may evolve **virus traits** that cause **virulence in non-bats**.




r^* within-host
virus growth
rate



Bat reservoirs may evolve **virus traits** that cause **virulence in non-bats**.



r^* within-host
virus growth
rate




Bat reservoirs may evolve **virus traits** that cause **virulence in non-bats**.





r^* within-host
virus growth
rate


Transmission
determined by
reservoir-evolved
virus growth rate



Reservoir



Spillover host



Limitations of the tradeoff model

- The ‘trade-off hypothesis’ offers an explanation for the disease inflicted by parasites and pathogens on their original hosts. While well-designed theoretically, it has not been historically well-supported empirically!




Limitations of the tradeoff model

- The ‘trade-off hypothesis’ offers an explanation for the disease inflicted by parasites and pathogens on their original hosts. While well-designed theoretically, it has not been historically well-supported empirically!
- This is partly due to challenges arising from the difficulty of measuring (and defining) transmission and virulence.
 - Virulence is a fitness cost that the parasite inflicts on the host, but these can take diverse forms, with differing consequences for the evolution of virulence.
 - For example: Fitness effects on reproduction vs. adult mortality
 - Sometimes, virulence is the result of the host’s immune response, rather than the direct impact of the parasite itself, further complicating dynamics

Limitations of the tradeoff model

- The ‘trade-off hypothesis’ offers an explanation for the disease inflicted by parasites and pathogens on their original hosts. While well-designed theoretically, it has not been historically well-supported empirically!
- This is partly due to challenges arising from the difficulty of measuring (and defining) transmission and virulence.
 - Virulence is a fitness cost that the parasite inflicts on the host, but these can take diverse forms, with differing consequences for the evolution of virulence.
 - For example: Fitness effects on reproduction vs. adult mortality
 - Sometimes, virulence is the result of the host’s immune response, rather than the direct impact of the parasite itself, further complicating dynamics

Sometimes, virulence is the result of the host's immune response, rather than the direct impact of the parasite itself



Self-reactive antibodies (ANA) **promote survival by downregulating worms in crash years** but **impede survival via immunopathology in peak years!**



Limitations of the tradeoff model

- The ‘trade-off hypothesis’ offers an explanation for the disease inflicted by parasites and pathogens on their original hosts. While well-designed theoretically, it has not been historically well-supported empirically!
- This is partly due to challenges arising from the difficulty of measuring (and defining) transmission and virulence.
 - Virulence is a fitness cost that the parasite inflicts on the host, but these can take diverse forms, with differing consequences for the evolution of virulence.
 - For example: Fitness effects on reproduction vs. adult mortality
 - Sometimes, virulence is the result of the host’s immune response, rather than the direct impact of the parasite itself, further complicating dynamics
 - In the case of zoonoses, the bulk of our measurements may be derived from a different host than the one in which the virus evolved

Limitations of the tradeoff model


- The ‘trade-off hypothesis’ offers an explanation for the disease inflicted by parasites and pathogens on their original hosts. While well-designed theoretically, it has not been historically well-supported empirically!
- This is partly due to challenges arising from the difficulty of measuring (and defining) transmission and virulence.
 - Virulence is a fitness cost that the parasite inflicts on the host, but these can take diverse forms, with differing consequences for the evolution of virulence.
 - For example: Fitness effects on reproduction vs. adult mortality
 - Sometimes, virulence is the result of the host’s immune response, rather than the direct impact of the parasite itself, further complicating dynamics
 - In the case of zoonoses, the bulk of our measurements may be derived from a different host than the one in which the virus evolved
- Many examples of cases in which transmission is decoupled from virulence, due to more complex transmission dynamics.
 - Ex: Ebola (transmission after death)
 - Ex: COVID (transmission high in the respiratory tract; morbidity low in the RT)



Limitations of the tradeoff model



- The ‘trade-off hypothesis’ offers an explanation for the disease inflicted by parasites and pathogens on their original hosts. While well-designed theoretically, it has not been historically well-supported empirically!
- This is partly due to challenges arising from the difficulty of measuring (and defining) transmission and virulence.
 - Virulence is a fitness cost that the parasite inflicts on the host, but these can take diverse forms, with differing consequences for the evolution of virulence.
 - For example: Fitness effects on reproduction vs. adult mortality
 - Sometimes, virulence is the result of the host’s immune response, rather than the direct impact of the parasite itself, further complicating dynamics
 - In the case of zoonoses, the bulk of our measurements may be derived from a different host than the one in which the virus evolved
- Many examples of cases in which transmission is decoupled from virulence, due to more complex transmission dynamics.
 - Ex: Ebola (transmission after death)
 - Ex: COVID (transmission high in the respiratory tract; morbidity low in the RT)
- A few specific applications of tradeoff theory...

Spatial structuring generally favors reduced pathogen virulence.




Spatial structuring generally favors reduced pathogen virulence.

High and low virulence are **both stable evolutionary outcomes** when pathogens generate acquired immunity in spatially structured populations.




Spatial structuring generally favors reduced pathogen virulence.



High and low virulence are **both stable evolutionary outcomes** when pathogens generate acquired immunity in spatially structured populations.



Here, **low virulence is favored in dense environments** and **high virulence in sparse environments**.



Imperfect vaccines can also support the evolution of virulence, depending on the lifestage of the pathogen that is targeted.



Is this ecology or evolution?


Is this ecology or evolution?

Phylogenetics unites both

Is this ecology or evolution? *Phyldynamics* unites both

Unifying the Epidemiological and Evolutionary
Dynamics of Pathogens

Bryan T. Grenfell,^{1,*} Oliver G. Pybus,² Julia R. Gog,¹ James L. N. Wood,³ Janet M. Daly,³ Jenny A. Mumford,³ Edward C. Holmes²



Population Biology


Conservation Biology

- Goal:
 - protect **populations** from extinction
- Concept:
 - **Minimum Viable Population** size (MVP)
- Approach:
 - protected area **reserves**

Single Large




Several Small




Disease Ecology


- Goal:
 - protect **populations** from disease via pathogen **extinction**
- Concept:
 - **Critical Community Size** (CCS)
- Approach:
 - sanitation
 - **vaccination**



Community = interacting populations of different species



How does fish abundance **vary** with changes in shark abundance?




Disease dynamics in the **broader community**

Disease dynamics in the **broader community**


Example: Lyme Disease

National Lyme disease risk map with four categories of risk






- Lyme disease is a vector-borne disease caused by the bacterium, *Borrelia burgdorferi*, vectored by *Ixodes* especially *Ixodes scapularis* ticks.
- Nymph ticks are borne in the spring, feed on small mammal hosts through the summer, then reproduce (particularly on deer) in the fall before going dormant in the winter.
- Human cases are largely concentrated in the spring and summer and result from infected tick bites.

Life Cycle of the *Ixodes scapularis* Tick



The **dilution effect** highlights buffering effects of biodiversity on disease transmission.

While a popular concept, it only holds in select cases!



In the case of Lyme, **increasing host biodiversity tends to be associated with decreasing Lyme prevalence.**

The **dilution effect** highlights buffering effects of biodiversity on disease transmission.

While a popular concept, it only holds in select cases!

