

# Introduction to Vector-Borne Diseases

UChicago Center in Paris

Paris, France

January 2025

# Goals for this lecture

- To introduce vector-borne diseases (VBD) broadly
- To introduce arboviruses specifically
- To describe the role of climate change in the expansion of arboviruses

# Goals for this lecture

- To introduce vector-borne diseases (VBD) broadly
- To introduce arboviruses specifically
- To describe the role of climate change in the expansion of arboviruses

# Mechanisms of disease transmission

- **Directly-transmitted** diseases – transmitted via exchange of bodily fluids
  - Droplet ( $> 5$  microns ) spread or direct contact, includes sexually-transmitted pathogens
  - Ex: Smallpox (*Variola* spp.), HIV, Mononucleosis (*Epstein Barr virus*)
- **Indirectly-transmitted** diseases – transmitted via droplets retained in air
  - Droplets  $< 5$  microns in diameter
  - Ex: Measles, COVID (*SARS-CoV-2*)
- **Vertically-transmitted** pathogens – transmitted mother-to-child *in utero*
  - Ex: HIV, *Herpes simplex virus*, *Cytomegalovirus*, Rubella, Zika
- **Environmentally-transmitted** pathogens – transmitted outside host (e.g. water, food)
  - Ex: Cholera (*Vibrio cholerae*), Salmonellosis (*Salmonella* spp. bacteria)
- **Vector-borne** diseases (a type of indirect transmission) are transmitted via blood-feeding arthropod (mosquitoes, ticks, fleas)
  - Ex: malaria, arboviruses (dengue, yellow fever), sleeping sickness, plague

Pathogens exhibit **diverse transmission mechanisms** that require tailored modeling structures

- **Vector-borne** diseases (a type of indirect transmission) are transmitted via blood-feeding arthropod (mosquitoes, ticks, fleas)
  - Euclidean **vector**: a quantity with a magnitude and direction  
→
  - Epidemiological **vector**: an agent that carries and transmits an infectious patient into another living organism



Pathogens exhibit **diverse transmission mechanisms** that require tailored modeling structures

- **Vector-borne** diseases (a type of indirect transmission) are transmitted via blood-feeding arthropod (mosquitoes, ticks, fleas)
  - Malaria: Mosquito-borne protozoan *Plasmodium spp.*
  - “Arboviruses”: Mosquito-borne viruses, including Dengue, Zika, Yellow fever virus, West Nile virus, Chikungunya virus
  - Sleeping sickness, also known as African trypanosomiasis: tsetse fly vector and protozoan pathogen (trypanosome)
  - Chagas disease: kissing bug vector and trypanosome pathogen
  - Plague: flea vector and bacterial pathogen (*Yersinia pestis*)

# Pathogens exhibit **diverse transmission mechanisms** that require tailored modeling structures

- **Vector-borne** diseases (a type of indirect transmission) are transmitted via blood-feeding arthropod (mosquitoes, ticks, fleas)
  - Malaria: Mosquito-borne protozoan *Plasmodium spp.*
  - “Arboviruses”: Mosquito-borne viruses, including Dengue, Zika, Yellow fever virus, West Nile virus, Chikungunya virus
  - Sleeping sickness, also known as African trypanosomiasis: tsetse fly vector and protozoan pathogen (trypanosome)
  - Chagas disease: kissing bug vector and trypanosome pathogen
  - **Plague**: flea vector and bacterial pathogen (*Yersinia pestis*) **Plague is BOTH vector-borne and zoonotic!**

# Pathogens exhibit **diverse transmission mechanisms** that require tailored modeling structures

- **Vector-borne** diseases (a type of indirect transmission) are transmitted via blood-feeding arthropod (mosquitoes, ticks, fleas)
  - **Malaria**: Mosquito-borne protozoan *Plasmodium spp.*
  - “Arboviruses”: Mosquito-borne viruses, including Dengue, Zika, Yellow fever virus, West Nile virus, Chikungunya virus
  - Sleeping sickness, also known as African trypanosomiasis: tsetse fly vector and protozoan pathogen (trypanosome)
  - Chagas disease: kissing bug vector and trypanosome pathogen
  - Plague: flea vector and bacterial pathogen (*Yersinia pestis*)




# Malaria

- 4 main human ***Plasmodium* parasites** (*falciparum*, *vivax*, *malariae*, *ovale*).
- Over 200 *Plasmodium* spp. globally, infecting birds, reptiles, and other mammals (rodents, bats, primates)

Distribution *Plasmodium falciparum*



Distribution *Plasmodium vivax*




# Malaria

- 4 main human **Plasmodium parasites** (*falciparum*, *vivax*, *malariae*, *ovale*).
- Over 200 *Plasmodium* spp. globally, infecting birds, reptiles, and other mammals (rodents, bats, primates)


Distribution *Plasmodium falciparum*



Distribution *Plasmodium vivax*




**Plasmodium parasites time RBC bursting such that they are at peak blood stage circulation when mosquito vectors are feeding at dusk!**




# Malaria has also shaped human DNA.

## Duffy antigen




- Modeled distribution of Duffy-negative human population

Distribution *Plasmodium falciparum*




Distribution *Plasmodium vivax*






# Malaria

- 4 main human ***Plasmodium* parasites** (*falciparum*, *vivax*, *malariae*, *ovale*).
- Over 200 *Plasmodium* spp. globally, infecting birds, reptiles, and other mammals (rodents, bats, primates)
- >400 global species of ***Anopheles* mosquito**, >100 that can transmit human malaria
- ~30-40 *Anopheles* spp. most commonly implicated in human malaria transmission!




# Malaria

- 4 main human ***Plasmodium* parasites** (*falciparum*, *vivax*, *malariae*, *ovale*).
- Over 200 *Plasmodium* spp. globally, infecting birds, reptiles, and other mammals (rodents, bats, primates)
- >400 global species of ***Anopheles* mosquito**, >100 that can transmit human malaria
- ~30-40 *Anopheles* spp. most commonly implicated in human malaria transmission!



Malaria has been eliminated from many regions where it was previously endemic, including the US.



Still one of the leading causes of child mortality globally – responsible for about half a million childhood deaths a year, 80% in Africa.

# Goals for this lecture





- To introduce vector-borne diseases (VBD) broadly
- To introduce arboviruses specifically
- To describe the role of climate change in the expansion of arboviruses


# Pathogens exhibit **diverse transmission mechanisms** that require tailored modeling structures





- **Vector-borne** diseases (a type of indirect transmission) are transmitted via blood-feeding arthropod (mosquitoes, ticks, fleas)
  - Malaria: Mosquito-borne protozoan *Plasmodium spp.*
  - “**Arboviruses**”: Mosquito-borne viruses, including Dengue, Zika, Yellow fever virus, West Nile virus, Chikungunya virus. ***Arbovirus is not a phylogenetic term!***
  - Sleeping sickness, also known as African trypanosomiasis: tsetse fly vector and protozoan pathogen (trypanosome)
  - Chagas disease: kissing bug vector and trypanosome pathogen
  - Plague: flea vector and bacterial pathogen (*Yersinia pestis*)







# Important arbovirus distributions


<b>Dengue virus</b>	
<i>Aedes</i> spp.	
<b>West Nile virus</b>	
<i>Culex pipiens</i>	
<b>Japanese encephalitis virus</b>	
<i>Culex pipiens</i>	
<b>Rift-Valley fever virus</b>	
<i>Culex</i> and <i>Aedes</i>	







<b>Yellow fever virus</b>		<i>Aedes</i> spp.
<b>Chikungunya virus</b>		<i>Aedes</i> spp.
<b>Venezuelan equine encephalitis virus</b>		<i>Culex</i> -> <i>Aedes</i>
<b>Crimean-Congo hemorrhagic fever virus</b>		<i>Hyalomma</i> ticks

# Important arbovirus distributions





<i>Aedes</i> spp.	<b>Dengue virus</b> 
<i>Culex pipiens</i>	<b>West Nile virus</b> 
<i>Culex pipiens</i>	<b>Japanese encephalitis virus</b> 
<i>Culex</i> and <i>Aedes</i>	<b>Rift-Valley fever virus</b> 








<b>Yellow fever virus</b> 	<i>Aedes</i> spp.
<b>Chikungunya virus</b> 	<i>Aedes</i> spp.
<b>Venezuelan equine encephalitis virus</b> 	<i>Culex</i> -> <i>Aedes</i>
<b>Crimean-Congo hemorrhagic fever virus</b> 	<i>Hyalomma</i> ticks



# Important arbovirus distributions

<b>Dengue virus</b>	
<i>Aedes</i> spp.	
<b>West Nile virus</b>	
<i>Culex pipiens</i>	
<b>Japanese encephalitis virus</b>	
<i>Culex pipiens</i>	
<b>Rift-Valley fever virus</b>	
<i>Culex</i> and <i>Aedes</i>	





<b>Yellow fever virus</b>		<i>Aedes</i> spp.
<b>Chikungunya virus</b>		<i>Aedes</i> spp.
<b>Venezuelan equine encephalitis virus</b>		<i>Culex</i> -> <i>Aedes</i>
<b>Crimean-Congo hemorrhagic fever virus</b>		<i>Hyalomma</i> ticks


# Important arbovirus distributions

*Aedes* spp.



Dengue virus




Yellow fever virus



*Aedes* spp.



Chikungunya virus



*Aedes* spp.



*Culex pipiens*

West Nile virus




*Culex pipiens*

Japanese encephalitis virus





Venezuelan equine encephalitis virus




*Culex* -> *Aedes*

*Culex* and *Aedes*

Rift-Valley fever virus




Crimean-Congo hemorrhagic fever virus





*Hyalomma* ticks

# Arboviruses infect a wide range of hosts and vectors






# Is flavivirus pathogenesis correlated with vector identity?



# Dengue vaccines


- Two licensed vaccines in circulation – both tetravalent live attenuated vaccines which confer immunity to all 4 serotypes simultaneously
- Dengvaxia – by Sanofi Pasteur
  - First licensed in 2016 but controversy developed after severe cases developed in those who were previously naïve in Philippines
  - Now recommended only in those who test seropositive
  - This policy has greatly diminished demand. Vaccine no longer widely used.
- Qdenga – TAK-003 – by Takeda
  - Pre-qualified for use in May 2024
  - Appears to be efficacious in both seronegative/seropositive individuals, though is ineffective at preventing infection for DENV-3/4 in seronegatives and lack of enhanced pathogenicity cannot be ruled out.
  - WHO recommends use in children 6-16 in high transmission settings only at this stage.
- NIH NIAID vaccine **TV005** also showing promise but not yet licensed.



# Global distribution of *Aedes aegypti*





# Global distribution of *Aedes albopictus*






Detailed probability of occurrence of *Aedes albopictus* in Europe and US, areas where the mosquito is most rapidly expanding its range


# Dengue fever infections, 2019

Estimated annual number of new dengue infections. Dengue is a viral infection transmitted through the bite of infected mosquitoes; symptoms include fever, headaches, and nausea. Most infections are asymptomatic or mild, but dengue can occasionally be severe or fatal.





Zika is also vectored by *Aedes aegypti* and its relatives











# Goals for this lecture

- To introduce vector-borne diseases (VBD) broadly
- To introduce arboviruses specifically
- To describe the role of climate change in the expansion of arboviruses





# Arboviruses and climate change

- Mosquito development rates are highly sensitive to temperature.
- As a result, many arboviruses are climate-constrained in distribution but ranges are expanding with climate change.

# Thermal performance curves for *Aedes aegypti*




Different vectors respond to temperature in different ways.





These vector differences have important consequences for disease dynamics.



These vector differences have important consequences for disease dynamics.

