


Seed dispersal by birds and lemurs in fragmented landscape



Nancia Raoelinjanakolona, University of Antananarivo

Background

Lemurs and birds: main seed dispersers in many plant species in Malagasy ecosystems

Species are facing a high risk of decline by forest fragmentation



???STATISTICAL MODEL


What factors influence the consumption of fruit by lemurs and birds in plant species?

??? DYNAMICAL MODEL

How does competition between birds and lemurs affect seed dispersal dynamics?



Acknowledgments : All INSTRUCTORS, Marius, Vero, Kim & Angelo

What factors influence the consumption of fruit by lemurs and birds



- Hypothesis: Color, size and shape of fruit influence the consumption of fruit by birds and lemurs
- Response variable: Eaten fruit by birds/lemurs per tree
- Predictor variables: Fruit traits: color, size, shape; frugivores traits: size, handling fruit, sites in both edge and interior of forest
- Family: “poisson”
- Link: log
- Rcode: `glmer(Fruit_number~color+size+shape+handling,family=poisson(link = “log”), data=tree.observation)`

How does competition between birds and lemurs affect seed dispersal dynamics?



β_B : death rate of bird
 β_L : death rate of lemur
 μ : fructification rate
 ω : feeding rate
 f_b : feeding rate by bird
 f_l : feeding rate by lemur
 B : number of bird
 L : number of lemurs

$$S_{t+1}: \mu(ST)(SD) - \omega(ST_t) [(f_{bt} * B_t) + ((1-f_{bt}) * L_t)]$$

$$SD_{t+1}: \omega(ST_t) [(f_{bt} * B_t) + ((1-f_{bt}) * L_t)]$$

What would be the future of the tree species if this competition disrupt by forest fragmentation?

- Collect data in the field: animal survey and camera trapping to estimate the diversity and abundance of birds and lemurs in the forest
- Built a model: simulation of the abundance of frugivores through time
- Built a model: simulation of loss of one frugivore (bird or lemur) or frugivore lower abundance through time