

# Relaxation Time

Ben Rosemeyer

July 18, 2013

## Spin Susceptibility

The presence of a magnetic field introduces a potential for particles with spin  $V = -\vec{m} \cdot \vec{H}$ . The magnetization due to this potential is given by  $M_\alpha(t) = -i \int_{-\infty}^t < [m_\alpha(t), V(t')] > dt'$ , and the magnetic susceptibility is  $\chi_{\alpha,\beta}(x, x', t) = i \int_{-\infty}^t < [m_\alpha(x, t), m_\beta(x', t')] > dt'$ . The magnetic moment is given by  $m_\alpha(x, t) = \mu_e \sum_{s,s'} \sigma_{s,s'}^\alpha \psi_s^\dagger(x, t) \psi_{s'}(x, t)$  (Mahan). Now we can proceed to calculate the susceptibility. In the case of uniform time, we can assume that the product  $m_\alpha(x, t), m_\beta(x', t')$  is a function of  $\tau = t - t'$ . In this limit we get:

$$\begin{aligned} \chi_{\alpha,\beta}(x, x', t) &= i \int_{-\infty}^t < [m_\alpha(x, 0), m_\beta(x', t')] > dt' \\ &= -i\mu_e^2 \sum_{s,s',t,t'} \sigma_{s,s'}^\beta \sigma_{t,t'}^\alpha \int_{-\infty}^t < [\psi_s^\dagger(x', t') \psi_{s'}(x' t'), \psi_t^\dagger(x, 0) \psi_t(x, 0)] > \end{aligned}$$

Using the Bogoliubov transformation and fourier transforming this to momentum and energy phase space we have:

$$\begin{aligned} \chi_{xx}(q, \omega) &= -\mu_B \sum_{s,k} \frac{f_{k,s} - f_{k+q,\bar{s}}}{\omega + E_{k,s} - E_{k+q,\bar{s}} + i\Gamma} (u_k^2 u_{k+q}^2 + u_k v_k u_{k+q} v_{k+q}) \\ &\quad - \frac{f_{k,\bar{s}} - f_{k+q,s}}{\omega - E_{k,\bar{s}} + E_{k+q,s} + i\Gamma} (v_k^2 v_{k+q}^2 + u_k v_k u_{k+q} v_{k+q}) \\ &\quad + \frac{1 - f_{k,\bar{s}} - f_{k+q,\bar{s}}}{\omega - E_{k,\bar{s}} - E_{k+q,\bar{s}} + i\Gamma} (v_k^2 u_{k+q}^2 - u_k v_k u_{k+q} v_{k+q}) \\ &\quad - \frac{1 - f_{k,s} - f_{k+q,s}}{\omega + E_{k,s} + E_{k+q,s} + i\Gamma} (u_k^2 v_{k+q}^2 - u_k v_k u_{k+q} v_{k+q}) \end{aligned}$$

The real part ( $\Gamma = 0$ ) in the  $\omega = 0$  limit is:

$$\chi'_{xx}(q) = -\mu_B \sum_{s,k} \frac{f_{k,s} - f_{k+q,\bar{s}}}{E_{k,s} - E_{k+q,\bar{s}}} (u_k u_{k+q} + v_k v_{k+q})^2 - \frac{1 - f_{k,s} - f_{k+q,s}}{E_{k,s} + E_{k+q,s}} (v_k u_{k+q} - u_k v_{k+q})^2$$

The imaginary part for finite  $\omega$  is:

$$\begin{aligned}
\chi''_{xx}(q, \omega) = \pi\mu_B \sum_{s,k} & (f_{k,s} - f_{k+q,\bar{s}})\delta(\omega + E_{k,s} - E_{k+q,\bar{s}})(u_k^2 u_{k+q}^2 + u_k v_k u_{k+q} v_{k+q}) \\
& - (f_{k,\bar{s}} - f_{k+q,s})\delta(\omega - E_{k,\bar{s}} + E_{k+q,s})(v_k^2 v_{k+q}^2 + u_k v_k u_{k+q} v_{k+q}) \\
& + (1 - f_{k,\bar{s}} - f_{k+q,\bar{s}})\delta(\omega - E_{k,\bar{s}} - E_{k+q,\bar{s}})(v_k^2 u_{k+q}^2 - u_k v_k u_{k+q} v_{k+q}) \\
& - (1 - f_{k,s} - f_{k+q,s})\delta(\omega + E_{k,s} + E_{k+q,s})(u_k^2 v_{k+q}^2 - u_k v_k u_{k+q} v_{k+q})
\end{aligned}$$

Now we use the derived formula for the relaxation time:

$$\frac{1}{TT_1} \propto \sum_q \left[ \frac{\chi''(q, \omega)}{\omega} \right]_{\omega \rightarrow 0}$$

$$\begin{aligned}
\frac{\chi''_{xx}(q, \omega)}{\omega} = \pi\mu_B \sum_{s,k} & \frac{f_{k,s} - f_{k+q,\bar{s}}}{-E_{k,s} + E_{k+q,\bar{s}}} \delta(\omega + E_{k,s} - E_{k+q,\bar{s}})(u_k^2 u_{k+q}^2 + u_k v_k u_{k+q} v_{k+q}) \\
& - \frac{f_{k,\bar{s}} - f_{k+q,s}}{E_{k,\bar{s}} - E_{k+q,s}} \delta(\omega - E_{k,\bar{s}} + E_{k+q,s})(v_k^2 v_{k+q}^2 + u_k v_k u_{k+q} v_{k+q}) \\
& + \frac{1 - f_{k,\bar{s}} - f_{k+q,\bar{s}}}{E_{k,\bar{s}} + E_{k+q,\bar{s}}} \delta(\omega - E_{k,\bar{s}} - E_{k+q,\bar{s}})(v_k^2 u_{k+q}^2 - u_k v_k u_{k+q} v_{k+q}) \\
& - \frac{1 - f_{k,s} - f_{k+q,s}}{-E_{k,s} - E_{k+q,s}} \delta(\omega + E_{k,s} + E_{k+q,s})(u_k^2 v_{k+q}^2 - u_k v_k u_{k+q} v_{k+q})
\end{aligned}$$

$$\begin{aligned}
\left[ \frac{\chi''_{xx}(q, \omega)}{\omega} \right]_{\omega \rightarrow 0} = \pi\mu_B \sum_{s,k} & - \frac{f_{k,s} - f_{k+q,\bar{s}}}{E_{k,s} - E_{k+q,\bar{s}}} \delta(E_{k,s} - E_{k+q,\bar{s}})(u_k u_{k+q} + v_k v_{k+q})^2 \\
& + \frac{1 - f_{k,s} - f_{k+q,s}}{E_{k,s} + E_{k+q,s}} \delta(E_{k,s} + E_{k+q,s})(v_k u_{k+q} - u_k v_{k+q})^2
\end{aligned}$$

Now we can plug this into the equation for the relaxation time and change coordinates from  $(k, q)$  to  $(k+q/2, k-q/2)$  and define  $(k+q/2, k-q/2) \Rightarrow (k, p)$ :

$$\frac{1}{TT_1} \propto \sum_{s,k,p} - \frac{f_{p,s} - f_{k,\bar{s}}}{E_{p,s} - E_{k,\bar{s}}} \delta(E_{p,s} - E_{k,\bar{s}})(u_p u_k + v_p v_k)^2 + \frac{1 - f_{p,s} - f_{k,s}}{E_{p,s} + E_{k,s}} \delta(E_{p,s} + E_{k,s})(v_p u_k - u_p v_k)^2$$

$$\frac{1}{TT_1} \propto - \sum_{s,k,p} \frac{f(E_{p,s}) - f(E_{k,\bar{s}})}{E_{p,s} - E_{k,\bar{s}}} \delta(E_{p,s} - E_{k,\bar{s}})(u_p u_k + v_p v_k)^2 + \frac{f(E_{p,s}) - f(-E_{k,s})}{E_{p,s} - (-E_{k,s})} \delta(E_{p,s} - (-E_{k,s}))(v_p u_k - u_p v_k)^2$$

$$\begin{aligned}
\frac{1}{TT_1} \propto - \sum_s \int & k dk d\theta_k p dp d\theta_p \frac{f(E_{p,s}) - f(E_{k,\bar{s}})}{E_{p,s} - E_{k,\bar{s}}} \delta(E_{p,s} - E_{k,\bar{s}})(u_p u_k + v_p v_k)^2 \\
& + \frac{f(E_{p,s}) - f(-E_{k,s})}{E_{p,s} - (-E_{k,s})} \delta(E_{p,s} - (-E_{k,s}))(v_p u_k - u_p v_k)^2
\end{aligned}$$

Now we wish to transform this integral into energy using the dispersion for quasi-particle energies  $E_{p,s} = \sqrt{(|p|^2 - 1)^2 + \Delta_{\theta_p}^2} + s\mu_e H$ . We use the Jacobian determinant to perform this transformation:

$$\begin{aligned}
dE_{p,s}d\theta &= \left| \det \begin{pmatrix} \frac{dE_{p,s}}{dp} & \frac{dE_{p,s}}{d\theta} \\ \frac{d\theta}{dp} & \frac{d\theta}{d\theta} \end{pmatrix} \right| dp d\theta \\
&= \frac{dE_{p,s}}{dp} dp d\theta = \left| \frac{2(p^2 - 1)p}{E_{p,s} - s\mu_e H} \right| dp d\theta \\
\Rightarrow pdpd\theta &= \frac{|E_{p,s} - s\mu_e H|}{2\sqrt{(E_{p,s} - s\mu_e H)^2 - \Delta_\theta^2}} dE_{p,s}d\theta = N(E_{p,s} - s\mu_e H, \theta) dE_{p,s}d\theta
\end{aligned}$$

We proceed with the transformation by defining the following variables. In the first term  $(E_1, \theta_1) = (E_{p,s}, \theta_p)$ ,  $(E_2, \theta_2) = (E_{k,\bar{s}}, \theta_k)$ , and in the second term,  $(E_1, \theta_1) = (E_{p,s}, \theta_p)$ ,  $(E_2, \theta_2) = (E_{k,s}, \theta_k)$

$$\begin{aligned}
\frac{1}{TT_1} &\propto - \sum_s \int d\theta_1 d\theta_2 dE_1 dE_2 \frac{f(E_1) - f(E_2)}{E_1 - E_2} \delta(E_1 - E_2) (u(E_1 - sb, \theta_1)u(E_2 + sb, \theta_2) + v(E_1 - sb, \theta_1)v(E_2 + sb, \theta_2))^2 N(E_1 - E_2) \\
&\quad + \frac{f(E_1) - f(-E_2)}{E_1 - (-E_2)} \delta(E_1 - (-E_2)) (v(E_1 - sb, \theta_1)u(E_2 - sb, \theta_2) - u(E_1 - sb, \theta_1)v(E_2 - sb, \theta_2))^2 N(E_1 + E_2)
\end{aligned}$$

Now we use the delta functions on the fermi functions and the denominators to get the following (we keep the subscripts on the u's and v's in order to better understand how the coherence factors reduce)

$$\begin{aligned}
\frac{1}{TT_1} &\propto - \sum_s \int d\theta_1 d\theta_2 dE_1 \frac{df}{dE} \Big|_{E_1} (u_1 u_2 + v_1 v_2)^2 N(E_1 - sb, \theta_1) N(E_1 + sb, \theta_2) \\
&\quad + \frac{df}{dE} \Big|_{E_1} (v_1 u_2 - u_1 v_2)^2 N(E_1 - sb, \theta_1) N(-E_1 - sb, \theta_2) \\
&= - \sum_s \int d\theta_1 d\theta_2 dE_1 \frac{df}{dE} \Big|_{E_1} N(E_1 - sb, \theta_1) N(E_1 + sb, \theta_2) [(u_1 u_2 + v_1 v_2)^2 + (v_1 u_2 - u_1 v_2)^2] \\
&= - \sum_s \int d\theta_1 d\theta_2 dE_1 \frac{df}{dE} \Big|_{E_1} N(E_1 - sb, \theta_1) N(E_1 + sb, \theta_2) [u_1^2(u_2^2 + v_2^2) + v_1^2(v_2^2 + u_2^2)] \\
&= - \sum_s \int d\theta_1 d\theta_2 dE_1 \frac{df}{dE} \Big|_{E_1} N(E_1 - sb, \theta_1) N(E_1 + sb, \theta_2) \\
&= -2 \int dE \frac{df}{dE} \bar{N}(E - b) \bar{N}(E + b)
\end{aligned}$$

Where we have defined  $b = \mu_e H$  and

$$\bar{N}(E) = \int d\theta \frac{|E|}{2\sqrt{E^2 - \Delta_\theta^2}}$$