

Homework 9 in L^AT_EX

Robert Brothers Mechanical Engineering Student @ UTSA

November 9, 2014

1 Question 1: The Lagrangian and Equations of Motion

1.1 (a) Write an Expression for the Lagrangian of the Particle

$$L = -1.0Rgm_1 \sin(q_1) + 0.5\dot{q}_1^2 (I_1 + R^2 m_1)$$

1.2 (b) Find the Equation of Motion of the Particle

$$\tau_k = 1.0I_1\ddot{q}_1 + 1.0R^2m_1\ddot{q}_1 + 1.0Rgm_1 \cos(q_1)$$

2 Question 2: Equations of Motion of a Two Link Manipulator

2.1 (a) Find the Lagrangian

$$L = 0.5I_1\dot{q}_1^2 + 2.0gm_2q_2 \cos(q_1) + 0.5m_2\dot{q}_1^2 + 6.12323399573677 \cdot 10^{-17}m_2\dot{q}_1\dot{q}_2 + 0.5m_2\dot{q}_2^2$$

2.2 (b) Find Equation of Motion

$$\begin{aligned}\tau_k = & 2.0gm_2q_2 \sin(q_1) - 2.0gm_2 \cos(q_1) \\ & + 1.0m_2\ddot{q}_2 + 1.0\ddot{q}_1(I_1 + m_2)\end{aligned}$$

3 Question 3: Equations of Motion of a Pendulum

3.1 (a) Find the Lagrangian of the Pendulum

$$L = l_1m_2(-1.0g \sin(q_1) + 0.5l_1\dot{q}_1^2) \quad (1)$$

3.2 (b) Find Equation of Motion

$$\tau_k = 1.0l_1m_2(g \cos(q_1) + l_1\ddot{q}_1) \quad (2)$$

```

\documentclass[10pt]{article}
\usepackage{setspace}
\usepackage{amsmath,amsfonts,amsthm,amssymb}
\usepackage{color}
\usepackage{fancyhdr}
\usepackage{chngpage}
\usepackage{enumerate}
\usepackage{graphicx}
\usepackage{boxedminipage}

\title{Homework 9 in \LaTeX}
\author{Robert Brothers Mechanical Engineering Student @ UTSA}
\begin{document}
\doublespacing
\footnotesize
\maketitle
\date

\section{Question 1: The Lagrangian and Equations of Motion}
\subsection{(a) Write an Expression for the Lagrangian of the Particle}
\begin{equation}
\begin{align}
L &= -1.0 R g m_{\{1\}} \sin{\left( q_{\{1\}} \right)} \nonumber \\
&\quad + 0.5 \dot{q}_{\{1\}}^2 \left( I_{\{1\}} + R^2 m_{\{1\}} \right) \nonumber
\end{align}
\end{equation}

\subsection{(b) Find the Equation of Motion of the Particle}
\begin{equation}
\begin{align}
\tau_{\{k\}} &= 1.0 I_{\{1\}} \ddot{q}_{\{1\}} \nonumber \\
&\quad + 1.0 R^2 m_{\{1\}} \ddot{q}_{\{1\}} \nonumber \\
&\quad + 1.0 R g m_{\{1\}} \cos{\left( q_{\{1\}} \right)} \nonumber
\end{align}
\end{equation}

\section{Question 2: Equations of Motion of a Two Link Manipulator}
\subsection{(a) Find the Lagrangian}
\begin{equation}
\begin{align}
L &= 0.5 I_{\{1\}} \dot{q}_{\{1\}}^2 + 2.0 g m_{\{2\}} q_{\{2\}} \cos{\left( q_{\{1\}} \right)} \nonumber \\
&\quad + 0.5 m_{\{2\}} \dot{q}_{\{1\}}^2 + 6.12323399573677 \cdot 10^{-17} m_{\{2\}} \dot{q}_{\{1\}} \dot{q}_{\{2\}} \nonumber \\
&\quad + 0.5 m_{\{2\}} \dot{q}_{\{2\}}^2 \nonumber
\end{align}
\end{equation}

\subsection{(b) Find Equation of Motion}
\begin{equation}
\begin{align}
\tau_{\{k\}} &= 2.0 g m_{\{2\}} q_{\{2\}} \sin{\left( q_{\{1\}} \right)} - 2.0 g m_{\{2\}} \cos{\left( q_{\{1\}} \right)} \nonumber \\
&\quad + 1.0 m_{\{2\}} \ddot{q}_{\{2\}} + 1.0 \ddot{q}_{\{1\}} \left( I_{\{1\}} + m_{\{2\}} \right) \nonumber
\end{align}
\end{equation}

\section{Question 3: Equations of Motion of a Pendulum}
\subsection{(a) Find the Lagrangian of the Pendulum}
\begin{equation}
\begin{align}
L &= l_{\{1\}} m_{\{2\}} \left( -1.0 g \sin{\left( q_{\{1\}} \right)} + 0.5 l_{\{1\}} \dot{q}_{\{1\}}^2 \right) \nonumber
\end{align}
\end{equation}

\subsection{(b) Find Equation of Motion}
\begin{equation}
\begin{align}
\tau_{\{k\}} &= 1.0 l_{\{1\}} m_{\{2\}} \left( g \cos{\left( q_{\{1\}} \right)} + l_{\{1\}} \ddot{q}_{\{1\}} \right) \nonumber
\end{align}
\end{equation}

```

`\end{document}`

```
#!/usr/bin/env python
```

```
import sys
sys.path.append(r"/Users/robertbrothers/Desktop/Fall 2014/Fundamentals_of_Robotics/r
obo_git/python/")
import robotics_functions as rf, sympy as sy, numpy as np

[R, t1, I1, m1, q1, qdot1, qddot1, I2] = sy.symbols("R t1 I1 m1 q1 qdot1 qddot1 I2")

link_list_cm = [[
    [ R, 0, 0, q1],
    ],[
        sy.Matrix(
            [
                [0], [0], [0], [1]
            ]
        )
    ]
]

[link_list, ocm] = link_list_cm
A0n = rf.sym_get_A0n(link_list)
J = rf.sym_pt_jacobian(link_list_cm)
qdot = sy.Matrix([
    [qdot1]
])

M = [
    sy.Matrix([
        [m1,0,0],
        [0,m1,0],
        [0,0,m1]
    ])
]

I = [
    sy.Matrix([
        [I1,0,0],
        [0,I1,0],
        [0,0,I1]
    ])
]

q = sy.Matrix([
    [q1],
])

tdv_vec = [
    (q1, qdot1),
    (qdot1, qddot1),
]

print "A matrices"
for A in A0n:
    sy.pprint(sy.trigsimp(A))
print "A Jacobian"
for j in J:
    sy.pprint(sy.trigsimp(j))
print "a) expression for the Lagrangian of the Particle: \n"
print sy.printing.latex(sy.trigsimp(rf.sym_lagrangian( link_list_cm, M, I, qdot)))
print "\n"
print "b) Find the Equations of Motion of the Particle: \n"
print sy.printing.latex(sy.simplify(sy.trigsimp(rf.sym_torque(link_list_cm, M, I, qdot, q, tdv_vec))))

[l1, l2, l3, t1, t2, t3, a1, a2, a3, d1, d2, d3] = sy.symbols("l1 l2 l3 t1 t2 t3 a1
a2 a3 d1 d2 d3")
[q1, q2, qdot1, qdot2, qddot1, qddot2, m1, m2, r1, r2] = sy.symbols("q1 q2 qdot1 qdo
t2 qddot1 qddot2 m1 m2 r1 r2")

link_list_cm = [[
    [0,np.pi/2, 0, q1],
    [0,0,q2,0],
    ],[
        sy.Matrix([
```

```

        [0],[0],[0],[1]]),
        sy.Matrix([
            [0],[0],[q2],[1]])
    ]
]

m = np.array([m1, m2])
l = np.array([l1, l2])
r = np.array([r1, r2])

M = [sy.Matrix([
    [m[i],0,0],
    [0,m[i],0],
    [0,0,m[i]]
]) for i in range(len(m))]

I = [sy.Matrix([
    [I1,0,0],
    [0,I1, 0],
    [0, 0,I1]
]),
sy.Matrix([
    [I2,0,0],
    [0,I2,0],
    [0,0,I2]
])
]
q = sy.Matrix([
    [q1],
    [q2]
])
qdot = sy.Matrix([
    [qdot1],
    [qdot2]
])
tdv_vec = [
    (qdot1,qddot1),
    (qdot2,qddot2),
    (q1, qdot1),
    (q2, qdot2),
]
link_list = link_list_cm[0]
A0n = rf.sym_get_A0n(link_list)
J = rf.sym_pt_jacobian(link_list_cm)

print "A matrices"
for A in A0n:
    sy.pprint(sy.trigsimp(A))

print "A Jacobian"
for j in J:
    sy.pprint(sy.trigsimp(j))
print "jacobian"
sy.pprint(rf.sym_pt_jacobian(link_list_cm))
print "\nLagrangian of the Manipulator"
print sy.printing.latex(sy.simplify(sy.trigsimp(rf.sym_lagrangian(link_list_cm, M, I
, qdot)[0])))

print "\nEquations of Motion of the Two Link Manipulator"
print sy.printing.latex(sy.simplify(sy.trigsimp(rf.sym_torque(link_list_cm, M, I, qd
ot, q, tdv_vec))))

link_list_cm = [[
    [l1, 0, 0, q1],
    ],[sy.Matrix([
        [0],[0],[0],[1]])
    ]
]

q = sy.Matrix([
    [q1],

```

```

    ])

qdot = sy.Matrix([
    [qdot1],
    ])
tdv_vec = [
    (qdot1,qddot1),
    (q1, qdot1),
    ]

M = [sy.Matrix([
    [m[i],0,0],
    [0,m[i],0],
    [0,0,m[i]]
    ])
    ]

I = [sy.Matrix([
    [0,0,0],
    [0,0,0],
    [0,0,0]
    ])
    ]

print "\nLagrangian of the pendulum"
print sy.printing.latex(sy.simplify(sy.trigsimp(rf.sym_lagrangian(link_list_cm, M, I
, qdot)[0])))

print "\nEquations of Motion of the Two Link Manipulator"
print sy.printing.latex(sy.simplify(sy.trigsimp(rf.sym_torque(link_list_cm, M, I, qd
ot, q, tdv_vec))))

```