

# Big Titles for \$400

Author one<sup>1</sup> and Author two<sup>2</sup>

<sup>1</sup>NOAA Great Lakes Environmental Research Lab, Ann Arbor, MI

<sup>2</sup>Cooperative Institute for Great Lakes Research, University of Michigan, Ann Arbor, MI

## Abstract

## 1 Introduction

Lake ice in the Great Lakes is highly variable and has impacts ranging from economy to ecology. The commercial shipping industry in the Great Lakes transports 135 million tons of cargo each year and supports a quarter million jobs (Martin, 2023). The presence vs absence and timing of ice cover influences the length of the shipping season and during high ice years shipping has been entirely suspended which interrupts supply chains that rely on raw materials such as Iron Ore, Grain, etc. The biota of the lakes also interact with ice cover which impacts nutrient loading (Nicholls, 1998), serves as a winter habitat for Diatoms (Twiss et al., 2012) and has been widely suggested to protect Lake Whitefish eggs during otherwise harsh winter conditions (Brown et al., 1993; Collingsworth et al., 2017; Lynch et al., 2015).

Previous studies have demonstrated that, consistent with trends in air and ocean temperatures, water temperatures in the Great Lakes, are also increasing both in the surface (Austin and Colman, 2007, many more!) and subsurface (Anderson et al., 2021). However, the rate and extent of warming in the Great Lakes varies considerably from lake to lake and even *within* each lake (Mason et al., 2016), partly due to the heterogeneity of characteristics within lakes (bathymetry, latitude, surface area) often referred to as lake morphometry. These characteristics can act as competing factors for changes in lake ice cover. For example, Sharma et al. (2019) found that mean depth influenced ice loss for lakes at similar latitudes. Richardson et al. (2024) hypothesized that, globally, lake ice phenology will respond nonlinearly to climate-change such that lakes in geographically colder (warmer) regions experience increased (decreased) interannual variability. This is mostly consistent with the findings of Lin et al. (2022), who showed that regionally the Great Lakes have been experiencing increased variability since 1998.

Assel (2005) classified ice seasons as either mild, typical or severe and found that. . .

## 2 Data & Methods

Ice cover in the Great Lakes has been recorded for over five decades with information from high resolution, satellite-based instruments.

While much previous work has identified the significant influence of teleconnections on (Great) Lakes ice cover (Bai et al., 2012; Richardson et al., 2024; Wang et al., 2018), our focus is on subregional ice formation (i.e. differences

### 3 Results

### References

- Anderson, E. J., Stow, C. A., Gronewold, A. D., Mason, L. A., McCormick, M. J., Qian, S. S., Ruberg, S. A., Beadle, K., Constant, S. A., & Hawley, N. (2021). Seasonal overturn and stratification changes drive deep-water warming in one of earth’s largest lakes. *Nature Communications*, 12(1). <https://doi.org/10.1038/s41467-021-21971-1>
- Assel, R. A. (2005). Classification of annual great lakes ice cycles: Winters of 1973–2002\*. *Journal of Climate*, 18(22), 4895–4905. <https://doi.org/10.1175/jcli3571.1>
- Austin, J. A., & Colman, S. M. (2007). Lake superior summer water temperatures are increasing more rapidly than regional air temperatures: A positive ice-albedo feedback. *Geophysical Research Letters*, 34(6). <https://doi.org/10.1029/2006gl029021>
- Bai, X., Wang, J., Sellinger, C., Clites, A., & Assel, R. (2012). Interannual variability of great lakes ice cover and its relationship to nao and enso. *Journal of Geophysical Research: Oceans*, 117(C3). <https://doi.org/10.1029/2010jc006932>
- Brown, R. W., Taylor, W. W., & Assel, R. A. (1993). Factors affecting the recruitment of lake whitefish in two areas of northern lake michigan. *Journal of Great Lakes Research*, 19(2), 418–428. [https://doi.org/10.1016/s0380-1330\(93\)71229-0](https://doi.org/10.1016/s0380-1330(93)71229-0)
- Collingsworth, P. D., Bunnell, D. B., Murray, M. W., Kao, Y.-C., Feiner, Z. S., Claramunt, R. M., Lofgren, B. M., Höök, T. O., & Ludsins, S. A. (2017). Climate change as a long-term stressor for the fisheries of the laurentian great lakes of north america. *Reviews in Fish Biology and Fisheries*, 27(2), 363–391. <https://doi.org/10.1007/s11160-017-9480-3>
- Lin, Y.-C., Fujisaki-Manome, A., & Wang, J. (2022). Recently amplified interannual variability of the great lakes ice cover in response to changing teleconnections. *Journal of Climate*, 35(19), 6283–6300. <https://doi.org/10.1175/jcli-d-21-0448.1>
- Lynch, A. J., Taylor, W. W., Beard, T. D., & Lofgren, B. M. (2015). Climate change projections for lake whitefish ( *coregonus clupeaformis* ) recruitment in the 1836 treaty waters of the upper great lakes. *Journal of Great Lakes Research*, 41(2), 415–422. <https://doi.org/10.1016/j.jglr.2015.03.015>
- Martin. (2023). Economic impacts of maritime shipping in the great lakes - st lawrence region. <http://marinedelivers.com>
- Mason, L. A., Riseng, C. M., Gronewold, A. D., Rutherford, E. S., Wang, J., Clites, A., Smith, S. D. P., & McIntyre, P. B. (2016). Fine-scale spatial variation in ice cover and surface temperature trends across the surface of the laurentian great lakes. *Climatic Change*, 138(1–2), 71–83. <https://doi.org/10.1007/s10584-016-1721-2>
- Nicholls, K. H. (1998). El niño, ice cover, and great lakes phosphorus: Implications for climate warming. *Limnology and Oceanography*, 43(4), 715–719. <https://doi.org/10.4319/lo.1998.43.4.0715>
- Richardson, D. C., Filazzola, A., Woolway, R. I., Imrit, M. A., Bouffard, D., Weyhenmeyer, G. A., Magnuson, J., & Sharma, S. (2024). Nonlinear responses in interannual variability of lake ice to climate change. *Limnology and Oceanography*, 69(4), 789–801. <https://doi.org/10.1002/lno.12527>
- Sharma, S., Blagrove, K., Magnuson, J. J., O’Reilly, C. M., Oliver, S., Batt, R. D., Magee, M. R., Straile, D., Weyhenmeyer, G. A., Winslow, L., & Woolway, R. I. (2019). Widespread loss of lake ice around the northern hemisphere in a warming world. *Nature Climate Change*, 9(3), 227–231. <https://doi.org/10.1038/s41558-018-0393-5>
- Twiss, M., McKay, R., Bourbonniere, R., Bullerjahn, G., Carrick, H., Smith, R., Winter, J., D’souza, N., Furey, P., Lashaway, A., Saxton, M., & Wilhelm, S. (2012). Diatoms abound in ice-covered lake erie: An investigation of offshore winter limnology in lake erie over the period 2007 to 2010. *Journal of Great Lakes Research*, 38(1), 18–30. <https://doi.org/10.1016/j.jglr.2011.12.008>

Wang, J., Kessler, J., Bai, X., Clites, A., Lofgren, B., Assuncao, A., Bratton, J., Chu, P., & Leshkevich, G. (2018). Decadal variability of great lakes ice cover in response to amo and pdo, 1963–2017. *Journal of Climate*, 31(18), 7249–7268. <https://doi.org/10.1175/jcli-d-17-0283.1>