

FIN 5330 - Homework 2

Tyler J. Brough

February 19, 2019

Problem 1

1. Simulate $T = 500$ observations from an AR(1) process for $\phi = \{0.25, 0.5, 0.75, 0.8, 0.9\}$.

$$y_t = \phi y_{t-1} + \epsilon_t$$

2. Treat the artificial from the simulations above as observed data by an econometrician.
3. Estimate each model via OLS.
4. Test the standard null hypothesis of $\phi = 0$ with a standard t -test for significance levels $\{0.01, 0.05, 0.10\}$ for each one and report the results in a table. Provide test statistics, standard errors, critical values, p-values, etc.
5. Pick one of the parameter values for the models above and do the following:
 - Use the Central Limit Theorem to derive a sampling distribution for $\hat{\phi}$. Present the parameter values of the sampling distribution. Produce a graph of the distribution.
 - Use parametric Monte Carlo to simulate the sampling distribution. Use $M = 10,000$ repetitions. Use the sample mean and standard deviation to estimate the parameter values of the distribution. Produce a histogram.
 - Use the IID Bootstrap to simulate the sampling distribution. Use $B = 10,000$ repetitions. Use the sample mean and standard deviation to estimate the parameter values of the distribution. Produce a histogram.
 - Compare all three methods.
 - Can you interpret the last two distributions as predictive densities?
6. Return to the problem in 5 above and redo the simulation from step one, but replace the error distribution with a Student-T distribution with $df = 5$ (degrees of freedom parameter). Even though we know at the generation stage that the errors come from the Student-T distribution, the econometrician assumes a normal distribution when using the CLT and parametric Monte Carlo. The bootstrap obviously does not need to make such assumptions. Compare to the results above.

Problem 2

- Simulate $T = 500$ time steps for the following two equations:

$$y_t = y_{t-1} + u_{1,t}$$

$$x_t = x_{t-1} + u_{2,t}$$

- where $u_{j,t}$ $j = 1, 2$ are independent standard white noise processes.
- Next regress y_t on x_t and estimate β (slope coefficient) via OLS in the following regression

$$y_t = \alpha + \beta x_t + \epsilon_t$$

- Test the null hypothesis $H_0 : \beta = 0$ against the alternative $H_a : \beta \neq 0$. Use the standard t -test with standard significance levels (0.01, 0.05, and 0.10). What should you find? What do you find?
- Repeat the process $M = 50,000$ times and store the β coefficients for each run of the simulation.

- Summarize the simulated sampling distribution for β .
- Make a histogram plot of the simulated coefficients.

Problem 3

- Repeat the exercise in Problem 1 above for $\phi = 1.0$.
- Comment on your findings.

Problem 4

- Simulate $T = 500$ time steps from the random walk model

$$x_t = x_{t-1} + u_{1,t}$$

- Next simulate $T = 500$ time steps from the model

$$y_t = \alpha + \beta x_t + \epsilon_t$$

- Where $\alpha = 0.22$ and $\beta = 2.50$.
- $\epsilon \sim N(0, 1)$ (white noise process)
- Use the Augmented Dickey-Fuller Test to check for the presence of a unit-root in both y_t and x_t . What do you find? What should you find?
- Implement the Engle-Granger two-step method by:
 - First, test for cointegration by submitting $\hat{\epsilon}_t$ to the ADF Test. What do you find?
 - Obtain $\hat{\beta}$ via OLS.
 - Estimate the error-correction model with $p = 1$ and include contemporaneous x_t .