



The Time Domain Spectroscopic Survey: Stellar Variables

Benjamin Roulston (BU/SAO/CfA)
SDSS-IV Meeting, 2019
Ensenada, Mexico

Paul Green (P-I, CfA), Scott Anderson (P-I, UWa), Michael Eracleous (PSU), Eric Morganson (UIUC), John Ruan (UWa), Jessie Runnoe (UMich), Niel Brandt (PSU), Don Schneider (PSU), Yue Shen (UIUC), the TDSS Team, the SDSS-IV Collaboration, and the Pan-STARRS1 Science Consortium



SDSS-IV TIME DOMAIN SPECTROSCOPIC SURVEY




- Adds dedicated spectra to era of time domain imaging.
- First large-scale, *inclusive* spectroscopic survey of variable sources
 - No color selection
 - Generic variability, i.e., no requirement for periodicity, flares, etc.
- Extending Survey Science to the Time Domain/Spectroscopy through:
 - ***Systematic discovery*** of celestial variables
 - ***Population studies*** of variable stars and quasars
 - ***Exploring the unknown:*** rare objects

Three principal components of TDSS, piggy-back on




- TDSS main (90%) survey: **SES**=single-epoch spectroscopy BOSS classification/characterization spectra of imaging variables from Pan-STARRS I, 3π imaging survey (PSI); $\sim 10^5$ fibers in SDSS-IV ($10/\text{deg}^2$)
- TDSS (10%) **FES**=few-epoch spectroscopy of potential spectral variables of interest (e.g., known from SDSS I-IV spectra) $\sim 10^{3-4}$ fibers in SDSS-IV
- TDSS **RQS**=repeat quasar spectroscopy of SDSS I-IV quasars; $\sim 10^4$ fibers in SDSS-IV



10% of TDSS fibers for special FES programs:



Target Type	Number	Observed (March '19)
Total (~1 deg⁻²)	9143	6261
WD/dM Binaries	1036	214
Dwarf Carbon Stars	830	370
Ultracool Dwarfs	402	556
Hypervariable Stars ($\geq 0.3m$)	1150	727
BAL Quasars	2900	1926
High S/N Quasars	1100	593
Double Peaked Emitters	900	576
QSO MGII Line Shifts	70	62
Hypervariable Quasars	1555	1237


(Target Selection Details in MacLeod et al. 2018)



TDSS/SES update (through DR14)


STAR subclasses (pipe) range from O/B, through A/F (green), to K/M (orange/red). Flaring M stars dominate, but thousands are in A/F-star regime that includes pulsating RR Lyrae. >710 have cataloged periodic light curves, e.g., also includes eclipsers. Large symbols highlight ~60 each CVs (purple x's) & variable WDs (blue +'s).









TDSS Stellar Science Opportunities


- Expect ~25,000 stellar variables in TDSS (pulsating, eclipsing, accreting, active)
 - ~ 4,000 off the main sequence
 - ~ 1,000 RR Lyr
 - ~ Handful of
 - WDs
 - CVs
 - Carbon stars
- ~10% of variables are periodic ($\log\text{Prob} < -10$)




TDSS Stellar Exotica: LARP?




Low accretion rate polar? Highly variable. GALEX detected.



TDSS Stellar Exotica: DBZ WDs



6.5m MMT followup spectrum. Analysis pending via EC Dufour.




Expected Periodic Stellar Variable Types

to $r < 18.5$ (*Drake et al. 2014, ApJ, 213, 1*)


Type	Percent	Description
Eclipsing:		
EW	50.5	contact eclipsing binary
EA	7.6	detached (Algol) eclipsing binary
beta Lyrae	0.5	semi-detached eclipsing binary
RR Lyr:		
RRab	27.3	RR Lyrae (fundamental mode pulsators)
RRc	8.9	RR Lyrae (multi-period pulsators)
RRd	0.8	RR Lyrae (both)
Blazkho	0.4	RR Lyrae (quasi-periodic)
CEPHEID:		
ACEP	0.1	anomalous Cepheid
Cep-II	0.2	type II Cepheid
HADS	0.4	high amplitude delta Scuti
MISC:		
RSCVn	2.5	spotted rotator
LPV	0.8	Mira or semi-regular AGB variables

CSS ID: | P = 0.163
 logProb=-89.964 | Amp= 0.616
 ngood=530 | nreject=2
 nabove=42 (7%) | nbelow=93 (17%)


Drake: P=0.325056 | Amp=0.8 | VarType=EA




$\log_{10}(P / \text{day}) = -0.79$
 $\log_{10}(\text{Amp} / \text{mag}) = -0.21$
 Skewness = 1.31



$M_i = 6.03$
 $g-i = 1.31$
 UpperLim Dist = 23185 pc
 LowerLim $M_i = 0.3$




RA: , DEC: | cz = $60.92 \pm 4.32 \text{ km s}^{-1}$ | SDSS Subclass = K3V
 PyHammer = K4, RV = 100.26 km s^{-1}
 prop. | Plate = MJD = Fiberid = | GaiaDR2 Dist = 1659 pc (SNR = 3.64) | GaiaDR2 PMtot = 5.38 mas/yr (SNR = 26.81)




CSS ID: | P = 0.139
 logProb=-131.996 | Amp= 0.610
 ngood=466 | nreject=4
 nabove=75 (15%) | nbelow=96 (20%)


Drake: P=0.278904 | Amp=0.74 | VarType= β \Lyrae



$\log_{10}(P / \text{day}) = -0.86$
 $\log_{10}(\text{Amp} / \text{mag}) = -0.21$
 Skewness = 0.86



$M_i = 5.31$
 $g-i = 1.02$
 UpperLim Dist = 22511 pc
 LowerLim Mi = 0.28



RA:


DEC:

$cz = 0.0 \pm 0.0 \text{ km s}^{-1}$ | SDSS Subclass = None

DyHammer = K3, RV = -51.87 km s^{-1}


| GaiaDR2 Dist = 2229 pc (SNR=4.95) | GaiaDR2 PMtot = 5.6 mas/yr (SNR = 35.2)

DR | Plate = MJD = Fiberid =





CSS ID: P= 0.105
 logProb=-148.751 | Amp= 0.627
 ngood=433 | nreject=3
 nabove=23 (5%) | nbelow=69 (15%)

Drake: P=0.210378 | Amp=0.53 | VarType=EW



$\log_{10}(P / \text{day}) = -0.98$
 $\log_{10}(\text{Amp} / \text{mag}) = -0.2$
 Skewness = 0.77

$M_i = 6.3$
 $g-i = 1.19$
 UpperLim Dist = 21712 pc
 LowerLim $M_i = 0.61$



RA:

DEC: -

DR | Plate =


MJD = ?


Fiberid =

$z = 14.73 \pm 6.52 \text{ km s}^{-1}$ | SDSS Subclass = K5Ve

PyHammer = K5, RV = -35.23 km s⁻¹


| GaiaDR2 Dist = 1583 pc (SNR=1.7) | GaiaDR2 PMtot = 5.82 mas/yr (SNR = 23.08)






TDSS Stellar Science Opportunities

- 1000 RR Lyr w/RVs and metallicity estimates, could help identify streams
- Flare stars
- Active WD+dM systems
 - Follow FES dC paper and get separations
 - new post-CE?
 - Statistically examine the activity variation of close pairs as a function of spectral type, measure the white dwarf cooling age, and (possibly the) metallicity of the system
- **identify SB2s within the sample**





PyHammer & Extension

- PyHammer ([Kesseli et al. \(2017\)](#), <https://github.com/BU-hammerTeam/PyHammer>)
- Add additional C and WD templates (from TDSS)
- Developing tool to identify SB2s within the sample
 - Stellar templates from: A (Pickles+ 1998)
FGKM (MaStar, Yan+ 2018)
WD (Levenhagen+ 2017)





TDSS PyHammer WDs






TDSS PyHammer Carbon Stars






Composite-spectrum Binaries (a.k.a. SB2)

Can sometimes be obvious!







More Typical Composite-Spectrum Binaries


Survey: *eboss* Program: *eboss Target*:

$cz = -88 \pm 3$ km/s Class=STAR K5Ve (118100)
No warnings.





PyHammer SB2 Extension




TDSS WIKI

<https://trac.sdss.org/wiki/TDSS>

- Telecon number/minutes
- Documents and brief primer
- Instructions/software info for VIP (Ruan et al.)
- spAllTDSS file (an spAll file for TDSS targets, with SDSS-IV spectro pipeline parameters): **spAllTDSS-v5_10_10-01Jul2018_v0.txt**
- PS1/SDSS-IV data-sharing file

TEAM



S. Anderson (UWa), C. Badenes (UPitt), M. Bershadsky (UWi), M. Blanton (NYU), N. Brandt (PSU), K. Chambers (IfA), J. Davenport (UW), K. Dawson (Utah), M. Eracleous (PSU), N. Filiz Ak (Erciyes), P. Garnavich (Notre Dame), A. Georgakakis (MPE), P. Green (CfA), J. Greene (Princeton), C. Grier (PSU), P. Hall (York), S. Hawley (UW), N. Kaiser (IfA), J.-P. Kneib (EPFL), C. MacLeod (CfA), Vivek Mariappan (PSU), I. McGreer (UA), A. Merloni (MPE), E. Morganson (UIUC), A. Myers (UWy), I. Paris (IAP), B. Roulston (SAO/BU), J. Ruan (McGill), J. Runnoe (UMich), M. Salvato (MPE), E. Schlaufly (MPIA), D. Schlegel (LBNL), S. Schmidt (AIP), D. Schneider (PSU), Axel Schwone (AIP), B. Sesar (Caltech), T. Shanks (Durham), S. Smartt (Queen's U. Belfast), K. Stassun (Vander.), P. Szkody (UWa), J. Tonry (IfA), F. Walter (MPIA), D. York (UC), ... YOUR NAME HERE