

Square lattice Ising model

文 Add languages

Tools Article Talk

From Wikipedia, the free encyclopedia



improve it to make it understandable to non-experts, without removing the technical details. (May 2019) (Learn how and when to remove this template message) In statistical mechanics, the two-dimensional square lattice Ising model is a simple lattice model of

This article may be too technical for most readers to understand. Please help

interacting magnetic spins. The model is notable for having nontrivial interactions, yet having an analytical solution. The model was solved by Lars Onsager for the special case that the external magnetic field $H=0.^{[1]}$ An analytical solution for the general case for $H\neq 0$ has yet to be found. Defining the partition function [edit]

the horizontal and vertical directions, which effectively reduces the topology of the model to a torus.

Generally, the horizontal coupling J and the vertical coupling J^* are not equal. With $eta=rac{1}{kT}$ and absolute temperature T and Boltzmann's constant k, the partition function

Consider a 2D Ising model on a square lattice Λ with N sites and periodic boundary conditions in both

 $Z_N(K\equiv eta J, L\equiv eta J^*) = \sum_{\{\sigma\}} \exp\left(K\sum_{\langle ij
angle_H} \sigma_i \sigma_j + L\sum_{\langle ij
angle_V} \sigma_i \sigma_j
ight).$

Critical temperature `[edit]`
The critical temperature
$$T_c$$
 can be obtained from the Kramers–Wannier duality relation. Denoting the free energy per site as $F(K,L)$, one has:

$eta F\left(K^{st},L^{st} ight) = eta F\left(K,L ight) + rac{1}{2}\log\left[\sinh(2K)\sinh(2L) ight]$

 $\sinh(2K^*)\sinh(2L)=1$

where

$$\sinh(2L^*)\sinh(2K)=1$$
 Assuming there is only one critical line in the (K,L) plane, the duality relation implies that this is given

by:

$$\sinh(2K)\sinh(2L)=1$$
 For the isotropic case $J=J^*$, one finds the famous relation for the critical temperature T_c

Dual lattice [edit]

 $e^{K(N-2s)+L(N-2r)}$


Construct a dual lattice Λ_D as

 $\frac{kT_c}{J} = \frac{2}{\ln(1+\sqrt{2})} \approx 2.26918531421$

Consider a configuration of spins
$$\{\sigma\}$$
 on the square lattice Λ . Let r and s denote the number of unlike neighbours in the vertical and horizontal directions respectively. Then the summand in Z_N corresponding to $\{\sigma\}$ is given by

depicted in the diagram. For every configuration $\{\sigma\}$, a polygon is associated to the lattice by drawing a line on the edge of the dual lattice if the

spins separated by the edge are unlike. Since by traversing a vertex of Λ the spins need to change an even number of times so that one arrives at the starting point with the same charge, every vertex of the dual lattice is connected to an even number of lines in the configuration, defining a polygon. This reduces the partition function to $Z_N(K,L) = 2e^{N(K+L)} \sum_{i} e^{-2Lr - 2Ks}$ summing over all polygons in the dual lattice, where r and s are factor of 2 arising from the inversion of spin configuration.



 $Z_N(K,L) = 2e^{N(K+L)} \; \sum \; e^{-2Lr-2Ks}$

Since $\sigma\sigma'=\pm 1$ one has

defines a low temperature expansion of $Z_N(K,L)$.

High-temperature expansion [edit]

 $e^{K\sigma\sigma'}=\cosh K+\sinh K(\sigma\sigma')=\cosh K(1+\tanh K(\sigma\sigma')).$ **Therefore**

where v= anh K and w= anh L . Since there are N horizontal and vertical edges, there are a

where the sum is over all polygons in the lattice. Since anh K, anh L o 0 as $T o \infty$, this gives the

 $Z_N(K,L) = (\cosh K \cosh L)^N \sum_{\{\sigma\}} \prod_{\langle ii
angle_{TT}} (1 + v \sigma_i \sigma_j) \prod_{\langle ii
angle_{TT}} (1 + w \sigma_i \sigma_j)$

At low temperatures, K, L approach infinity, so that as $T \to 0, \ e^{-K}, e^{-L} \to 0$, so that

total of 2^{2N} terms in the expansion. Every term corresponds to a configuration of lines of the lattice, by associating a line connecting i and j if the term $v\sigma_i\sigma_j$ (or $w\sigma_i\sigma_j$) is chosen in the product. Summing

over the configurations, using $\sum_{\sigma:=\pm 1} \sigma_i^n = egin{cases} 0 & ext{for } n ext{ odd} \ 2 & ext{for } n ext{ even} \end{cases}$

shows that only configurations with an even number of lines at each vertex (polygons) will contribute to the partition function, giving $Z_N(K,L) = 2^N (\cosh K \cosh L)^N \sum_{D \subset \Lambda} v^r w^s$

The Helmholtz free energy per site F can be expressed as

and the spontaneous magnetization is, for $T < T_c$,

 $M = \left\lceil 1 - \sinh^{-4}(2 eta J)
ight
ceil^{1/8}$

and M=0 for $T\geq T_c$.

high temperature expansion of $Z_N(K,L)$.

 $k = rac{1}{\sinh(2K)\sinh(2L)}$

The free energy per site in the limit $N o \infty$ is given as follows. Define the parameter k as

 $-eta F = rac{\log(2)}{2} + rac{1}{2\pi} \int_0^\pi \log \left[\cosh(2K) \cosh(2L) + rac{1}{k} \sqrt{1 + k^2 - 2k \cos(2 heta)}
ight] d heta$ For the isotropic case $J=J^st$, from the above expression one finds for the internal energy per site:

 $U = -J \coth(2eta J) \left| 1 + rac{2}{\pi} (2 anh^2 (2eta J) - 1) \int_0^{\pi/2} rac{1}{\sqrt{1 - 4k(1+k)^{-2} \sin^2(heta)}} d heta
ight|$

Notes [edit]

doi:10.1103/revmodphys.39.883 2. ISSN 0034-6861 2.

8121/ac0983 2. ISSN 1751-8113 2. S2CID 232290629 2.

MR 0690578 2 • Baxter, Rodney J. (2016). "The bulk, surface and corner free energies of the square lattice Ising model". Journal of Physics A: Mathematical and Theoretical. 50 (1). IOP Publishing: 014001. arXiv:1606.02029 doi:10.1088/1751-8113/50/1/014001 2 ISSN 1751-8113 2 S2CID 2467419 2.

• Huang, Kerson (1987), Statistical mechanics (2nd edition), Wiley, ISBN 978-0471815181

375201. arXiv:2103.10776 3. Bibcode:2021JPhA...54K5201H 2. doi:10.1088/1751-

• Hucht, Alfred (2021). "The square lattice Ising model on the rectangle III: Hankel and Toeplitz

determinants". Journal of Physics A: Mathematical and Theoretical. 54 (37). IOP Publishing:

• Ising, Ernst (1925), "Beitrag zur Theorie des Ferromagnetismus", Z. Phys., 31 (1): 253–258,

• Barry M. McCoy and Tai Tsun Wu (1973), The Two-Dimensional Ising Model. Harvard University Press, Cambridge Massachusetts, ISBN 0-674-91440-6 • Montroll, Elliott W.; Potts, Renfrey B.; Ward, John C. (1963), "Correlations and spontaneous

Cookie statement Mobile view

doi:10.1103/PhysRev.65.117 2, MR 0010315 2

0521408059

• Onsager, Lars (1944), "Crystal statistics. I. A two-dimensional model with an order-disorder transition", *Phys. Rev.*, Series II, **65** (3–4): 117–149, Bibcode:1944PhRv...65..1170 2,

• Yang, C. N. (1952), "The spontaneous magnetization of a two-dimensional Ising model", *Physical* Review, Series II, 85 (5): 808–816, Bibcode:1952PhRv...85..808Y ₺, doi:10.1103/PhysRev.85.808 2, MR 0051740 2 Categories: Statistical mechanics | Exactly solvable models | Lattice models

• Onsager, Lars (1949), "Discussion", Supplemento al Nuovo Cimento, 6: 261

Transition" ☑. Physical Review. 65 (3–4): 117–149. doi:10.1103/PhysRev.65.117 ☑. References [edit] • Baxter, Rodney J. (1982), Exactly solved models in statistical mechanics (PDF), London: Academic Press Inc. [Harcourt Brace Jovanovich Publishers], ISBN 978-0-12-083180-7,

1. ^ Onsager, Lars (1944-02-01). "Crystal Statistics. I. A Two-Dimensional Model with an Order-Disorder

• Kurt Binder (2001) [1994], "Ising model" , Encyclopedia of Mathematics, EMS Press • BRUSH, STEPHEN G. (1967-10-01). "History of the Lenz-Ising Model". Reviews of Modern Physics. 39 (4). American Physical Society (APS): 883–893. Bibcode:1967RvMP...39..883B 2.

- Bibcode:1925ZPhy...31..25312, doi:10.1007/BF029805772, S2CID 1221573192 • Itzykson, Claude; Drouffe, Jean-Michel (1989), Théorie statistique des champs, Volume 1, Savoirs actuels (CNRS), EDP Sciences Editions, ISBN 978-2868833600 • Itzykson, Claude; Drouffe, Jean-Michel (1989), Statistical field theory, Volume 1: From Brownian motion to renormalization and lattice gauge theory, Cambridge University Press, ISBN 978-
- magnetization of the two-dimensional Ising model" , Journal of Mathematical Physics, 4 (2): 308-322, Bibcode:1963JMP.....4..308M 2, doi:10.1063/1.1703955 2, ISSN 0022-2488 2, MR 0148406 ☑, archived from the original ☑ on 2013-01-12

• John Palmer (2007), Planar Ising Correlations. Birkhäuser, Boston, ISBN 978-0-8176-4248-8.

This page was last edited on 23 February 2024, at 02:45 (UTC). Text is available under the Creative Commons Attribution-ShareAlike License 4.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.

Privacy policy About Wikipedia Disclaimers Contact Wikipedia Code of Conduct Developers Statistics