
a MEG-MASC (NEMAR nm000229), linear probe on frozen MEG-XL features; primary cell 7 % train data; one probe seed per arm
chance
paper ±5 % (pre-registered band)
value
kept epoch / epochs run
Paper (Fig. 4, MEG-MASC, 150 s)
0.2197 ± 0.0096
Authors' code, attention dropout zeroed
0.2130
1 / 11
braindecode MEGXL, authors' sensor geometry
0.2321
5 / 15
braindecode MEGXL, each recording's chs_info
0.2249
5 / 15
Secondary / controls (not in any verdict)
Authors' code as released (dropout replay, 12 draws)
0.2263
6 / 16
Authors' code, 25 % train (config default)
0.2330
2 / 12
braindecode MEGXL, 25 % train
0.2449
3 / 13
0.20
0.21
0.22
0.23
0.24
balanced top-10 accuracy, 50-word retrieval (test, 882 segments, 27 subjects)
b, c: post-hoc reviewer diagnostic (not part of the pre-registered verdict). Dashed lines: chance.
b Test value per epoch
test balanced top-10 acc. (50 words)
epoch
c Early-stopping criterion (validation)
val balanced top-10 acc. (250 words)
epoch-1 value
+ min_delta 0.001
epoch
Authors' code, attention dropout zeroed
braindecode MEGXL, authors' sensor geometry
braindecode MEGXL, each recording's chs_info
kept checkpoint (early stopping)
authors' code, repeat run (same command, seed 42; log, 4 dp)