

Maintenance Scheduling System

Christian Brunbjerg Jespersen

Technical University of Denmark

January 1, 1980

Agenda

- ▶ Introduction to Maintenance Scheduling
- ▶ Architecture of a Scheduling System
- ▶ Possible Contributions to Operation Research

Mathematical Notation: Sets

$a \in$

$A_{b,c}^m(t, x, y)$

- ▶ a: set element
- ▶ A: set itself
- ▶ b: set element from set B
- ▶ c: set element from set C
- ▶ m: model formulation m
- ▶ t: time
- ▶ x: value of decision variable from a different model
- ▶ y: value of decision variable from a different model

Mathematical Notation: Parameters

*name_of_parameter*_{a,b}(*t*, *x*, *y*)

- ▶ parameters are functions of set elements and input parameters
- ▶ a: set element from A
- ▶ b: set element from B
- ▶ t: time
- ▶ x: value of decision variable from another model
- ▶ y: value of decision variable from another model

Mathematical Notation: Variables

$$x_{a,b}^m(t)$$

- ▶ variables are functions of set elements, specified model, and time
- ▶ x: decision variable
- ▶ a: set element from A
- ▶ b: set element from B
- ▶ m: specifying the model
- ▶ t: time
- ▶ Notice: decision variables cannot depend on other decision variables as it would make them belong to the same model.

Strategic

Meta variables:

$$s \in S \quad (1)$$

$$\beta(\tau) \quad (2)$$

$$\tau \in [0, \infty] \quad (3)$$

Maximize:

$$\begin{aligned} & \sum_{w \in W(\tau)} \sum_{p \in P(\tau)} \text{strategic_value}_{wp}(\tau) \cdot \alpha_{wp}(\tau) \\ & - \sum_{p \in P(\tau)} \sum_{r \in R(\tau)} \text{strategic_penalty} \cdot \epsilon_{pr}(\tau) \\ & + \sum_{p \in P(\tau)} \sum_{w1 \in W(\tau)} \sum_{w2 \in W(\tau)} \text{clustering_value}_{w1, w2} \cdot \alpha_{w1p}(\tau) \cdot \alpha_{w2p}(\tau) \end{aligned} \quad (4)$$

Subject to:

$$\sum_{w \in W(\tau)} \text{work_order_work}_{wr} \cdot \alpha_{wp}(\tau) \leq \text{resource}_{pr}(\tau, \beta(\tau)) + \epsilon_{pr}(\tau) \quad \forall p \in P(\tau) \quad \forall r \in R(\tau) \quad (5)$$

$$\sum_{w \in W(\tau)} \alpha_{wp}(\tau) = 1 \quad \forall p \in P(\tau) \quad (6)$$

$$\alpha_{wp}(\tau) = 0 \quad \forall (w, p) \in \text{exclude}(\tau) \quad (7)$$

$$\alpha_{wp}(\tau) = 1 \quad \forall (w, p) \in \text{include}(\tau) \quad (8)$$

$$\alpha_{wp}(\tau) \in \{0, 1\} \quad \forall w \in W(\tau) \quad \forall p \in P(\tau) \quad (9)$$

$$\epsilon_{pr}(\tau) \in \mathbb{R}^+ \quad \forall p \in P(\tau) \quad \forall r \in R(\tau) \quad (10)$$

Tactical

Meta variables:

$$s \in S \quad (11)$$

$$\alpha(\tau) \quad (12)$$

$$\tau \in [0, \infty] \quad (13)$$

Minimize:

$$\sum_{o \in O(\tau, \alpha(\tau))} \sum_{d \in D(\tau)} \text{tactical_value}_{do}(\tau) \cdot \beta_{do}(\tau) + \sum_{r \in R(\tau)} \sum_{d \in D(\tau)} \text{tactical_penalty} \cdot \mu_{rd}(\tau) \quad (14)$$

Subject to:

$$\sum_{o \in O(\tau, \alpha(\tau))} \text{work}_o(\tau) \cdot \beta_{do}(\tau) \leq \text{tactical_resource}_{dr}(\tau) + \mu_{rd}(\tau) \forall d \in D(\tau) \quad \forall r \in R(\tau) \quad (15)$$

$$\sum_{d=\text{earliest_start}_o(\tau)}^{\text{latest_finish}_o(\tau)} \sigma_{do}(\tau) = \text{duration}_o(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad (16)$$

$$\sum_{d^* \in D_{\text{duration}_o(\tau)}(\tau)} \sigma_{d^*o}(\tau) = \text{duration}_o(\tau) \cdot \eta_{do}(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad \forall d \in D(\tau) \quad (17)$$

$$\sum_{o \in O(\tau, \alpha(\tau))} \eta_{do}(\tau) = 1, \quad \forall d \in D(\tau) \quad (18)$$

$$\sum_{d \in D(\tau)} d \cdot \sigma_{do1}(\tau) + \Delta_o(\tau) = \sum_{d \in D(\tau)} d \cdot \sigma_{do2}(\tau) \quad \forall (o1, o2) \in \text{finish_start}_{o1, o2} \quad (19)$$

$$\sum_{d \in D(\tau)} d \cdot \sigma_{do1}(\tau) = \sum_{d \in D(\tau)} d \cdot \sigma_{do2}(\tau) \quad \forall (o1, o2) \in \text{start_start}_{o1, o2} \quad (20)$$

$$\beta_{do}(\tau) \leq \text{number}_o(\tau) \cdot \text{operating_time}_o \quad \forall d \in D(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad (21)$$

$$\beta_{do}(\tau) \in \mathbb{R} \quad \forall d \in D(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad (22)$$

$$\mu_{rd}(\tau) \in \mathbb{R} \quad \forall r \in R(\tau) \quad \forall d \in D(\tau) \quad (23)$$

$$\sigma_{do}(\tau) \in \{0, 1\} \quad \forall d \in D(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad (24)$$

$$\eta_{do}(\tau) \in \{0, 1\} \quad \forall d \in D(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad (25)$$

$$\Delta_o(\tau) \in \{0, 1\} \quad \forall o \in O(\tau, \alpha(\tau)) \quad (26)$$

Supervisor

Meta variables:

$$z \in Z \quad (26)$$

$$\alpha(\tau) \quad (27)$$

$$\theta(\tau) \quad (28)$$

$$\tau \in [0, \infty] \quad (29)$$

Maximize:

$$\sum_{a \in A(\tau, \alpha(\tau))} \sum_{t \in T(\tau)} \text{supervisor_value}_{at}(\tau, \lambda_t(\tau), \Lambda_t(\tau)) \cdot \gamma_{at}(\tau) \quad (30)$$

Subject to:

$$\sum_{a \in A_o(\tau, \alpha(\tau))} \rho_a(\tau) = \text{work}_o(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad (31)$$

$$\sum_{t \in T(\tau)} \sum_{a \in A_o(\tau, \alpha(\tau))} \gamma_{at}(\tau) = \phi_o(\tau) \cdot \text{number}_o(\tau) \quad \forall o \in O(\tau, \alpha(\tau)) \quad (32)$$

$$\sum_{o \in O_w(\tau, \alpha(\tau))} \phi_o(\tau) = |Q_w(\tau, \alpha(\tau))| \quad \forall w \in W(\tau, \alpha(\tau)) \quad (33)$$

$$\sum_{a \in A_o(\tau, \alpha(\tau))} \gamma_{at}(\tau) \leq 1 \quad \forall o \in O(\tau, \alpha(\tau)) \quad \forall t \in T(\tau) \quad (34)$$

$$\gamma_{at}(\tau) \leq \text{feasible}_{at}(\theta(\tau)) \quad \forall o \in O(\tau, \alpha(\tau)) \quad \forall t \in T(\tau) \quad (35)$$

$$\gamma_{at}(\tau) \in \{0, 1\} \quad \forall o \in O(\tau, \alpha(\tau)) \quad \forall t \in T(\tau) \quad (36)$$

$$\rho_a(\tau) \in [\text{lower_activity_work}_a(\tau), \text{work}_a(\tau)] \quad \forall a \in A(\tau, \alpha(\tau)) \quad (37)$$

Operational

Meta variables:

$$t \in T(\tau) \quad (38)$$

$$\alpha(\tau) \quad (39)$$

$$\gamma(\tau) \quad (40)$$

$$\tau \in [0, \infty] \quad (41)$$

Maximize:

$$\sum_{a \in A(\tau, \gamma_t(\tau))} \sum_{k \in K(\gamma(\tau))} \delta_{ak}(\tau) \quad (42)$$

Subject to:

$$\sum_{k \in K(\gamma(\tau))} \delta_{ak}(\tau) \cdot \pi_{ak}(\tau) = \text{activity_work}_a(\tau, \rho(\tau)) \cdot \theta \quad (\tau) \forall a \in A(\tau, \gamma_t(\tau)) \quad (43)$$

$$\lambda_{a21}(\tau) \geq \Lambda_{a1, \text{last}(a1)}(\tau) + \text{preparation}_{a1, a2} \quad \forall a1 \in A(\tau, \gamma_t(\tau)) \quad \forall a2 \in A(\tau, \gamma_t(\tau)) \quad (44)$$

$$\lambda_{ak}(\tau) \geq \Lambda_{ak-1}(\tau) - \text{constraint_limit} \cdot (2 - \pi_{ak}(\tau) + \pi_{ak-1}(\tau)) \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad (45)$$

$$\delta_{ak}(\tau) = \Lambda_{ak}(\tau) - \lambda_{ak}(\tau) \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad (46)$$

$$\lambda_{ak}(\tau) \geq \text{event}_{ie} + \text{duration}_{ie} - \text{constraint_limit} \cdot (1 - \omega_{akie}(\tau)) \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad \forall i \in I(\tau) \quad \forall e \in E(\tau) \quad (47)$$

$$\Lambda_{ak}(\tau) \leq \text{event}_{ie} + \text{constraint_limit} \cdot \omega_{akie}(\tau) \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad \forall i \in I(\tau) \quad \forall e \in E(\tau) \quad (48)$$

$$\lambda_{a1}(\tau) \geq \text{time_window_start}_a(\beta(\tau)) \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad (49)$$

$$\Lambda_{a, \text{last}(a)}(\tau) \leq \text{time_window_finish}_a(\beta(\tau)) \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad (50)$$

$$\pi_{ak}(\tau) \in \{0, 1\} \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad (51)$$

$$\lambda_{ak}(\tau) \in [\text{availability_start}(\tau), \text{availability_finish}(\tau)] \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad (52)$$


$$\Lambda_{ak}(\tau) \in [\text{availability_start}(\tau), \text{availability_finish}(\tau)] \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad (53)$$

$$\delta_{ak}(\tau) \in [0, \text{work}_{a, \text{to_o}(a)}(\tau)] \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad (54)$$

$$\omega_{akie}(\tau) \in \{0, 1\} \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad \forall k \in K(\gamma(\tau)) \quad \forall i \in I(\tau) \quad \forall e \in E(\tau) \quad (55)$$

$$\theta_a(\tau) \in \{0, 1\} \quad \forall a \in A(\tau, \gamma_t(\tau)) \quad (56)$$

Architecture



Managing metaheuristics



Solution sharing (At




Metaheuristics (Mathe

Data storage (Mem

Message passing (M

Architecture



Managing metaheuristics



Solution sharing (At




Metaheuristics (Mathe

Data storage (Mem

Message passing (M

1. Reactive Versus Static Constraints
2. Dynamic
3. Business
4. Technical
5. Academic


Dynamic versus Static Models



- ▶ Mathematical models guide direction but does not provide direct solutions.
- ▶ Static solutions are rarely fully executable.
- ▶ Dynamic models are less constrained and ensure a contained optimal solution.
- ▶ Remember: The real optimal solution is ever knowable at time $= 0$


Meaningful
Model
Complexity

Actual
Uncertainty




Correct Model for that business process

Time horizon




- An engineering trade off will have to be made

Model setup



Quality of Solutions



Business