

Conditioning a Record on Its Own Future: Targeted Accordance and Survivorship-Conditioned Path Measures

Ajax Benander*

September 2026 — Working Draft, rev. 5

Abstract

A future condition can reshape the statistical profile of earlier events without anything in the future acting on the past: conditioning a law over histories on a future event reweights the entire path, and the Doob transform is the standard form of that reweighting. This paper states the attribute-level consequence and keeps it separate from the inference the consequence does not license. The claim in one sentence: if the realized member of a binary partition of the past is drawn under a law conditioned on a future target, then for every $\kappa > 1$ the realized member is, with probability at least $1 - 1/(1 + \kappa)$, within a κ -fold exceedance of being the more target-productive member, a statement indifferent to the target's tense; and whether a given record is drawn under such a law is a separate hypothesis, tested by comparing measures. Four things follow at their stated grades. The theorem, with its one-line proof. Its specialization to survival targets, where the critical regime forbids conditioning on a null event and the right object is the horizon family $\{T > t\}$ with the Q -process as its limit. The between-model layer: the Bayes factor between a survivorship-conditioned and an unconditioned law, which at finite information is the Doob density itself, so that a historical model comparison and a stochastic h -transform are one object; the realized log-evidence fluctuates and the monotone quantity is the relative entropy, whose rate is the information flux. And the limit on certification: the two laws are mutually absolutely continuous on every finite horizon and singular only at infinity, so no event of bounded duration separates them with probability one, while statistical discrimination remains available and is what the Bayes factor measures. A conjunction of predeclared attributes sharpens the theorem multiplicatively under submultiplicativity, and the multiplicity guard that any retrospective scan owes is stated as part of the result.

1 Introduction

Conditioning a stochastic process on a future event is standard. The process conditioned to avoid absorption indefinitely, the Q -process, is constructed as a limit of finite-horizon conditionings and realized as a Doob h -transform by the persistence-harmonic function; the penalization literature treats a broad class of such conditionings uniformly [3, 4, 5]. What is standard in probability has a consequence in the interpretation of records that is not usually drawn, and this paper

*Email: akb9408@rit.edu. The survivorship model and its Q -process limit are proved in a companion in applied probability [1] and imported here; the attribute-level apparatus descends from the Nexic companion [2]. The empirical hypothesis that the record of natural history is drawn under the survivorship-conditioned law, and its consequences, are developed in a separate downstream companion on the Route World hypothesis, not cited here so that the dependency runs one way, and the theorems here and the hypothesis there carry their different grades separately. Drafting collaboration: Anthropic Claude (Opus 4.x, Fable 5), under the author's direction.

draws it at the level of attributes. If a history is sampled under a law conditioned on a future target, then each realized attribute of that history, read against its complement, is constrained by the target in a way the unconditioned law does not impose: the realized member is typically the more target-productive of its pair, and a quantitative bound says how typically. Nothing in the bound requires the future to act on the past. The conditioning is a statement about which histories are in the support and with what weight, and the past of a weighted history carries the weight’s fingerprints.

The paper’s discipline is to keep two statements apart that are easy to run together. The first is a theorem: *under* a target-conditioned draw, realized attributes accord with the target. The second is an inference: *because* a realized attribute accords with a target, the record was drawn under conditioning on it. The second does not follow from the first. It is a comparison between rival laws for the record, and it is answered by a Bayes factor, not by the theorem. Both are stated here, in that order, and the theorem’s proof is placed beside the disclaimer of the inference so that the two cannot be confused.

The claim, in one sentence. If the realized member of a binary partition of the past is drawn under a law conditioned on a future target, then for every $\kappa > 1$ it is, with probability at least $1 - 1/(1 + \kappa)$, within a κ -fold exceedance of being the more target-productive member, a statement indifferent to the target’s tense; whether a given record is drawn under such a law is a separate hypothesis, tested by comparing measures.

What is imported. The survivorship companion [1] proves the model in which survival targets are specialized here, the finite-horizon conditioned laws and their Q -process limit in the critical regime, and the theorem that the limit is locally absolutely continuous with respect to the unconditioned law on survivors and globally singular. The Nexic companion [2] supplies the attribute-level setting: an information state as a set of stated attributes with the unstated blank, the postulate that a realized attribute is drawn by its parent-conditional weights, and the exceedance modality. Each is restated in Section 2 at its grade.

What is new. The theorem in its general form (Section 3), stated for an arbitrary target and proved in one line, with its prior-bound corollary in joint-event form and its conjunction form under the correct dependence assumption. The specialization to survival targets through the horizon family (Section 4), which removes the zero-probability conditioning that a naive statement of the survival target commits. The between-model layer (Section 5): the Bayes factor between conditioned and unconditioned laws identified with the Doob density at finite information, and the information flux as the monotone statistic the realized evidence lacks. The limit on certification (Section 6) stated at its exact strength. And the protocol under which the theorem’s application to a real record is a prospective test rather than a retrospective fit (Section 7).

Position in the branch. This paper’s core theorem (Sections 2–3) depends on nothing Golden and on no sibling; it is a statement about arbitrary target conditioning. Its survival specialization (Section 4) imports the survivorship companion [1], which is the root of the branch; a downstream technoanatomy companion imports this paper and the survivorship one for its mechanism; and a terminal Route World companion imports all three to state its hypothesis. Neither downstream paper is cited here, so that the citation graph, like the logical one, runs one way. No downstream paper is cited here as support for anything.

What is refused. No claim that any actual record is drawn under a future-conditioned law is made here; that is the downstream companion’s hypothesis, and it is carried there at a hypothesis’s grade. No retrocausal mechanism is invoked anywhere. And no result below

certifies anything about a record from finite data, which Section 6 proves impossible.

2 Setting

Definition 2.1 (Laws over histories, and partitions of the past). Let $(\Omega, \mathcal{F}, \mathbb{P})$ carry a law over histories, with a filtration $(\mathcal{F}_t)$ recording the history to time t . An *attribute* is an $\mathcal{F}_\infty$ -measurable event; a *binary partition* of the past by an attribute is a pair $\{A, A'\}$ with $A' = \Omega \setminus A$ and $0 < \mathbb{P}(A) < 1$. The *realized member* A^* is the member of the pair that obtains, and $A^{*'} its complement. A *target* is an attribute B with $\mathbb{P}(B) > 0$, and the *B-conditioned law* is $\mathbb{P}_B := \mathbb{P}(\cdot | B)$. Nothing in the definition orders A and B in time: an attribute of the history's future is an attribute.$

Definition 2.2 (Information states and the draw; from [2]). An information state fixes a set of stated attributes and treats the unstated as blank rather than as unspecified. The *draw* is the postulate that at each open attribute the realized member is distributed by the parent-conditional weights of the law in force; under $\mathbb{P}$ the weights are $\mathbb{P}(A), \mathbb{P}(A')$, and under $\mathbb{P}_B$ they are $\mathbb{P}(A | B), \mathbb{P}(A' | B)$. The modality $\lesssim$ used below is the *exceedance form*: $X \lesssim Y$ abbreviates the family of statements that $X \geq \kappa Y$ has probability at most $1/(1 + \kappa)$ for every $\kappa > 1$, the probability taken over the draw.

Definition 2.3 (The survivorship model; from [1]). A history carries a hazard process $h_t > 0$ and a killing time T with $\mathbb{P}(T > t | \mathcal{F}_\infty^h) = e^{-\Lambda_t}$, $\Lambda_t = \int_0^t h$. The finite-horizon conditioned laws are $Q_t := \mathbb{P}(\cdot | T > t)$. In the *critical regime*, $\mathbb{P}(T = \infty) = 0$ while $\mathbb{E}[T] = \infty$, and the Q -process is the limit $Q := \lim_{t \rightarrow \infty} Q_t$ on each $\mathcal{F}_s$. Everything in this paper is written for the abstract density

$$Z_s := \left. \frac{dQ}{d\mathbb{P}} \right|_{\mathcal{F}_s},$$

and two specializations are imported at their grades: the hard-wall law, proved, $Z_s^{\text{hard}} = \mathbf{1}_{\{\tau > s\}} \mathfrak{h}(X_s) / \mathfrak{h}(X_0)$ with $\mathfrak{h}$ the persistence-harmonic function; and the soft-wall law, conditional on the companion's ratio hypothesis (its Ansatz B), $Z_s^{\text{soft}} = e^{-\int_0^s q(X_u) du} g(X_s) / g(X_0)$, which coincides with the hard-wall density exactly when g is proportional to $\mathfrak{h}$ after a credit translation. No result below depends on which specialization is in force.

Theorem 2.4 (Locally ours, globally elsewhere; [P] for the hard-wall law, [C] for the soft-wall law, in [1]). *In the critical regime $Q|_{\mathcal{F}_s} \ll \mathbb{P}|_{\mathcal{F}_s}$ for every finite s , with mutual absolute continuity on the survival event, while $Q \perp \mathbb{P}$ on $\mathcal{F}_\infty$.*

3 Future-targeted accordance

Theorem 3.1 (Future-targeted accordance). *Let $\{A, A'\}$ be a binary partition of the past and B a target, and suppose the realized member A^* is drawn under $\mathbb{P}_B$. Then for every $\kappa > 1$,*

$$\mathbb{P}_B \left[\frac{\mathbb{P}(B | A^{*'})}{\mathbb{P}(B | A^*)} \geq \kappa \frac{\mathbb{P}(A^*)}{\mathbb{P}(A^{*'})} \right] \leq \frac{1}{1 + \kappa}. \quad (1)$$

Equivalently, under the B-conditioned draw, $\mathbb{P}(B | A^{'}) / \mathbb{P}(B | A^*) \lesssim \mathbb{P}(A^*) / \mathbb{P}(A^{*'})$.*

Proof. By Bayes, for either member C of the pair, $\mathbb{P}(B \mid C')/\mathbb{P}(B \mid C) = (\mathbb{P}(C' \mid B)/\mathbb{P}(C \mid B)) \cdot (\mathbb{P}(C)/\mathbb{P}(C'))$, so the event in (1) is $\{\mathbb{P}(A^* \mid B)/\mathbb{P}(A^* \mid B) \geq \kappa\}$, that the realized member carries B -conditioned weight $w := \mathbb{P}(A^* \mid B) \leq 1/(1 + \kappa)$. For $\kappa > 1$ the threshold is below one half, so at most one member of the pair has conditioned weight below it. Under $\mathbb{P}_B$ that member, if it exists, is realized with probability equal to its conditioned weight, which is at most $1/(1 + \kappa)$; if it does not exist the event is empty. $\square$

Remark 3.2 (What the theorem says, and what it does not). The outer probability in (1) is $\mathbb{P}_B$. The theorem says that a record drawn under conditioning on B has realized attributes that typically accord with B : the more B -productive member of each pair is typically the one realized, to within an exceedance the draw prices. It does not say that a record whose realized attributes accord with B was drawn under $\mathbb{P}_B$. That is a comparison between $\mathbb{P}_B$ and rival laws for the record, and it is Section 5. The theorem is a within-model statement; the inference is between-model; and the whole discipline of applying this result to an actual record is to keep them apart. Nothing in the statement or the proof distinguishes a target in the past from one in the future: B is an attribute, and the draw is the same draw.

Corollary 3.3 (The prior bound, in joint-event form). *Fix a member A of the pair with $\mathbb{P}(A) = p < \frac{1}{2}$. Under the B -conditioned draw, the joint event that A is realized and is the less B -productive member has probability at most p :*

$$\mathbb{P}_B[A^* = A \text{ and } \mathbb{P}(B \mid A') \geq \mathbb{P}(B \mid A)] \leq p. \quad (2)$$

Proof. $\mathbb{P}(B \mid A') \geq \mathbb{P}(B \mid A)$ holds exactly when $w := \mathbb{P}(A \mid B) \leq p$, by the Bayes identity in the proof above with $\kappa = (1 - p)/p$. Under $\mathbb{P}_B$ the member A is realized with probability w , so the joint event has probability $w \mathbf{1}\{w \leq p\} \leq p$. $\square$

The corollary is the precise form of the slogan that a rare realized attribute establishes its own significance: given the B -conditioned draw, a rare member is realized-and-unproductive with probability at most its own prior. The slogan is a within-model diagnostic, and it is not evidence for the draw.

Proposition 3.4 (Conjunction: the stockpile). *Let $F_1, \dots, F_n$ be a predeclared family of attributes with priors $p_i = \mathbb{P}(F_i)$, let $F = \bigcap_i F_i$, and suppose the family is submultiplicative, $\mathbb{P}(F) \leq \prod_i p_i$, which holds with equality under independence and holds under negative association. Then, under the B -conditioned draw of the partition $\{F, F'\}$,*

$$\mathbb{P}_B[A^* = F \text{ and } \mathbb{P}(B \mid F') \geq \mathbb{P}(B \mid F)] \leq \mathbb{P}(F) \leq \prod_i p_i : \quad (3)$$

the joint event that the conjunction is realized and is the less B -productive member of its pair has probability at most the product of the priors, which tightens multiplicatively in n while the complement, the union of the F'_i , accumulates weight $1 - \prod_i p_i$. Positive association gives $\mathbb{P}(F) \geq \prod_i p_i$ and defeats the bound; it is not a sufficient condition.

Proof. Corollary 3.3 applied to the fixed member F of the partition $\{F, F'\}$ gives the first inequality when $\mathbb{P}(F) < \frac{1}{2}$; the second is submultiplicativity. $\square$

Proposition 3.5 (The multiplicity guard: expectation and tail). *If n partitions are interrogated under the B -conditioned draw and N counts those whose realized member exhibits a κ -fold*

exceedance of (1), then $\mathbb{E}[N] \leq n/(1 + \kappa)$; and, with no assumption on the dependence among the partitions,

$$\Pr[N \geq m] \leq \frac{n}{m(1 + \kappa)}, \quad \Pr[N \geq 1] \leq \frac{n}{1 + \kappa}, \quad (4)$$

by Markov's inequality and the union bound respectively. Under independence or a stated dependence structure the tail sharpens. Exceeding the expectation is not significance: a declared family is scored against the tail (4), and a family whose size n is not declared cannot be scored at all.

Proof. Linearity of expectation over the n indicators, each of probability at most $1/(1 + \kappa)$ by Theorem 3.1; Markov on N ; and the union bound on the n events. $\square$

4 Survival targets and the horizon family

The target of most interest is indefinite survival, and in the critical regime it cannot be conditioned on directly: $\{T = \infty\}$ is null, and $\mathbb{P}(\cdot \mid T = \infty)$ is undefined. The right objects are the horizon events.

Theorem 4.1 (Horizon-indexed accordance; the second statement at the grade of the specialization in force). *Let $B_t = \{T > t\}$ and $Q_t = \mathbb{P}(\cdot \mid B_t)$. For every finite t and every binary partition $\{A, A'\}$ of $\mathcal{F}_s$, $s < t$, if the realized member is drawn under Q_t then (1) holds with $B = B_t$ and $\mathbb{P}_B = Q_t$. In the critical regime, as $t \rightarrow \infty$ the conditioned law Q_t converges to the Q -process on $\mathcal{F}_s$, and for each cell*

$$\frac{\mathbb{P}(T > t \mid A)}{\mathbb{P}(T > t)} \longrightarrow \frac{Q(A)}{\mathbb{P}(A)} = \mathbb{E}_{\mathbb{P}}[Z_s \mid A], \quad (5)$$

so that

$$\frac{\mathbb{P}(T > t \mid A')}{\mathbb{P}(T > t \mid A)} \longrightarrow \frac{Q(A')/\mathbb{P}(A')}{Q(A)/\mathbb{P}(A)}. \quad (6)$$

Proof. The first statement is Theorem 3.1 with the target B_t . For (5), $\mathbb{P}(T > t \mid A)/\mathbb{P}(T > t) = \mathbb{E}_{\mathbb{P}}[\mathbf{1}_A \mathbb{P}(T > t \mid \mathcal{F}_s)/\mathbb{P}(T > t)]/\mathbb{P}(A)$, and by Definition 2.3 $\mathbb{P}(T > t \mid \mathcal{F}_s)/\mathbb{P}(T > t) \rightarrow Z_s$ in $L^1(\mathbb{P})$ on the survival event, giving $\mathbb{E}_{\mathbb{P}}[Z_s \mathbf{1}_A]/\mathbb{P}(A) = Q(A)/\mathbb{P}(A)$; (6) is the quotient of two instances. $\square$

Remark 4.2 (Grade propagation in this section). Every statement in this section that consumes the survivorship companion's conditioned law inherits that law's grade: with the hard-wall specialization the statements are at $[\mathbf{P}]$; with the soft-wall specialization they are conditional on the companion's ratio hypothesis, Ansatz B, and carry that conditionality wherever they are used. Theorem 3.1 and its corollaries depend on neither and remain unconditional.

Remark 4.3 (The ideal limit is a limit). Every statement about “conditioning on indefinite survival” in this paper and its companions is to be read through Theorem 4.1: a statement at finite horizon, and then a limit. The Q -process is not a conditional probability on a null event, and expressions of the form $\mathbb{P}(G \mid A)$ with G the event of indefinite survival are shorthand for $\lim_t \mathbb{P}(B_t \mid A)$ where that limit exists. In the survivorship model the persistence-harmonic function $\mathfrak{h}$ is what the limit produces, and the attribute-level ratios are ratios of its averages.

5 Rival laws for a record: the between-model layer

Definition 5.1 (Hypotheses about the outer measure). For a record with law $\mathbb{P}$ and a target family B_t , let $H_{\mathbb{P}}$ be the hypothesis that the record’s realized attributes are drawn under $\mathbb{P}$ (no conditioning), and H_Q the hypothesis that they are drawn under Q (the survivorship-conditioned law, through its horizon family). Other targets, an observer-existence condition among them, define other hypotheses in the same way.

Proposition 5.2 (The Bayes factor is the Doob density). *For a predeclared record statistic $E \in \mathcal{F}_s$, and with both laws conditioned on the common baseline C of Remark 5.3 wherever an observer’s record is compared, the Bayes factor between the conditioned and unconditioned laws is*

$$\text{BF}_{Q:\mathbb{P}}(E) = \frac{Q(E \mid C)}{\mathbb{P}(E \mid C)}, \quad (7)$$

which reads $Q(E)/\mathbb{P}(E)$ when no baseline is shared; and on the full information $\mathcal{F}_s$ it is the likelihood ratio Z_s of Definition 2.3, the hard-wall Doob density or the soft Feynman–Kac density according to the specialization in force. If E is required by the target, $Q(E) \approx 1$, and rare under $\mathbb{P}$, $\mathbb{P}(E) = q$, then $\text{BF}_{Q:\mathbb{P}}(E) \approx 1/q$.

Proof. (7) is the definition of the Bayes factor for a simple event; on $\mathcal{F}_s$ the factor is the Radon–Nikodym density by definition, and the density’s form is Definition 2.3. $\square$

Remark 5.3 (The null for a surviving observer is the survivor law, not raw $\mathbb{P}$). Under raw $\mathbb{P}$ there is positive mass on $\{T \leq s\}$, where $Z_s = 0$ and $\log Z_s = -\infty$. That is correct as mathematics and wrong as a null for an observer whose record already exists at s : current survival is evidence both hypotheses share, and a comparison that leaves it in credits the conditioned law for the trivial fact that the observation was made. The null for the observed surviving cohort is $\mathbb{P}_s^{\text{surv}} := \mathbb{P}(\cdot \mid T > s)$, against which the likelihood ratio on $\mathcal{F}_s$ is

$$\left. \frac{dQ}{d\mathbb{P}_s^{\text{surv}}} \right|_{\mathcal{F}_s} = \mathbb{P}(T > s) Z_s, \quad (8)$$

which in the hard-wall specialization is $\mathbb{P}(T > s) \mathfrak{h}(X_s)/\mathfrak{h}(X_0)$, finite on the whole cohort. More generally every rival law in a comparison is conditioned on the common baseline C the record fixes, observers existing now and survival to the observation time, and the Bayes factor is $\mathbb{P}(E \mid C, H_1)/\mathbb{P}(E \mid C, H_2)$, so that no part of it is earned by evidence both rivals already had to explain.

Remark 5.4 (Why this and not the theorem carries the inference). A low-probability realization under $H_{\mathbb{P}}$ is not a contradiction of $H_{\mathbb{P}}$; it is evidence against it, and (7) says how much. The attribute-level accordance of Theorem 3.1 tells one what to expect of a record if H_Q holds; the Bayes factor tells one whether H_Q holds better than a rival. A record can accord with Q at every interrogated attribute and still be better explained by a rival law that also predicts those attributes, which is why the comparison must name its rival and predeclare its statistic. The same object, the Doob density, serves both layers: within the model it is the weight by which the conditioning tilts each path, and between models it is the evidence the tilted path carries.

Definition 5.5 (Realized evidence, information, and flux). Let $Z_t = dQ|_{\mathcal{F}_t}/d\mathbb{P}|_{\mathcal{F}_t}$ and $\ell_t = \log Z_t$, the realized log-evidence. Since Z is a $\mathbb{P}$ -martingale, ℓ_t is not monotone along a path: new observations can favor Q , then $\mathbb{P}$. The monotone quantity is the *information*

$$I_t := \mathbb{E}_Q[\ell_t] = D_{\text{KL}}(Q|_{\mathcal{F}_t} \parallel \mathbb{P}|_{\mathcal{F}_t}), \quad (9)$$

nondecreasing in t as the filtration grows, with $\mathbb{E}_{\mathbb{P}}[\ell_t] = -D_{\text{KL}}(\mathbb{P}|_{\mathcal{F}_t}||Q|_{\mathcal{F}_t}) \leq 0$. Where differentiable, $\mathcal{J}_t := dI_t/dt$ is the *information flux*.

Proposition 5.6 (Discriminating rival laws by the law of the evidence). *Under H_Q the realized log-evidence has mean $I_t \geq 0$ and, in the critical regime of the survivorship model, $I_t \rightarrow \infty$; under $H_{\mathbb{P}}$ it has mean $-D_{\text{KL}}(\mathbb{P}_t||Q_t) \leq 0$. The two hypotheses are therefore discriminated by the distribution of ℓ_t , of which the flux is the summary, and not by any monotone accumulation along a single path, which does not exist.*

Proof. The means are (9) and its $\mathbb{P}$ -counterpart; divergence under H_Q in the critical regime follows from Theorem 2.4, since mutual singularity on $\mathcal{F}_{\infty}$ forces the relative entropy to diverge. $\square$

6 The limit on certification

Theorem 6.1 (No bounded observation certifies the law; grade as Theorem 2.4). *In the critical regime, for every finite s , no event in $\mathcal{F}_s$ has probability one under Q and probability zero under $\mathbb{P}$ on the survival event, nor conversely. Consequently no observation of bounded duration certifies whether a surviving record is drawn under Q or under $\mathbb{P}$; certification is available only on $\mathcal{F}_{\infty}$, where the laws are singular.*

Proof. Mutual absolute continuity on the survival event at every finite s is Theorem 2.4; an event of probability one under one law and zero under the other would violate it. $\square$

Remark 6.2 (Certification against discrimination). The theorem forbids perfect separation and nothing more. Mutually absolutely continuous laws can be statistically far apart, and Proposition 5.2 is exactly the measure of how far: a Bayes factor of 10^8 on a finite window is discrimination of a high order, and it is not certification, because the same window has positive probability under both laws. The theorem is turned on this paper’s own subject matter deliberately: whatever a record’s attributes say in favor of H_Q , they say it at a Bayes factor and never at probability one.

7 A protocol for applying the theorem to a record

The theorem applies to any record with a law and a target. Its application to an actual record faces the retrospective-target objection in its full force: any realized history, read backward, contains rare attributes that are prerequisites of whatever it happens to contain, so a set of attributes selected after the fact and found to accord with a target chosen after the fact establishes nothing. Two of the objection’s three freedoms are closed by the theorem’s form: the target is fixed before the record is read, and the alternative to each attribute is its complement, not an analyst’s choice of class. The third freedom, which attributes to interrogate, is not closed by any theorem, and the multiplicity guard prices it only when the scan is declared. The protocol is therefore prospective.

Definition 7.1 (Prospective test of a target hypothesis). Declare, before the observation interval: (i) the target family B_t and the rival law; (ii) the attribute family $F_1, \dots, F_n$, with n fixed; (iii) the baseline priors p_i under the rival law and the dependence assumption on the family (independence, or negative association, or a stated bound on $\mathbb{P}(\bigcap F_i)$); (iv) the record statistic

E whose Bayes factor (7) will be reported; (v) the stopping rule and the multiplicity budget $n/(1 + \kappa)$ at the κ to be scored. Then observe. A retrospective exercise on an existing record, however suggestive, is hypothesis-generating and is reported as such.

8 Open problems

Every open item is gathered here. The soft-wall ratio hypothesis on which the survivorship model’s Q -process rests is open in the companion [1], and Theorem 4.1’s second statement inherits it. The convergence of Q_t on a window shifted to the present, the law of $(X_{t+u})_{u \leq s}$ given survival to t , is a different statement from convergence on an early window and is not established; every claim about a surviving record’s *future* behavior waits on it. A finite, monotone evidence budget under which a Copernican position statement would be well posed is not defined, and no Lindy-type result is claimed. And the extension of the multiplicity guard from a fixed family to a sequentially chosen one, with a stopping rule, is stated as a protocol requirement rather than proved as a bound.

9 Conclusion

The claim stated at the outset stands as proved. If the realized member of a binary partition of the past is drawn under a law conditioned on a future target, then for every $\kappa > 1$ it is, with probability at least $1 - 1/(1 + \kappa)$, within a κ -fold exceedance of being the more target-productive member, and the statement is indifferent to the target’s tense; whether a given record is drawn under such a law is a separate hypothesis, tested by comparing measures, and the measure of that comparison is the Doob density. The future does nothing to the past in any of this. A conditioned law has a support and a weight, and the past of a weighted history carries the weight; that is the whole of what “conditioning a record on its own future” means, and it is enough to make the companion’s hypothesis about the record of natural history a hypothesis with a stated test rather than a reading with a stated feeling.

References

- [1] Ajax Benander. *Survivorship Conditioning for Hazard Rates Driven at the Log-Derivative Level*. Working draft. The Golden branch’s root extraction; the two-condition viability theorem with its piecewise driver-viability functional, the hard-wall and soft-wall conditioned laws, the certification ceiling, and the half-time theorem. 2026. URL: <https://aethus.org>.
- [2] Ajax Benander. “Nexic Reasoning: Defining a Generalized Calculus Over Anthropic Parameters”. In: (Dec. 2024). Manuscript; the anthropic wing’s master, successive deposits December 2024–2026. Also at <https://osf.io/jvkcp>. This version: September 2026. DOI: [10.31219/osf.io/jvkcp](https://doi.org/10.31219/osf.io/jvkcp). URL: <https://philpapers.org/rec/BENNRD>.
- [3] B. Roynette, P. Vallois, and M. Yor. “Some penalisations of the Wiener measure”. In: *Japanese Journal of Mathematics* 1 (2006), pp. 263–290.
- [4] Sylvie Méléard and Denis Villemonais. “Quasi-stationary distributions and population processes”. In: *Probability Surveys* 9 (2012), pp. 340–410.
- [5] Nicolas Champagnat and Denis Villemonais. “Exponential convergence to quasi-stationary distribution and Q -process”. In: *Probability Theory and Related Fields* 164 (2016). AUTHOR TO VERIFY page range, pp. 243–283.