

MATH 455 (Honours Analysis 4)

J. Han

January 13, 2024

The following notes are adapted from lectures given by Dmitry Jakobson in Winter 2024. All errors in interpretation, reasoning, coherence, and articulation are my own.

This document's source code, located at <https://github.com/brunefig/math455/blob/main/notes.org>, can be converted into Anki flashcards with the [anki-editor](#) package for [GNU Emacs](#). Flashcard [cloze deletions](#) are typeset in [magenta](#).

preliminaries

topological spaces 2

2024-01-05 topology & metric spaces

connectedness 2

examples of metric spaces 3

p -adic numbers 3

convexness 4

2024-01-10 contraction mappings & product topologies

total disconnectedness 4

contraction mappings 4

iterated function systems and fixed point sets 6

basis for a topology 6

product topologies 7

projection maps 7

2024-01-12 Hilbert spaces & Hausdorff spaces

inner product spaces 8

Hilbert spaces 8

Hausdorff spaces 9

preliminaries

topological spaces

A **topological space** (X, τ) is a **set** X endowed with a **topology** τ .

A **topology** τ on a set X is a collection of sets that contains \emptyset and X and is closed under **finite intersections and arbitrary unions**. Membership in τ defines an **open set** in the topological space (X, τ) .

2024-01-05 topology & metric spaces

connectedness

A topological space (X, τ) is a **connected space** if and only if $\neg \exists U, U' \in \tau \setminus \{\emptyset\} : U \cap U' = \emptyset \wedge X = U \cup U'$.

For a topological space (X, τ) , $A \subseteq X$ is a **connected set** if and only if $\neg \exists U, U' \in \tau : A \cap U \neq \emptyset, A \cap U' \neq \emptyset, U \cap U' = \emptyset, A = (U \cap A) \cup (U' \cap A)$.

A topological space X is a **path connected space** if and only if $\forall x, x' \in X : \exists \text{ continuous } f : [0, 1] \rightarrow X : [f(0) = x \wedge f(1) = x']$.

All path connected spaces are **connected spaces**.

For **open sets** in \mathbb{R}^n , **connectedness** is equivalent to **path connectedness**.

A topological space X can be expressed as **the disjoint union of maximal connected subsets**, where a connected subset is called maximal if and only if **it has no connected superset in X** . These subsets are the **connected components** of X .

A topological space X can be expressed as **the disjoint union of maximal path connected subsets**, where a path connected subset is called maximal if and only if **it has no path connected superset in X** . These subsets are the **path components** of X .

A **path connected** space has exactly one **path component**.

$\left\{ \left(x, \sin\left(\frac{1}{x}\right) \right) \mid x \in (0, 1] \right\} \cup \{(0, 0)\} \subseteq \mathbb{R}^2$ is **connected** but not **path connected** because **it has two path components**.

If $f : X \rightarrow \mathbb{R}$ is continuous, $A \subseteq X$ is connected, and $x, x' \in f(A)$ such that $x < x'$, then $[x, x'] \subseteq f(A)$.

Proof. Suppose $\exists c \in (f(x'), f(x)) : c \notin f(A)$. Since $f^{-1}((-\infty, c))$ and $f^{-1}((c, \infty))$ are open, $A \subseteq f^{-1}((-\infty, c)) \cup f^{-1}((c, \infty))$ is not connected. \square

examples of metric spaces

Any normed vector space is a metric space with the induced metric $d(x, x') := \|x - x'\|$.

For $p \in (0, \infty)$, $l_p := \{(x_n)_{n \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} \mid \sum_{n \in \mathbb{N}} |x_n|^p < \infty\}$ is a normed vector space with $\|x\|_p := (\sum_{n \in \mathbb{N}} |x_n|^p)^{1/p}$.

The sequence $(\frac{1}{n})_{n \in \mathbb{Z}_+}$ is a member of l_p if and only if $p > 1$.

Proof. $(\frac{1}{n})_{n \in \mathbb{Z}_+} \in l_p \iff \sum_{n \in \mathbb{Z}_+} (\frac{1}{n})^p < \infty \iff p > 1$. \square

For $p \in [1, \infty)$, $L^p([a, b]) := \left\{ f(x) \mid \int_a^b |f(x)|^p dx < \infty \right\}$ is a normed vector space with $\|f\|_p := \left(\int_a^b |f(x)|^p dx \right)^{1/p}$.

$d(A, A') := \text{vol}_n(A \Delta A')$ is a possible metric on \mathbb{R}^n .

For a metric space (X, d) , a set $A \subseteq X$, and $\epsilon > 0$, let $A_\epsilon := \bigcup_{x \in A} B(x, \epsilon)$. Then the Hausdorff metric is $d_H(A, A') := \inf\{\epsilon > 0 \mid A' \subseteq A_\epsilon \wedge A \subseteq A'_\epsilon\}$.

p -adic numbers

Given a fixed prime p ,

$$\forall q \in \mathbb{Q} \setminus \{0\} : \exists (a, b, n) \in \mathbb{Z}^3 : \left[q = p^n \cdot \frac{a}{b} \wedge \gcd(a, p) = \gcd(b, p) = 1 \right]$$

and \mathbb{Q} is a normed vector space with $\|q\|_p := \begin{cases} p^{-n} & q \neq 0 \\ 0 & q = 0 \end{cases}$.

The 2-adic norm of $\frac{96}{7}$ is $\frac{1}{32}$.

The 3-adic norm of $3^{-2024} \cdot \frac{8}{13}$ is 3^{2024} .

The p -adic norm $\|q\|_p$ is small if q is divisible by a large power of p .

The p -adic norm of 0 is 0 because 0 is divisible by any power of p .

(p -adic product formula.) If $q \in \mathbb{Q} \setminus \{0\}$, then $|q| \cdot \prod_{p \text{ prime}} \|q\|_p = 1$.

convexness —————

A set X is **convex** if and only if the line segment joining any two points in X lies within X .

2024-01-10 contraction mappings & product topologies

$\alpha \in I$ henceforth refers to members of a possibly uncountable index set I .

If $\{A_\alpha\}_{\alpha \in I}$ is a family of **connected** sets in a topological space X such that $\forall \alpha, \alpha' \in I : A_\alpha \cap A_{\alpha'} \neq \emptyset$, then $\bigcup_{\alpha \in I} A_\alpha$ is **connected**.

Proof. Suppose $A := \bigcup_{\alpha \in I} A_\alpha$ is not connected. Let $U, U' \subseteq X$ be open and nonempty relative to A such that $U \cap U' = \emptyset$ and $A = U \cup U'$. Since each A_α is connected, there is no α for which $A_\alpha \cap U \neq \emptyset \neq A_\alpha \cap U'$. But then $\exists \alpha, \alpha' \in I : A_\alpha \subseteq U \wedge A_{\alpha'} \subseteq U'$, which contradicts that $A_\alpha \cap A_{\alpha'} \neq \emptyset$. \square

total disconnectedness —————

A topological space X is **totally disconnected** if and only if every connected subset of X is a singleton.

A **totally disconnected** space in \mathbb{R} contains **only points** and **no intervals**.

The **Cantor set** is a totally disconnected space in \mathbb{R} .

contraction mappings —————

For a metric space (X, d) , $T : X \rightarrow X$ is a **contraction mapping** if and only if $\exists c \in [0, 1) : \forall x, x' \in X : d(T(x), T(x')) \leq c \cdot d(x, x')$.

All contraction mappings are **continuous**.

(Orbit lemma.) For $x \in X$, the **orbit** $(T^n(x))_{n \in \mathbb{N}}$ of a **contraction mapping** T on X is a **Cauchy sequence**.

Proof. For $n \in \mathbb{Z}_+$,

$$\begin{aligned} d(T^n(x), T^{n+1}(x)) &< c \cdot d(T^{n-1}(x), T^n(x)) \\ &< c^2 \cdot d(T^{n-2}(x), T^{n-1}(x)) \\ &\dots < c^n \cdot d(x, T(x)). \end{aligned}$$

Let $m \geq n$. By the triangle inequality,

$$\begin{aligned} d(T^n(x), T^m(x)) &\leq \sum_{k=n}^{m-1} d(T^k(x), T^{k+1}(x)) \\ &\leq d(x, T(x)) \sum_{k=n}^{m-1} c^k \\ &= c^n \cdot d(x, T(x)) \sum_{k=0}^{m-n-1} c^k \\ &\leq \frac{c^n \cdot d(x, T(x))}{1-c}. \end{aligned}$$

For $\epsilon > 0$, choosing $n > \log_c \left(\frac{\epsilon}{2} \cdot \frac{1-c}{d(x, T(x))} \right)$ and $m, m' \geq n$ guarantees

$$\begin{aligned} d(T(m), T(m')) &\leq \\ d(T(n), T(m)) + d(T(n), T(m')) &< \epsilon. \end{aligned}$$

□

(*Contraction mapping theorem.*) If (X, d) is a *nonempty* and *complete* metric space and T is a *contraction mapping on X* , then

$$\exists! z \in X : T(z) = z,$$

i.e. z is *the unique fixed point*, and

$$\forall x \in X : \lim_{n \rightarrow \infty} T^n(x) = z,$$

where $\forall n \in \mathbb{N} : T^n(x) := T(T(\dots T(x)))$.
n times

Proof. For $x \in X$, $T(x)$ is Cauchy by the orbit lemma, and since (X, d) is complete it converges to some point $z \in X$. Then z is a fixed point of T because

$$\begin{aligned} T^n(x) \rightarrow z &\implies T(T^n(x)) \rightarrow T(z) \\ &\implies T(T^n(x)) = T^{n+1}(x) \rightarrow z = T(z), \end{aligned}$$

and is its unique fixed point because

$$T(z) = z \wedge T(z') = z' \implies d(z, z') \leq c \cdot d(z, z') \iff d(z, z') = 0. \quad \square$$

$x \mapsto \frac{x}{2}$ is a contraction mapping on $(\mathbb{R}, |\cdot|)$ with fixed point $z = 0$.

iterated function systems and fixed point sets

$\mathcal{K}(X)$ henceforth denotes the set of compact subsets of a set X .

If $m \in \mathbb{Z}_+$, $\{T_n : \mathbb{R}^d \rightarrow \mathbb{R}^d\}_{n \in [m]}$ are contraction mappings, and

$$F : (\mathcal{K}(\mathbb{R}^d), d_H) \rightarrow (\mathcal{K}(\mathbb{R}^d), d_H) : A \mapsto \bigcup_{n \in [m]} T_n(A),$$

then $\exists! A \in \mathcal{K}(\mathbb{R}^d) : F(A) = A$. This is the **fixed point set** of F .

Proof. F is a contraction mapping (to be demonstrated in problem set 2), and $(\mathcal{K}(X), d_H)$ is a complete metric space (???). A unique fixed point set thus exists by the contraction mapping theorem. \square

Let $T_0, T_1 : \mathbb{R} \rightarrow \mathbb{R}$ such that $T_0(x) := \frac{x}{3}$ and $T_1(x) := \frac{x+2}{3}$ and $F(A) : \mathcal{K}(\mathbb{R}) \rightarrow \mathcal{K}(\mathbb{R})$ such that $F(A) := T_0(A) \cup T_1(A)$. Then $F([0, 1]) = T_0([0, 1]) \cup T_1([0, 1]) = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$, and the **fixed point set** of F is the **middle-thirds Cantor set**.

If $T_0, T_1 : \mathbb{R}^d \rightarrow \mathbb{R}^d$ are **contraction mappings** and $F : (\mathcal{K}(\mathbb{R}^d), d_H) \rightarrow (\mathcal{K}(\mathbb{R}^d), d_H)$ such that $F(A) = T_0(A) \cup T_1(A)$, then the composition of F with itself is

$$F^2(A) = T_0(T_0(A)) \cup T_0(T_1(A)) \cup T_1(T_0(A)) \cup T_1(T_1(A)).$$

basis for a topology

A **basis** \mathcal{B} for a topology τ is a **subset of τ such that $\forall A \in \tau : \exists \{A_\alpha\}_{\alpha \in I} \subseteq \mathcal{B} : A = \bigcup_{\alpha \in I} A_\alpha$** .

The **open intervals** form a basis for the **standard topology** on \mathbb{R} .

For a metric space X , $\{B(x, \epsilon) \mid x \in X, \epsilon > 0\}$ is a basis for the **open sets in X** .

If \mathcal{B} is a basis for the topology on X , then $\forall x \in X : \exists U \in \mathcal{B} : x \in U$.

If \mathcal{B} is a basis for the topology τ on X , then $[U, U' \in \mathcal{B} \wedge x \in U \cap U'] \implies \exists U'' \in \tau : x \in U'' \subseteq U \cap U'$.

If $f : X \rightarrow Y$ for **topological spaces** (X, τ_X) and (Y, τ_Y) and \mathcal{B} is a basis for τ_Y , then f is **continuous** if and only if $\forall U \in \mathcal{B} : f^{-1}(U) \in \tau_X$.

product topologies

For **topological spaces** $\{(X_\alpha, \tau_\alpha)\}_{\alpha \in I}$, $\prod_{\alpha \in I} A_\alpha$ is a **cylinder set** in $X := \prod_{\alpha \in I} X_\alpha$ if and only if $\forall \alpha \in I : A_\alpha \in \tau_\alpha$.

If $X := \mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$, then \forall **open intervals** $(a, b), (a', b') \subseteq \mathbb{R} : (a, b) \times (a', b')$ is a **cylinder set** in X .

For **topological spaces** $\{X_\alpha\}_{\alpha \in I}$, consider a **cylinder set** $\prod_{\alpha \in I} A_\alpha$ in $X := \prod_{\alpha \in I} X_\alpha$ such that $\exists I' \subseteq I : [|I'| \in \mathbb{N} \wedge \forall \alpha \in I \setminus I' : A_\alpha = X_\alpha]$. These base cylinder sets form a basis for **the product topology on X** .

projection maps

For a vector $x := (x_\alpha)_{\alpha \in I}$ in possibly uncountable dimensions, the function $\pi_\alpha(x) := x_\alpha$ is a **projection map**.

For topological spaces $\{X_\alpha\}_{\alpha \in I}$ let $f : Y \rightarrow \prod_{\alpha \in I} X_\alpha$, and for $\alpha \in I$ let $f_\alpha : Y \rightarrow X_\alpha : y \mapsto \pi_\alpha(f(y))$. Then f is **continuous** if and only if $\forall \alpha \in I : f_\alpha$ is **continuous**.

Proof. Suppose f is continuous. Then, for $\alpha' \in I$ and an open set $U \subseteq X_{\alpha'}$,

$$f_{\alpha'}^{-1}(U) = f^{-1} \left(U \times \prod_{\alpha \in I \setminus \{\alpha'\}} X_\alpha \right)$$

is open as the preimage of a base cylinder set in $\prod_{\alpha \in I} X_\alpha$.

Suppose $\forall \alpha \in I : f_\alpha$ is continuous. It suffices to verify that the preimage of f for a base cylinder set U is open. Let $I' \subseteq I$ be a finite index subset for which

$$U = \prod_{\alpha \in I'} A_\alpha \times \prod_{\alpha \in I \setminus I'} X_\alpha.$$

Then

$$\begin{aligned} f^{-1}(U) &= \left(\bigcap_{\alpha \in I'} f_\alpha^{-1}(A_\alpha) \right) \cap \left(\bigcap_{\alpha \in I \setminus I'} f_\alpha^{-1}(X_\alpha) \right) \\ &= \bigcap_{\alpha \in I'} f_\alpha^{-1}(A_\alpha) \end{aligned}$$

is open in Y as the finite intersection of open sets in Y . □

2024-01-12 Hilbert spaces & Hausdorff spaces

inner product spaces

The closed unit ball in l_p is not **compact** because it contains $(x_n)_{n \in \mathbb{N}}$ where

$$\forall n, k \in \mathbb{N} : \pi_k(x_n) = \begin{cases} 1 & k = n \\ 0 & k \neq n \end{cases},$$

which has no convergent subsequence.

An **inner product space** V is a **normed vector space** with an **inner product operation** $\langle v, v' \rangle$ for $v, v' \in V$ that defines its **norm** $\|v\|^2 = \langle v, v \rangle$.

(**Parallelogram law**.) The **norm** in a vector space V is **induced by an inner product** if and only if

$$\forall v, v' \in V : \|v + v'\|^2 + \|v - v'\|^2 = 2(\|v\|^2 + \|v'\|^2)$$

holds.

(**Hanner's inequality**.) For a measure space X , $p \in [1, \infty) \setminus \{2\}$, and $f, g \in L^p(X)$,

$$\|f + g\|_p^p + \|f - g\|_p^p * (\|f\|_p + \|g\|_p)^p + \left| \|f\|_p - \|g\|_p \right|^p$$

where $*$:= $\begin{cases} \geq & p < 2 \\ \leq & p > 2 \end{cases}$. If $p = 2$, then this becomes **the parallelogram law**.

Hilbert spaces

A **Hilbert space** is a **complete inner product space**.

l_2 is a **Hilbert space** with $\langle (a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \rangle = \sum_{n \in \mathbb{N}} |a_n \overline{b_n}|$.

For a measure space X , $L^2(X)$ is a **Hilbert space** with $\langle f, g \rangle := \int_X f(x) \overline{g(x)} dx$.

(**Parseval's identity**.) For $f \in L^2([0, 2\pi])$,

$$\|f\|_2^2 = \int_0^{2\pi} |f(x)|^2 dx = 2\pi \sum_{n=-\infty}^{\infty} |\hat{f}(n)|^2 = \langle f, f \rangle$$

where $\hat{f}(n) = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-inx} dx$.

(Consequence of **Riesz's lemma**.) The **closed unit ball** in a **Hilbert space** V is **compact** if and only if V is **finite-dimensional**.

Hausdorff spaces

A topological space (X, τ) is a **Hausdorff space** if and only if $\forall x \neq x' \in X : \exists U, U' \in \tau : [U \cap U' = \emptyset \wedge (x, x') \in U \times U']$.

All **metric spaces** are Hausdorff spaces.

*Singltons in a **Hausdorff space** X with topology τ are **closed**.*

Proof. Let $x' \in X \implies$. Then

$$\forall x \in X : \exists U_x \in \tau : [x \in U_x \wedge x' \notin U_x] \implies X \setminus \{x'\} = \bigcup_{x' \in X \setminus \{x'\}} U_x \in \tau.$$

□

$\tau := \{\emptyset, X\}$ is the **trivial topology** on X .

If τ is the **trivial topology** and $|X| > 1$, then (X, τ) is **not a Hausdorff space**.

The **subspace topology** on a subspace Y of a topological space (X, τ) is $\tau_Y := \{U \cap Y \mid U \in \tau\}$.

A subspace of a **Hausdorff space** is a **Hausdorff space** under the **subspace topology**.