

MATH 455 (Honours Analysis 4)

J. Han

January 18, 2024

The following notes are adapted from lectures given by Dmitry Jakobson in Winter 2024. All errors in interpretation, reasoning, coherence, and articulation are my own.

This document's source code, located at <https://github.com/brunefig/math455/blob/main/notes.org>, can be converted into Anki flashcards with the [anki-editor](#) package for [GNU Emacs](#). Flashcard [cloze deletions](#) are typeset in [magenta](#).

preliminaries

topological spaces 2

2024-01-05 topology & metric spaces

connectedness 2

examples of metric spaces 3

p -adic numbers 3

convexness 4

2024-01-10 contraction mappings & product topologies

total disconnectedness 4

contraction mappings 5

iterated function systems and fixed point sets 6

basis for a topology 7

product topologies 7

projection maps 7

2024-01-12 Hilbert spaces & Hausdorff spaces

inner product spaces 8

Hilbert spaces 9

compact sets & continuity 9

Hausdorff spaces	10
normal spaces	11
Lebesgue number	11
2024-01-17 product spaces & separable spaces	
metrizable	12
countable product of compact metric spaces	13
separability	14

preliminaries

topological spaces

A **topological space** (X, τ) is a **set** X endowed with a **topology** τ .

A **topology** τ on a set X is a collection of sets that contains \emptyset and X and is closed under **finite intersections and arbitrary unions**. Membership in τ defines an **open set** in the topological space (X, τ) .

2024-01-05 topology & metric spaces

connectedness

A topological space (X, τ) is a **connected space** if and only if $\neg \exists U, U' \in \tau \setminus \{\emptyset\} : U \cap U' = \emptyset \wedge X = U \cup U'$.

For a topological space (X, τ) , $A \subseteq X$ is a **connected set** if and only if $\neg \exists U, U' \in \tau : A \cap U \neq \emptyset, A \cap U' \neq \emptyset, U \cap U' = \emptyset, A = (U \cap A) \cup (U' \cap A)$.

A topological space X is a **path connected space** if and only if $\forall x, x' \in X : \exists$ continuous $f : [0, 1] \rightarrow X : [f(0) = x \wedge f(1) = x']$.

All path connected spaces are **connected spaces**.

For **open** sets in \mathbb{R}^n , **connectedness** is equivalent to **path connectedness**.

A topological space X can be expressed as the **disjoint union of maximal connected subsets**, where a connected subset is called maximal if and only if it has no **connected superset in X** . These subsets are the **connected components** of X .

A topological space X can be expressed as the disjoint union of maximal path connected subsets, where a path connected subset is called maximal if and only if it has no path connected superset in X . These subsets are the path components of X .

A path connected space has exactly one path component.

$\left\{ \left(x, \sin\left(\frac{1}{x}\right) \right) \mid x \in (0, 1] \right\} \cup \{(0, 0)\} \subseteq \mathbb{R}^2$ is connected but not path connected because it has two path components.

If $f : X \rightarrow \mathbb{R}$ is continuous, $A \subseteq X$ is connected, and $x, x' \in f(A)$ such that $x < x'$, then $[x, x'] \subseteq f(A)$.

Proof. Suppose $\exists c \in (f(x'), f(x)) : c \notin f(A)$. Since $f^{-1}((-\infty, c))$ and $f^{-1}((c, \infty))$ are open, $A \subseteq f^{-1}((-\infty, c)) \cup f^{-1}((c, \infty))$ is not connected. \square

examples of metric spaces

Any normed vector space is a metric space with the induced metric $d(x, x') := \|x - x'\|$.

For $p \in (0, \infty)$, $l_p := \{(x_n)_{n \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} \mid \sum_{n \in \mathbb{N}} |x_n|^p < \infty\}$ is a normed vector space with $\|x\|_p := (\sum_{n \in \mathbb{N}} |x_n|^p)^{1/p}$.

The sequence $(\frac{1}{n})_{n \in \mathbb{Z}_+}$ is a member of l_p if and only if $p > 1$.

Proof. $(\frac{1}{n})_{n \in \mathbb{Z}_+} \in l_p \iff \sum_{n \in \mathbb{Z}_+} (\frac{1}{n})^p < \infty \iff p > 1$. \square

For $p \in [1, \infty)$, $L^p([a, b]) := \left\{ f(x) \mid \int_a^b |f(x)|^p dx < \infty \right\}$ is a normed vector space with $\|f\|_p := \left(\int_a^b |f(x)|^p dx \right)^{1/p}$.

$d(A, A') := \text{vol}_n(A \Delta A')$ is a possible metric on \mathbb{R}^n .

For a metric space (X, d) , a set $A \subseteq X$, and $\epsilon > 0$, let $A_\epsilon := \bigcup_{x \in A} B(x, \epsilon)$. Then the Hausdorff metric is $d_H(A, A') := \inf\{\epsilon > 0 \mid A' \subseteq A_\epsilon \wedge A \subseteq A'_\epsilon\}$.

p -adic numbers

Given a fixed prime p ,

$$\forall q \in \mathbb{Q} \setminus \{0\} : \exists (a, b, n) \in \mathbb{Z}^3 : \left[q = p^n \cdot \frac{a}{b} \wedge \gcd(a, p) = \gcd(b, p) = 1 \right]$$

and \mathbb{Q} is a normed vector space with $\|q\|_p := \begin{cases} p^{-n} & q \neq 0 \\ 0 & q = 0 \end{cases}$.

The 2-adic norm of $\frac{96}{7}$ is $\frac{1}{32}$.

The 3-adic norm of $3^{-2024} \cdot \frac{8}{13}$ is 3^{2024} .

The p -adic norm $\|q\|_p$ is small if q is divisible by a large power of p .

The p -adic norm of 0 is 0 because 0 is divisible by any power of p .

(p -adic product formula.) If $q \in \mathbb{Q} \setminus \{0\}$, then $|q| \cdot \prod_{p \text{ prime}} \|q\|_p = 1$.

convexness —————

A set X is convex if and only if the line segment joining any two points in X lies within X .

2024-01-10 contraction mappings & product topologies

$\alpha \in I$ henceforth refers to members of a possibly uncountable index set I .

If $\{A_\alpha\}_{\alpha \in I}$ is a family of connected sets in a topological space X such that $\forall \alpha, \alpha' \in I : A_\alpha \cap A_{\alpha'} \neq \emptyset$, then $\bigcup_{\alpha \in I} A_\alpha$ is connected.

Proof. Suppose $A := \bigcup_{\alpha \in I} A_\alpha$ is not connected. Let $U, U' \subseteq X$ be open and nonempty relative to A such that $U \cap U' = \emptyset$ and $A = U \cup U'$. Since each A_α is connected, there is no α for which $A_\alpha \cap U \neq \emptyset \neq A_\alpha \cap U'$. But then $\exists \alpha, \alpha' \in I : A_\alpha \subseteq U \wedge A_{\alpha'} \subseteq U'$, which contradicts that $A_\alpha \cap A_{\alpha'} \neq \emptyset$. \square

total disconnectedness —————

A topological space X is totally disconnected if and only if every connected subset of X is a singleton.

A totally disconnected space in \mathbb{R} contains only points and no intervals.

The Cantor set is a totally disconnected space in \mathbb{R} .

contraction mappings

For a metric space (X, d) , $T : X \rightarrow X$ is a **contraction mapping** if and only if $\exists c \in [0, 1) : \forall x, x' \in X : d(T(x), T(x')) \leq c \cdot d(x, x')$.

All contraction mappings are **continuous**.

(Orbit lemma.) For $x \in X$, the **orbit** $(T^n(x))_{n \in \mathbb{N}}$ of a **contraction mapping** T on X is a **Cauchy sequence**.

Proof. For $n \in \mathbb{Z}_+$,

$$\begin{aligned} d(T^n(x), T^{n+1}(x)) &< c \cdot d(T^{n-1}(x), T^n(x)) \\ &< c^2 \cdot d(T^{n-2}(x), T^{n-1}(x)) \\ &\dots < c^n \cdot d(x, T(x)). \end{aligned}$$

Let $m \geq n$. By the triangle inequality,

$$\begin{aligned} d(T^n(x), T^m(x)) &\leq \sum_{k=n}^{m-1} d(T^k(x), T^{k+1}(x)) \\ &\leq d(x, T(x)) \sum_{k=n}^{m-1} c^k \\ &= c^n \cdot d(x, T(x)) \sum_{k=0}^{m-n-1} c^k \\ &\leq \frac{c^n \cdot d(x, T(x))}{1-c}. \end{aligned}$$

For $\epsilon > 0$, choosing $n > \log_c \left(\frac{\epsilon}{2} \cdot \frac{1-c}{d(x, T(x))} \right)$ and $m, m' \geq n$ guarantees

$$\begin{aligned} d(T(m), T(m')) &\leq \\ d(T(n), T(m)) + d(T(n), T(m')) &< \epsilon. \end{aligned}$$

□

(**Contraction mapping theorem.**) If (X, d) is a **nonempty** and **complete** metric space and T is a **contraction mapping on X** , then

$$\exists! z \in X : T(z) = z,$$

i.e. z is *the unique fixed point*, and

$$\forall x \in X : \lim_{n \rightarrow \infty} T^n(x) = z,$$

where $\forall n \in \mathbb{N} : T^n(x) := T(T(\dots T(x)))$.
 n times

Proof. For $x \in X$, $T(x)$ is Cauchy by the orbit lemma, and since (X, d) is complete it converges to some point $z \in X$. Then z is a fixed point of T because

$$\begin{aligned} T^n(x) \rightarrow z &\implies T(T^n(x)) \rightarrow T(z) \\ &\implies T(T^n(x)) = T^{n+1}(x) \rightarrow z = T(z), \end{aligned}$$

and is its unique fixed point because

$$T(z) = z \wedge T(z') = z' \implies d(z, z') \leq c \cdot d(z, z') \iff d(z, z') = 0. \quad \square$$

$x \mapsto \frac{x}{2}$ is a contraction mapping on $(\mathbb{R}, |\cdot|)$ with fixed point $z = 0$.

iterated function systems and fixed point sets —————

$\mathcal{K}(X)$ henceforth denotes the set of compact subsets of a set X .

If $m \in \mathbb{Z}_+$, $\{T_n : \mathbb{R}^d \rightarrow \mathbb{R}^d\}_{n \in [m]}$ are contraction mappings, and

$$F : (\mathcal{K}(\mathbb{R}^d), d_H) \rightarrow (\mathcal{K}(\mathbb{R}^d), d_H) : A \mapsto \bigcup_{n \in [m]} T_n(A),$$

then $\exists! A \in \mathcal{K}(\mathbb{R}^d) : F(A) = A$. This is the *fixed point set* of F .

Proof. F is a contraction mapping (to be demonstrated in problem set 2), and $(\mathcal{K}(X), d_H)$ is a complete metric space (???). A unique fixed point set thus exists by the contraction mapping theorem. \square

Let $T_0, T_1 : \mathbb{R} \rightarrow \mathbb{R}$ such that $T_0(x) := \frac{x}{3}$ and $T_1(x) := \frac{x+2}{3}$ and $F(A) : \mathcal{K}(\mathbb{R}) \rightarrow \mathcal{K}(\mathbb{R})$ such that $F(A) := T_0(A) \cup T_1(A)$. Then $F([0, 1]) = T_0([0, 1]) \cup T_1([0, 1]) = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$, and the *fixed point set* of F is *the middle-thirds Cantor set*.

If $T_0, T_1 : \mathbb{R}^d \rightarrow \mathbb{R}^d$ are *contraction mappings* and $F : (\mathcal{K}(\mathbb{R}^d), d_H) \rightarrow (\mathcal{K}(\mathbb{R}^d), d_H)$ such that $F(A) = T_0(A) \cup T_1(A)$, then the composition of F with itself is

$$F^2(A) = T_0(T_0(A)) \cup T_0(T_1(A)) \cup T_1(T_0(A)) \cup T_1(T_1(A)).$$

basis for a topology

A **basis** \mathcal{B} for a topology τ is a subset of τ such that $\forall A \in \tau : \exists \{A_\alpha\}_{\alpha \in I} \subseteq \mathcal{B} : A = \bigcup_{\alpha \in I} A_\alpha$.

The open intervals form a basis for the standard topology on \mathbb{R} .

For a metric space X , $\{B(x, \epsilon) \mid x \in X, \epsilon > 0\}$ is a basis for the open sets in X .

If \mathcal{B} is a basis for the topology on X , then $\forall x \in X : \exists U \in \mathcal{B} : x \in U$.

If \mathcal{B} is a basis for the topology τ on X , then $[U, U' \in \mathcal{B} \wedge x \in U \cap U'] \implies \exists U'' \in \tau : x \in U'' \subseteq U \cap U'$.

If $f : X \rightarrow Y$ for topological spaces (X, τ_X) and (Y, τ_Y) and \mathcal{B} is a basis for τ_Y , then f is continuous if and only if $\forall U \in \mathcal{B} : f^{-1}(U) \in \tau_X$.

product topologies

For topological spaces $\{(X_\alpha, \tau_\alpha)\}_{\alpha \in I}$, $\prod_{\alpha \in I} A_\alpha$ is a cylinder set in $X := \prod_{\alpha \in I} X_\alpha$ if and only if $\forall \alpha \in I : A_\alpha \in \tau_\alpha$.

If $X := \mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$, then \forall open intervals $(a, b), (a', b') \subseteq \mathbb{R} : (a, b) \times (a', b')$ is a cylinder set in X .

For topological spaces $\{X_\alpha\}_{\alpha \in I}$, consider a cylinder set $\prod_{\alpha \in I} A_\alpha$ in $X := \prod_{\alpha \in I} X_\alpha$ such that $\exists I' \subseteq I : [|I'| \in \mathbb{N} \wedge \forall \alpha \in I \setminus I' : A_\alpha = X_\alpha]$. These base cylinder sets form a basis for the product topology on X .

projection maps

For a vector $x := (x_\alpha)_{\alpha \in I}$ in possibly uncountable dimensions, the function $\pi_\alpha(x) := x_\alpha$ is a projection map.

For topological spaces $\{X_\alpha\}_{\alpha \in I}$ let $f : Y \rightarrow \prod_{\alpha \in I} X_\alpha$, and for $\alpha \in I$ let $f_\alpha : Y \rightarrow X_\alpha : y \mapsto \pi_\alpha(f(y))$. Then f is continuous if and only if $\forall \alpha \in I : f_\alpha$ is continuous.

Proof. Suppose f is continuous. Then, for $\alpha' \in I$ and an open set $U \subseteq X_{\alpha'}$,

$$f_{\alpha'}^{-1}(U) = f^{-1}\left(U \times \prod_{\alpha \in I \setminus \{\alpha'\}} X_\alpha\right)$$

is open as the preimage of a base cylinder set in $\prod_{\alpha \in I} X_\alpha$.

Suppose $\forall \alpha \in I : f_\alpha$ is continuous. It suffices to verify that the preimage of f for a base cylinder set U is open. Let $I' \subseteq I$ be a finite index subset for which

$$U = \prod_{\alpha \in I'} A_\alpha \times \prod_{\alpha \in I \setminus I'} X_\alpha.$$

Then

$$\begin{aligned} f^{-1}(U) &= \left(\bigcap_{\alpha \in I'} f_\alpha^{-1}(A_\alpha) \right) \cap \left(\bigcap_{\alpha \in I \setminus I'} f_\alpha^{-1}(X_\alpha) \right) \\ &= \bigcap_{\alpha \in I'} f_\alpha^{-1}(A_\alpha) \end{aligned}$$

is open in Y as the finite intersection of open sets in Y . □

2024-01-12 Hilbert spaces & Hausdorff spaces

inner product spaces

The closed unit ball in l_p is not **compact** because it contains $(x_n)_{n \in \mathbb{N}}$ where

$$\forall n, k \in \mathbb{N} : \pi_k(x_n) = \begin{cases} 1 & k = n \\ 0 & k \neq n \end{cases},$$

which has no convergent subsequence.

An **inner product space** V is a **normed vector space** with an **inner product operation** $\langle v, v' \rangle$ for $v, v' \in V$ that induces its **norm** $\|v\|^2 =: \langle v, v \rangle$.

(**Parallelogram law**.) The **norm** in a vector space V is induced by an **inner product** if and only if

$$\forall v, v' \in V : \|v + v'\|^2 + \|v - v'\|^2 = 2(\|v\|^2 + \|v'\|^2)$$

holds.

(**Hanner's inequality**.) For a measure space X , $p \in [1, \infty) \setminus \{2\}$, and $f, g \in L^p(X)$,

$$\|f + g\|_p^p + \|f - g\|_p^p * (\|f\|_p + \|g\|_p)^p + \left| \|f\|_p - \|g\|_p \right|^p$$

where $*$:= $\begin{cases} \geq & p < 2 \\ \leq & p > 2 \end{cases}$. If $p = 2$, then this becomes **the parallelogram law**.

Hilbert spaces

A Hilbert space is a complete inner product space.

l_2 is a Hilbert space with $\langle (a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \rangle = \sum_{n \in \mathbb{N}} |a_n \overline{b_n}|$.

For a measure space X , $L^2(X)$ is a Hilbert space with $\langle f, g \rangle := \int_X f(x) \overline{g(x)} dx$.

(Parseval's identity.) For $f \in L^2([0, 2\pi])$, which has a basis $\{e^{inx}\}_{n \in \mathbb{Z}}$,

$$\begin{aligned} \|f\|_2^2 &= \int_0^{2\pi} |f(x)|^2 dx \\ &= 2\pi \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2 = \langle f, f \rangle \end{aligned}$$

where $\hat{f}(n) = \langle f, e^{inx} \rangle = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-inx} dx$.

(Consequence of Riesz's lemma.) The closed unit ball in a Hilbert space V is compact if and only if V is finite-dimensional.

compact sets & continuity

Continuous functions map compact sets to compact sets.

Proof. Let f be continuous and A a compact set in its domain. The preimages of an open cover U of $f(A)$ form an open cover U' of A . Then a finite subcover of U' exists, and the images of sets in U' form a finite subcover of U . \square

(Tychonoff.) If $\forall \alpha \in I : X_\alpha$ is compact, then $\prod_{\alpha \in I} X_\alpha$ is compact in the product topology.

For topological spaces X and Y , a bijection $f : X \rightarrow Y$ is a homeomorphism if and only if f and f^{-1} are both continuous.

For a subset A of a metric space (X, d) and $x \in X$, $d(x, A) := \inf_{x' \in A} d(x, x')$ is continuous.

Proof. Let $\epsilon, \delta > 0$. Then $\exists a \in A : d(x, a) < \delta$, so for $x' \in B(x, \epsilon)$

$$\begin{aligned} d(x', A) &\leq d(x', a) \\ &\leq d(x, a) + d(x, x') \\ &< d(x, A) + \delta + \epsilon. \end{aligned}$$

Taking the limit as $\delta \rightarrow 0$ gives $d(x', A) \leq d(x, A) + \epsilon$. By parallel reasoning, $d(x, A) \leq d(x', A) + \epsilon$, hence $|d(x, A) - d(x', A)| \leq \epsilon$. \square

Hausdorff spaces

A topological space (X, τ) is a **Hausdorff space** if and only if $\forall x \neq x' \in X : \exists U, U' \in \tau : [U \cap U' = \emptyset \wedge (x, x') \in U \times U']$.

All **metric spaces** are Hausdorff spaces.

*Singletons in a **Hausdorff space** X with topology τ are **closed**.*

Proof. Let $x' \in X$. Then

$$\forall x \in X : \exists U_x \in \tau : [x \in U_x \wedge x' \notin U_x] \implies X \setminus \{x'\} = \bigcup_{x' \in X \setminus \{x'\}} U_x \in \tau. \quad \square$$

$\tau := \{\emptyset, X\}$ is the **trivial topology** on X .

If τ is the **trivial topology** and $|X| > 1$, then (X, τ) is **not a Hausdorff space**.

The **subspace topology** on a subspace Y of a topological space (X, τ) is $\tau_Y := \{U \cap Y \mid U \in \tau\}$.

Any subspace of a **Hausdorff space** is a **Hausdorff space** under the **subspace topology**.

*For Hausdorff spaces (X, τ) and (X', τ') , $X \times X'$ under the product topology is a **Hausdorff space**.*

Proof. Let \mathcal{B} be the cylinder set basis for the product topology and let $(x_0, x'_0) \neq (x_1, x'_1) \in X \times Y$, assuming without loss of generality that $x_0 \neq x_1$. Then

$$\exists U_0, U_1 \in \tau : [U_0 \cap U_1 = \emptyset \wedge (x_0, x_1) \in U_0 \times U_1],$$

so $U_0 \times X'$ and $U_1 \times X'$ are disjoint open sets that contain (x_0, x'_0) and (x_1, x'_1) respectively. \square

*Any **compact** subspace of a Hausdorff space is **closed**.*

Proof. Let (X, τ) be a Hausdorff space, $A \subseteq X$ a compact subset, and $x' \in X \setminus A$. For $x \in A$, let $U_x, U'_x \in \tau$ such that $U_x \cap U'_x = \emptyset$ and $(x, x') \in U_x \times U'_x$.

Since $\{U_x\}_{x \in A}$ is an open cover of A , let $\{x_n\}_{n \in [m]} \subseteq A$ such that $\{U_{x_n}\}_{n \in [m]}$ covers A . Then $x' \in \bigcap_{n \in [m]} U'_{x_n} \subseteq X \setminus A$, and this intersection is open and disjoint to $\bigcup_{n \in [m]} U_{x_n}$. Thus $X \setminus A$ is open. \square

A *continuous bijection* $f : X \rightarrow Y$ is a *homeomorphism* if X is *compact* and Y is a *Hausdorff space*.

Proof. Let $A \subseteq X$ be closed. Then A is compact, so $(f^{-1})^{-1}(A) = f(A)$ is closed as a compact subset of a Hausdorff space. It follows that f^{-1} is continuous. \square

normal spaces

A Hausdorff space (X, τ) is a *normal space* if and only if

$$\exists U, U' \in \tau : [U \cap U' \neq \emptyset \wedge A \subseteq U \wedge A' \subseteq U']$$

for any *disjoint closed sets* $A, A' \subseteq X$.

All *compact Hausdorff space* are *normal spaces*.

Proof. Let (X, τ) be a compact Hausdorff space and $A, A' \subseteq X$ closed sets with $A \cap A' = \emptyset$. For $x \in A$, let $U_x, U'_x \in \tau$ such that $U_x \cap U'_x = \emptyset$, $x \in U_x$, and $A' \subseteq U'_x$.

Since $\{U_x\}_{x \in A}$ is an open cover of A and A is compact as the closed subset of a compact set, let $\{x_n\}_{n \in [m]} \subseteq A$ such that $\{U_{x_n}\}_{n \in [m]}$ covers A . It follows that $\bigcup_{n \in [m]} U_{x_n}$ and $\bigcap_{n \in [m]} U'_{x_n} \supseteq A'$ are open and disjoint. \square

All *metric spaces* are *normal spaces*.

Proof. Let $A, A' \subseteq X$ be disjoint and closed. Then

$$U_A := \{x \mid d(x, A) < d(x, A')\}, \quad U'_A := \{x \mid d(x, A') < d(x, A)\},$$

are disjoint, and they are respective open supersets of A and A' because $x \in A \implies d(x, A) = 0 < d(x, A')$ (and vice versa) and

$$U_A = f^{-1}((-\infty, 0)), \quad U'_A = f^{-1}((0, \infty))$$

where $f : x \mapsto d(x, A) - d(x, A')$ is continuous as the difference between continuous functions. \square

Lebesgue number

A *Lebesgue number* of an open cover $\{A_\alpha\}_{\alpha \in I}$ of X is $\epsilon > 0$ such that $\forall x \in X : \exists \alpha \in I : B(x, \epsilon) \subseteq A_\alpha$.

Any open cover of *a compact metric space* has *a Lebesgue number*.

Proof. Let X be a compact metric space and $\{U_\alpha\}_{\alpha \in I}$ an open cover of X with finite subcover $\{U_{\alpha_n}\}_{n \in [m]}$. For $n \in [m]$ and $x \in X$, let

$$d_n(x) := d(x, X \setminus U_n),$$

noting that $B(x, \epsilon) \subseteq U_n \iff d_n(x) \geq \epsilon$ and that $\exists n \in [m] : x \in U_n \implies d_n(x) > 0$.

Let $f(x) := \max_{n \in [m]} d_n(x)$, which is continuous as the maximum of continuous functions. Then $f(X) \subseteq (0, \infty)$ is compact hence closed and bounded. It follows that $\exists \delta > 0 : f(X) \subseteq [\delta, \infty)$ and δ is a Lebesgue number for $\{U_\alpha\}_{\alpha \in I}$. \square

If $f : X \rightarrow Y$ is *continuous*, X and Y are metric spaces, and X is *compact*, then f is *uniformly continuous*.

Proof. For $\epsilon > 0$, let δ be a Lebesgue number for $\{f^{-1}(B(y, \epsilon))\}_{y \in Y}$. Then $\forall x \in X : \exists y \in Y : f(B(x, \delta)) \subseteq B(y, \epsilon)$, so for $x' \in B(x, \delta)$

$$\begin{aligned} d(f(x), f(x')) &\leq \\ d(f(x), y) + d(f(x'), y) &< 2\epsilon. \end{aligned}$$

\square

2024-01-17 product spaces & separable spaces

metrizability

A topological space (X, τ) is *metrizable* if and only if *there exists a metric d on X that induces τ* . In other words, *the set of open balls with respect to d is a basis for τ* .

Any topological space that is not *normal* is non-metrizable.

If $\forall n \in \mathbb{N} : (X_n, d_n)$ are *metric spaces*, then $X := \prod_{n \in \mathbb{N}} X_n$ is *metrizable under the product topology*.

Proof. For $n \in \mathbb{N}$ and $x_n, y_n \in X_n$ let $\tilde{d}(x_n, y_n) := \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \leq 1$, noting that this is an equivalent metric to d_n . Then define

$$D : X \rightarrow \mathbb{R} : (x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \mapsto \sum_{n \in \mathbb{N}} \left(\frac{\tilde{d}_n(x_n, y_n)}{2^n} \right).$$

As shown in problem set 1, D induces the product topology on X . \square

countable product of compact metric spaces

$\tau := \mathcal{P}(X)$ is the discrete topology on X .

If τ is the discrete topology and $\forall n \in \mathbb{N} : X_n := (\{0, 1\}, \tau)$, then $\prod_{n \in \mathbb{N}} X_n$ is not compact under the discrete topology because it is an open cover of itself but has no finite subcover.

(Tychonoff for countable products.) If $\forall n \in \mathbb{N} : (X_n, \tau_n)$ is a compact topological space, then $X := \prod_{n \in \mathbb{N}} X_n$ is compact under the product topology.

Proof. Suppose X is not compact and let $C := \{C_\alpha\}_{\alpha \in I}$ be an open cover of X with no finite subcover.

Suppose for every $x_0 \in X_0$ that there is a set $U_{x_0} \in \tau_0$ such that

$$x_0 \in U_{x_0} \wedge \exists I' \subseteq I : |I'| \in \mathbb{N} \wedge \left(U_{x_0} \times \prod_{n \in \mathbb{Z}_+} X_n \right) \subseteq \bigcup_{\alpha \in I'} C_\alpha.$$

Then

$$X \subseteq \bigcup_{x_0 \in X_0} \left(U_{x_0} \times \prod_{n \in \mathbb{Z}_+} X_n \right).$$

Since X_0 is compact, this open cover has a finite subcover and contradicts that C has no finite subcover. Thus there is some $x_0 \in X_0$ for which no such U_{x_0} exists.

By similar reasoning, there exists $x_1 \in X_1$ such that no cylinder set of the form $U_{x_0} \times U_{x_1} \times \prod_{n \in \mathbb{Z}_+ \setminus \{1\}} X_n$ both satisfies $(x_0, x_1) \in U_{x_0} \times U_{x_1}$ and has a finite open cover in C ; otherwise,

$$\left(U_{x_0} \times \prod_{n \in \mathbb{Z}_+} X_n \right) \subseteq \bigcup_{x_1 \in X_1} \left(U_{x_0} \times U_{x_1} \times \prod_{n \in \mathbb{Z}_+ \setminus \{1\}} X_n \right)$$

would have a finite subcover by compactness of X_1 and thus $(U_{x_0} \times \prod_{n \in \mathbb{Z}_+} X_n)$ would have a finite open cover in C .

Selecting $x_n \in X_n$ in this fashion for every $n \in \mathbb{N}$ provides a point $x := (x_n)_{n \in \mathbb{N}} \in X$ such that, for any $m \in \mathbb{N}$, every cylinder set of the form

$$\prod_{n=0}^m U_n \times \prod_{n=m+1}^{\infty} X_n,$$

where $(x_n)_{n=0}^m \in \prod_{n=0}^m U_n$, has no finite open cover in C . But this cannot be; $\exists C_\alpha \in C : x \in C_\alpha$, implying that x lies in some cylinder set $V := \prod_{n \in \mathbb{N}} V_n \subseteq C_\alpha$ in

the product topology basis. Taking m to be the maximum index n for which $V_n \neq X_n$ gives

$$(x_n)_{n \in \mathbb{N}} \in \left(\prod_{n=0}^m V_n \times \prod_{n=m+1}^{\infty} X_n \right) \subseteq C_\alpha,$$

which is certainly a finite cover. □

separability

A metric space is **separable** if and only if it has a countable dense subset.

All **compact** metric spaces are separable.

Proof. Let X be a compact metric space. For $n \in \mathbb{Z}_+$, $X \subseteq \bigcup_{x \in X} B(x, \frac{1}{n})$, and by compactness, let $N_n \subseteq X$ be a finite subset such that $X \subseteq \bigcup_{x \in N_n} B(x, \frac{1}{n})$. Thus $\bigcup_{n \in \mathbb{Z}_+} N_n \subseteq X$ is dense and countable as the countable union of finite sets. □

A metric space X with topology τ is **separable** if and only if $\exists \{U_n\}_{n \in \mathbb{N}} \subseteq \tau : \forall U \in \tau : \exists N \subseteq \mathbb{N} : U = \bigcup_{n \in N} U_n$; in other words, τ has a countable basis.

Any **subspace** of a separable metric space is **separable**.

\mathbb{R}^n , $C([a, b])$, and $L^p([a, b])$ for $p \in [1, \infty)$ are examples of separable spaces.

$L^\infty([a, b])$ is an example of a **non-separable** space.