



## Biochemical and Biophysical Research Communications

Volume 257, Issue 3, 21 April 1999, Pages 708-713

Regular Article

# Fabry Disease: Identification of Novel Alpha-Galactosidase A Mutations and Molecular Carrier Detection by Use of Fluorescent Chemical Cleavage of Mismatches ☆

Dominique P. Germain<sup>1</sup>, Livia Poenaru

[Show more](#) ✓



Share



Cite

<https://doi.org/10.1006/bbrc.1999.0310> ↗

[Get rights and content](#) ↗

## Abstract

Fabry disease (FD) (angiokeratoma corporis diffusum) is an X-linked inborn error of glycosphingolipid metabolism caused by defects in the lysosomal  $\alpha$ -galactosidase A gene (*GLA*). The enzymatic defect leads to the systemic accumulation of neutral glycosphingolipids with terminal  $\alpha$ -galactosyl moieties. Clinically, affected hemizygous males have angiokeratoma, severe acroparesthesia, renal failure, and vasculopathy of the heart and brain. While demonstration of  $\alpha$ -galactosidase deficiency in leukocytes is diagnostic in affected males, enzymatic detection of female carriers is often inconclusive, due to random X-chromosomal inactivation, underlining the need of molecular investigations for accurate genetic counseling. By use of chemical cleavage of mismatches adapted to fluorescence-based detection systems, we have characterized the mutations underlying  $\alpha$ -Gal A deficiency in 16 individuals from six unrelated families with FD. The mutational spectrum included five missense mutations (C202W, C223G, N224D, R301Q, and

Q327K) and one splice-site mutation [IVS3 G(-1) → C]. Studies at the mRNA level showed that the latter led to altered pre-mRNA splicing with consequent alteration of the mRNA translational reading frame and generation of a premature termination codon of translation. By use of this strategy, carrier status was accurately assessed in all seven at-risk females tested, whereas enzymatic dosages failed to diagnose or exclude heterozygosity.

[Recommended articles](#)

---

## References (23)

T. Ohshima *et al.*

Gene (1995)

R.W. Gotlib *et al.*

Biochem. Mol. Med. (1996)

R.G.H. Cotton

Trends Genet. (1997)

R.J. Desnick *et al.*

The Metabolic and Molecular Bases of Inherited Diseases(1995)

H. Sakuraba *et al.*

Am. J. Hum. Genet. (1990)

W. Scheidt *et al.*

N. Engl. J. Med. (1991)

R. Desnick *et al.*

J. Lab. Clin. Med. (1973)

D.F. Bishop *et al.*

Proc. Natl. Acad. Sci USA (1986)

R. Kornreich *et al.*

Nucleic Acids Res. (1989)

C.M. Eng *et al.*

Am. J. Hum. Genet. (1993)

[View more references](#)

---

## Cited by (49)

### [Screening for Fabry disease in male patients with end-stage renal disease in western France](#)

2021, Nephrologie et Therapeutique

*Citation Excerpt :*

...Fabry disease (FD; OMIM #301500) is a rare X-linked genetic disease due to pathogenic variants in the GLA gene coding for the lysosomal  $\alpha$ -galactosidase A [1–3]....

[Show abstract](#) ✓

### [Standardising clinical outcomes measures for adult clinical trials in Fabry disease: A global Delphi consensus](#)

2021, Molecular Genetics and Metabolism

*Citation Excerpt :*

...Two clinical phenotypes have been described: “classical” and “later-onset” [3]. These cannot be predicted exclusively by genotype [4,5]. Classical FD is characterised by a multi-system involvement with early acroparaesthesia, sweating abnormalities and gastrointestinal disturbance, whereas most of the later onset phenotypes may be an attenuated form of the disease, commonly involving a single organ system, usually cardiac or sometimes renal [3]....

[Show abstract](#) ✓

### [Why systematic literature reviews in Fabry disease should include all published evidence](#)

2019, European Journal of Medical Genetics

*Citation Excerpt :*

...The X-chromosome inactivation pattern that is observed in female patients with Fabry disease correlates with the variation in phenotype and disease course (Echevarria et al., 2016). Fabry disease is further characterised by a large number of pathogenic variants in the GLA gene (Germain et al., 1999), including variants associated with the classic presentation of the disease, later onset or atypical disease presentations (Germain, 2001; Germain et al., 1996, 2018) and of uncertain significance. Thus,

pooling results from hemizygous male patients and heterozygous female patients is another source of bias in study data (Dobrovolny et al., 2005; Echevarria et al., 2016; Germain, 2007)....

[Show abstract](#) ✓

## Fabry disease: Four case reports of meningioma and a review of the literature on other malignancies

2017, Molecular Genetics and Metabolism Reports

*Citation Excerpt :*

...However, little has been reported on the occurrence of meningioma in association with metabolic disorders. These are the first reported cases of meningioma occurring in Fabry disease, an X-linked metabolic disorder caused by a deficiency of lysosomal alpha-galactosidase, resulting in cellular accumulation of the lipid globotriaosylceramide (GL-3) and its deacylated product lyso-Gb3 [6,7]. We review the existing reports of cancer occurring in patients with Fabry disease, and discuss the possible association of metabolic lipid disorders with respect to the evolution of malignancy....

[Show abstract](#) ✓

## The alpha-galactosidase A p.Arg118Cys variant does not cause a Fabry disease phenotype: Data from individual patients and family studies

2015, Molecular Genetics and Metabolism

[Show abstract](#) ✓

## An intronic haplotype in $\alpha$ galactosidase A is associated with reduced mRNA expression in males with cryptogenic stroke

2014, Gene

*Citation Excerpt :*

...When intronic variants are involved, one might entertain the possibility that these induce altered splice sites and skipping of whole exons and thereby may be predictive of disease expression. Studies at the mRNA level have shown that intronic variants induce altered splicing with consequent alteration of the mRNA translational reading frame (Germain and Poenaru, 1999); using this strategy, female patients were identified despite the presence of residual enzyme (Germain and Poenaru, 1999). On the other hand, none of the intronic variants studied here appeared to aberrantly affect transcript splicing....

[Show abstract](#) ✓



View all citing articles on Scopus ↗

---

☆ Sriver, C. R.Beaudet, A. L.Sly, W. S.Valle, D.

- 1 To whom correspondence should be addressed at Laboratoire de Génétique. CHU Broussais, 96, rue Didot. 75014 Paris, France. Fax: +33 1 45 41 02 34. E-mail:dominique.germain@brs.ap-hop-paris.fr.

View full text

Copyright © 1999 Academic Press. All rights reserved.



All content on this site: Copyright © 2024 Elsevier B.V., its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply.

