

Changing Primaries

This appendix describes the operation of transforming one set of primaries into another. The mathematical name for this operation is a *change of basis*.

To convert a color from one set of primary lights to another, it is first necessary to define a conversion between the primaries themselves. We can think of this as matching each of the new primary lights using the old primary system. Suppose we designate our original set of primaries P_1 , P_2 , and P_3 and the new set of primaries Q_1 , Q_2 , and Q_3 . We now use our original primaries to create matches with each of the new primaries in turn. Let us call the amount of each of the P primaries c_{ij} .

Thus,

$$\begin{aligned} Q_1 &\equiv c_{11}P_1 + c_{12}P_2 + c_{13}P_3 \\ Q_2 &\equiv c_{21}P_1 + c_{22}P_2 + c_{23}P_3 \\ Q_3 &\equiv c_{31}P_1 + c_{32}P_2 + c_{33}P_3 \end{aligned} \tag{A.1}$$

If we denote the matrix of c_{ij} values C , then

$$P = CQ \tag{A.2}$$

To reverse the transformation, invert the matrix:

$$P \equiv C^{-1}Q \tag{A.3}$$

This same matrix can now be used to convert any set of values expressed in one set of primaries to the other set of primaries. Thus, the values p_1 , p_2 , and p_3 represent the amounts of the lights in primary system P needed to make a match.

$$\text{Sample} \equiv p_1 P_1 + p_2 P_2 + p_3 P_3 \quad (\text{A.4})$$

Then we can calculate the values q in primary system Q simply by solving

$$q = Cp \quad (\text{A.5})$$